



# Measure of non-compactness and limiting interpolation with slowly varying functions

Fernando Cobos<sup>1</sup>  · Luz M. Fernández-Cabrera<sup>2</sup> · Manvi Grover<sup>3</sup>

Received: 20 October 2023 / Accepted: 9 February 2024  
© The Author(s) 2024

## Abstract

We give estimates for the measure of non-compactness of an operator interpolated by the limiting methods involving slowly varying functions. As applications we establish estimates for the measure of non-compactness of operators acting between Lorentz–Karamata spaces.

**Keywords** Limiting interpolation · Slowly varying functions · Measure of non-compactness · Lorentz–Karamata spaces · Generalized Lorentz–Zygmund spaces

**Mathematics Subject Classification** 46M35 · 46E30 · 47B10

## 1 Introduction

The real interpolation method  $(A_0, A_1)_{\theta, q}$  has found important applications in Operator Theory, Approximation Theory, Function Spaces and Harmonic Analysis. See, for

---

To Professor William B. Johnson on the occasion of his 80th birthday.

---

Communicated by Mieczyslaw Mastylo.

---

✉ Fernando Cobos  
cobos@mat.ucm.es

Luz M. Fernández-Cabrera  
luz\_fernandez-c@mat.ucm.es

Manvi Grover  
grover.manvi94@gmail.com

- <sup>1</sup> Departamento de Análisis Matemático y Matemática Aplicada, Facultad de Matemáticas, Universidad Complutense de Madrid, Plaza de Ciencias 3, 28040 Madrid, Spain
- <sup>2</sup> Sección Departamental del Departamento de Análisis Matemático y Matemática Aplicada, Facultad de Estudios Estadísticos, Universidad Complutense de Madrid, 28040 Madrid, Spain
- <sup>3</sup> Department of Mathematical Analysis, Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic

example, the monographs by Butzer and Berens [8], Bergh and Löfström [4], Triebel [43, 44], König [32] and Bennett and Sharpley [3]. The real method is very flexible, admitting several equivalent definitions, what is very useful in applications.

The real method applied to the couple of Lebesgue spaces  $(L_1, L_\infty)$  yields Lorentz spaces  $L_{p,q}$ . It is possible to obtain more general spaces if we modify the definition of the real method. So, logarithmic perturbations of the real method produce Lorentz–Zygmund spaces  $L_{p,q}(\log L)_a$  (see [20, 27, 28]) and perturbations involving slowly varying functions  $(L_1, L_\infty)_{\theta,q;b}$  give Lorentz–Karamata spaces  $L_{p,q;b}$  (see [29]).

We are interested here in the limit cases when  $\theta = 0, 1$  of the perturbations with slowly varying functions  $(A_0, A_1)_{\theta,q;b}$ . These spaces are very close to  $A_0$  when  $\theta = 0$  and to  $A_1$  when  $\theta = 1$ . They have received attention from a number of authors either to study limiting embeddings between function spaces or to establish limiting properties of operators (see, for example, [26, 29, 36]).

Among the classical problems for any interpolation method, a prominent one is to describe the behavior of properties that operators may have. First of all boundedness but then other useful properties of operators. For example, techniques used by Davis, Figiel, Johnson and Pelczyński [23] in the proof of their famous factorization theorem for weakly compact operators motivated the investigation on the behavior of weak compactness under interpolation (see, for example, [1, 18, 31, 34, 35]).

The behavior under interpolation of compactness have been also deeply studied (see [9, 17, 22] and the references given there). Quantitative estimates in terms of the measure of non-compactness have been also established. Concerning the real method, the first result in this direction is due to Edmunds and Teixeira [42]. They assume an approximation condition for the couple in the target. The case of general Banach couples has been studied by Cobos, Fernández-Martínez and Martínez [15]. Results for the real method with a function parameter and  $0 < \theta < 1$  are due to Cordeiro [21], Szwedek [41] and Cobos, Fernández-Cabrera and Martínez [11]. Besides, the case of limiting methods involving logarithms have been considered by Cobos, Fernández-Cabrera and Martínez [12, 14] and Besoy and Cobos [5].

Our aim here is to establish estimates for the measure of non-compactness of operators interpolated by the limiting perturbations of the real method involving slowly varying functions. As applications we derive estimates for the measure of non-compactness of operators acting between certain Lorentz–Karamata spaces. In particular, one of our results can be considered as a quantitative extension of a compactness result of Edmunds and Opic [26] for operators acting between Lorentz–Zygmund spaces.

We work with quasi-Banach couples  $(A_0, A_1)$ . Our techniques are based on the vector-valued sequence spaces that come up with the definition of  $(A_0, A_1)_{\theta,q;b}$  and with its description as a  $J$ -space. These ideas originated in the papers on compactness by Cobos and Peetre [19] and Cobos, Kühn and Schonbek [17]. In the context of the measure of non-compactness, they were developed by Cobos, Fernández-Martínez and Martínez [15], Cobos, Fernández-Cabrera and Martínez [14] and Besoy and Cobos [5] among other authors.

## 2 Limiting real interpolation spaces

Let  $(A, \|\cdot\|_A)$  be a quasi-Banach space and let  $c_A \geq 1$  be its constant in the quasi-triangle inequality. Let  $0 < p \leq 1$  such that  $c_A = 2^{1/p-1}$ . According to the Aoki-Rolewicz theorem (see [33, Section 15.10]) there is another quasi-norm  $|||\cdot|||$  on  $A$  which is equivalent to  $\|\cdot\|_A$  and such that  $|||\cdot|||^p$  satisfies the triangle inequality. Then  $(A, |||\cdot|||)$  is called a  $p$ -Banach space. Note that if  $0 < r < p$ , then  $(A, |||\cdot|||)$  is also an  $r$ -Banach space and that any  $p$ -Banach space satisfies the quasi-triangle inequality with constant  $2^{1/p-1}$ .

If  $B$  is another quasi-Banach space, we write  $A = B$  if  $A \hookrightarrow B$  and  $B \hookrightarrow A$ , where  $\hookrightarrow$  means continuous embedding.

For  $0 < q \leq \infty$ , let  $\ell_q$  be the space of  $q$ -summable sequences with  $\mathbb{Z}$  as index set. If  $(w_m)_{m \in \mathbb{Z}}$  is a sequence of positive numbers, we denote by  $\ell_q(w_m)$  the space of all scalars sequences  $(\xi_m)$  such that  $(w_m \xi_m) \in \ell_q$ .

Let  $(W_m)$  be a sequence of quasi-Banach spaces with the same constant in the quasi-triangle inequality. We put

$$\begin{aligned} \ell_q(w_m W_m) &= \{w = (w_m) : w_m \in W_m \text{ and} \\ &\|w\|_{\ell_q(w_m W_m)} = \|(w_m \|w_m\|_{W_m})\|_{\ell_q} < \infty\}. \end{aligned}$$

A quasi-Banach space  $(\Gamma, \|\cdot\|_\Gamma)$  of real valued sequences with  $\Gamma \hookrightarrow \ell_q + \ell_q(2^{-m})$  is said to be a *quasi-Banach sequence lattice* if  $\Gamma$  contains all the sequences with only finitely many non-zero coordinates and whenever  $(\eta_m) \in \Gamma$  and  $|\xi_m| \leq |\eta_m|$  for each  $m \in \mathbb{Z}$ , then  $(\xi_m) \in \Gamma$  and  $\|(\xi_m)\|_\Gamma \leq \|(\eta_m)\|_\Gamma$ .

We define  $\Gamma(W_m)$  as the collection of all sequences  $w = (w_m)$  such that  $w_m \in W_m$  and  $\|w\|_{\Gamma(W_m)} = \|(\|w_m\|_{W_m})\|_\Gamma < \infty$ .

Subsequently, if  $b$  and  $v$  are non-negative functions on  $(0, \infty)$ , we say that  $b$  and  $v$  are *equivalent* (and write  $b(t) \approx v(t)$ ) if there are positive constants  $c, C$  such that  $cb(t) \leq v(t) \leq Cb(t)$  for any  $t > 0$ .

A positive, finite and Lebesgue-measurable function  $b$  on  $(0, \infty)$  is said to be *slowly varying* ( $b \in SV(0, \infty)$ ) if, for each  $\varepsilon > 0$ ,  $t^\varepsilon b(t)$  is equivalent to a positive non-decreasing measurable function and  $t^{-\varepsilon} b(t)$  is equivalent to a positive non-increasing measurable function. Important examples of slowly varying functions are powers of iterated logarithms and broken logarithmic functions  $v(t) = \ell^\mathbb{A}(t)$  where  $\ell(t) = (1 + |\log(t)|)$ ,  $\mathbb{A} = (\alpha_0, \alpha_\infty) \in \mathbb{R}^2$ ,  $\ell^\mathbb{A}(t) = \ell^{\alpha_0}(t)$  if  $0 < t \leq 1$  and  $\ell^\mathbb{A}(t) = \ell^{\alpha_\infty}(t)$  if  $1 < t < \infty$ .

We refer to [29] for properties of slowly varying functions. We only recall here that if  $\varepsilon > 0$ , then there are positive constant  $c_\varepsilon, C_\varepsilon$  such that

$$c_\varepsilon \min\{s^{-\varepsilon}, s^\varepsilon\}b(t) \leq b(st) \leq C_\varepsilon \max\{s^\varepsilon, s^{-\varepsilon}\}b(t) \text{ for every } s, t > 0, \quad (2.1)$$

(see [29, Proposition 2.2]). Put

$$\bar{b}(s) = \sup_{t>0} \frac{b(st)}{b(t)}.$$

The function  $\bar{b}$  satisfies that  $\bar{b}(st) \leq \bar{b}(s)\bar{b}(t)$ . Moreover, using (2.1) with  $\varepsilon = 1/2$ , we have

$$s\bar{b}(s) \leq C_{1/2}s^{1/2} \rightarrow 0 \text{ as } s \rightarrow 0. \tag{2.2}$$

Another consequence of (2.1), this time with  $\varepsilon = 1$ , is that

$$c_1/2 \leq \bar{b}(s) \leq 2C_1 \text{ for any } 1/2 \leq s \leq 1. \tag{2.3}$$

Let  $\mathbb{A} = (\alpha_0, \alpha_\infty) \in \mathbb{R}^2$ ,  $v(t) = \ell^{\mathbb{A}}(t)$  and  $\mathbb{B} = (\alpha_0^+ + (-\alpha_\infty)^+, \alpha_\infty^+ + (-\alpha_0)^+)$  with  $\alpha^+ = \max\{0, \alpha\}$ . It follows from [14, Lemma 2.1] and [5, (2.6)] that  $\bar{v}(s) \leq \ell^{\mathbb{B}}(s)$ ,  $s \in (0, \infty)$ .

For  $0 < q \leq \infty$  and  $b \in SV(0, \infty)$ , the quasi-Banach sequence space  $\ell_q(b(2^m))$  will be of special interest for us.

If  $k \in \mathbb{Z}$ , the *shift operator*  $\tau_k$  is defined by  $\tau_k\xi = (\xi_{m+k})_{m \in \mathbb{Z}}$  for  $\xi = (\xi_m)$ . We have

$$\|\tau_k\xi\|_{\ell_q(b(2^m))} = \|(b(2^m)|\xi_{m+k}|)\|_{\ell_q} \leq \bar{b}(2^{-k})\|(b(2^{m+k})|\xi_{m+k}|)\|_{\ell_q}.$$

Hence  $\tau_k : \ell_q(b(2^m)) \rightarrow \ell_q(b(2^m))$  is bounded with

$$\|\tau_k\|_{\ell_q(b(2^m)), \ell_q(b(2^m))} \leq \bar{b}(2^{-k}). \tag{2.4}$$

We say that  $\bar{A} = (A_0, A_1)$  is a (*p-Banach*) *quasi-Banach couple* if  $A_0$  and  $A_1$  are (*p-Banach*) quasi-Banach spaces which are continuously embedded in the same Hausdorff topologic vector space.

For  $t > 0$  and  $a \in A_0 + A_1$ , the Peetre’s *K-functional* is given by

$$K(t, a) = K(t, a; A_0, A_1) = \inf\{\|a_0\|_{A_0} + t\|a_1\|_{A_1} : a = a_0 + a_1, a_j \in A_j\}.$$

If  $a \in A_0 \cap A_1$ , the *J-functional* of Peetre is

$$J(t, a) = J(t, a; A_0, A_1) = \max\{\|a\|_{A_0}, t\|a\|_{A_1}\}.$$

Note that  $K(1, \cdot)$  and  $J(1, \cdot)$  are the quasi-norms of  $A_0 + A_1$  and  $A_0 \cap A_1$ , respectively.

If  $(A_j, \|\cdot\|_{A_j})$  is a *p-Banach* space for  $j = 0, 1$ , then  $J(t, \cdot)$  is also a *p-norm*, as well as

$$\begin{aligned} K_p(t, a) &= K_p(t, a; A_0, A_1) \\ &= \inf\{(\|a_0\|_{A_0}^p + t^p\|a_1\|_{A_1}^p)^{1/p} : a = a_0 + a_1, a_j \in A_j\}. \end{aligned}$$

This last functional is equivalent to the *K-functional*. In fact

$$K(t, a) \leq K_p(t, a) \leq 2^{1/p-1}K(t, a), \quad a \in A_0 + A_1. \tag{2.5}$$

Note that if  $\xi = (\xi_m) \in \ell_p + \ell_p(2^{-m})$  then

$$K_p(2^r, \xi; \ell_p, \ell_p(2^{-m})) = \left( \sum_{m=-\infty}^{\infty} [\min\{1, 2^{r-m}\} |\xi_m|^p] \right)^{1/p}.$$

This expression will be useful later.

A quasi-Banach space  $A$  is said to be an *intermediate space* with respect to the couple  $\bar{A}$  if  $A_0 \cap A_1 \hookrightarrow A \hookrightarrow A_0 + A_1$ . We write  $A^\circ$  for the closure of  $A_0 \cap A_1$  in  $A$ . The fundamental lemma (see [4, Lemma 3.3.2] and [37, Lemma 2.4]) yields that

$$a \in (A_0 + A_1)^\circ \text{ if and only if } \min \left\{ 1, \frac{1}{t} \right\} K(t, a) \rightarrow 0$$

as  $t \rightarrow 0$  and as  $t \rightarrow \infty$ . (2.6)

For  $0 \leq \theta \leq 1$ ,  $0 < q \leq \infty$  and  $b \in SV(0, \infty)$ , the space  $\bar{A}_{\theta,q;b} = (A_0, A_1)_{\theta,q;b}$  consists of all those  $a \in A_0 + A_1$  that have a finite quasi-norm

$$\|a\|_{\bar{A}_{\theta,q;b}} = \|a\|_{(A_0, A_1)_{\theta,q;b}} = \left( \sum_{m=-\infty}^{\infty} [2^{-\theta m} b(2^m) K(2^m, a)]^q \right)^{1/q}$$

(the sum should be replaced by the supremum when  $q = \infty$ ). See [29, 37]. If  $b \equiv 1$  and  $0 < \theta < 1$ , we recover the classical real interpolation space  $(A_0, A_1)_{\theta,q;b}$  (see [3, 4, 8, 43]). If  $0 < \theta < 1$  then  $(A_0, A_1)_{\theta,q;b}$  is a special case of the real method with function parameter (see [30, 40]). If  $\theta = 0, 1$ ,  $\alpha_0, \alpha_\infty \in \mathbb{R}$  and

$$b(t) = \begin{cases} (1 + |\log t|)^{\alpha_0} & \text{if } 0 < t \leq 1, \\ (1 + |\log t|)^{\alpha_\infty} & \text{if } 1 < t < \infty, \end{cases}$$

then we recover the logarithmic interpolation spaces  $\bar{A}_{\theta,q;(\alpha_0, \alpha_\infty)}$  (see [10, 16, 20, 27, 28]).

We are mainly interested here in the limiting spaces  $\bar{A}_{0,q;b}$  and  $\bar{A}_{1,q;b}$ . Since  $K(t, a; A_0, A_1) = t K(t^{-1}, a; A_1, A_0)$ , they are related by the equality

$$(A_0, A_1)_{0,q;b} = (A_1, A_0)_{1,q;v} \text{ where } v(t) = b(1/t). \tag{2.7}$$

Note that  $v$  is also slowly varying on  $(0, \infty)$ . Due to equality (2.7), in what follows we focus on the case  $\theta = 0$ .

As it is shown in [28],  $(A_0, A_1)_{0,q;b}$  is an intermediate space with respect to  $\bar{A}$  if and only if

$$\left( \int_1^\infty b(t)^q dt/t \right)^{1/q} < \infty. \tag{2.8}$$

Let  $\bar{B} = (B_0, B_1)$  be another quasi-Banach couple. We write  $T \in \mathcal{L}(\bar{A}, \bar{B})$  to mean that  $T$  is a bounded linear operator from  $A_0 + A_1$  into  $B_0 + B_1$  such that the restrictions  $T : A_j \rightarrow B_j$  are bounded for  $j = 0, 1$ . Then the restriction

$$T : (A_0, A_1)_{0,q;b} \rightarrow (B_0, B_1)_{0,q;b}$$

is also bounded. Indeed, if  $M_j$  is bigger than or equal to the norm of  $T : A_j \rightarrow B_j$ ,  $j = 0, 1$ , then

$$K(t, Ta; B_0, B_1) \leq M_0 K\left(\frac{tM_1}{M_0}, a; A_0, A_1\right).$$

Therefore, if  $M_1 \leq M_0$ , we obtain that  $\|T\|_{\bar{A}_{0,q;b}, \bar{B}_{0,q;b}} \leq M_0$ . If  $M_0 < M_1$  then we can find  $r \in \mathbb{N} \cup \{0\}$  such that  $2^r \leq M_1/M_0 < 2^{r+1}$ . Hence

$$\begin{aligned} \|Ta\|_{\bar{B}_{0,b;q}} &\leq \left( \sum_{m=-\infty}^\infty [b(2^m)M_0 K(2^{(m+r+1)}, a)]^q \right)^{1/q} \\ &\leq M_0 \bar{b}(2^{-r-1}) \|a\|_{\bar{A}_{0,b;q}} \\ &\leq cM_0 \bar{b} \left( \frac{M_0}{M_1} \right) \|a\|_{\bar{A}_{0,b;q}} \end{aligned}$$

where we have used (2.3) in the last inequality. Therefore

$$\|T\|_{\bar{A}_{0,q;b}, \bar{B}_{0,q;b}} \leq \begin{cases} M_0 & \text{if } M_1 \leq M_0, \\ cM_0 \bar{b} \left( \frac{M_0}{M_1} \right) & \text{if } M_0 < M_1, \end{cases} \tag{2.9}$$

where  $c > 0$  is a constant depending only on  $b$ .

If  $(T_n) \subseteq \mathcal{L}(A_0 + A_1, B_0 + B_1)$  with

$$\sup \{ \|T_n\|_{A_1, B_1} : n \in \mathbb{N} \} < \infty \text{ and } \lim_{n \rightarrow \infty} \|T_n\|_{A_0, B_0} = 0,$$

then it follows from (2.9) and (2.1) that

$$\lim_{n \rightarrow \infty} \|T_n\|_{\bar{A}_{0,q;b}, \bar{B}_{0,q;b}} = 0. \tag{2.10}$$

Next we show a sufficient condition on  $b$  for the inclusion  $(A_0, A_1)_{0,q;b} \subseteq (A_0 + A_1)^\circ$ . Let  $0 < q \leq \infty$  and take any  $a \in (A_0, A_1)_{0,q;b}$ . Then

$$\left( \sum_{n=-\infty}^\infty \left[ \frac{b(2^n)}{\min\{1, 2^{-n}\}} \min\{1, 2^{-n}\} K(2^n, a) \right]^q \right)^{1/q} = \|a\|_{\bar{A}_{0,q;b}} < \infty. \tag{2.11}$$

Since  $tb(t)$  is equivalent to a non-decreasing function, we have  $(\sum_{n=0}^\infty [2^n b(2^n)]^q)^{1/q} = \infty$ . Hence, from (2.11) it follows that

$$\lim_{t \rightarrow \infty} \frac{1}{t} K(t, a) = \lim_{n \rightarrow \infty} \frac{1}{2^n} K(2^n, a) = 0.$$

On the other hand, if we assume

$$\begin{cases} (\int_0^1 b(t)^q dt/t)^{1/q} = \infty & \text{if } 0 < q < \infty, \\ \lim_{t \rightarrow 0} b(t) = \infty & \text{if } q = \infty, \end{cases} \tag{2.12}$$

then we also have that  $\lim_{t \rightarrow 0} K(t, a) = 0$ . Having in mind (2.6), it turns out that if (2.12) is satisfied then  $(A_0, A_1)_{0,q;b} \subseteq (A_0 + A_1)^\circ$ .

The Gagliardo completion  $A_j^\sim$  of  $A_j$  consists of all those  $a \in A_0 + A_1$  having a finite quasi-norm

$$\|a\|_{A_j^\sim} = \sup\{t^{-j} K(t, a) : t > 0\} \quad (\text{see [4,3]}).$$

We have that  $A_j \hookrightarrow A_j^\sim$  for  $j = 0, 1$ . The quasi-Banach couple  $\bar{A}$  is called *mutually closed* if  $A_j = A_j^\sim$  for  $j = 0, 1$ .

If  $\Gamma$  is a quasi-Banach sequence lattice and  $\bar{A} = (A_0, A_1)$  is a  $p$ -Banach couple, then the  $J$ -space  $\bar{A}_{\Gamma;J} = (A_0, A_1)_{\Gamma;J}$  is formed by all sums  $a = \sum_{m=-\infty}^\infty u_m$  (convergence in  $A_0 + A_1$ ), where  $(u_m) \subseteq A_0 \cap A_1$  and  $(J(2^m, u_m)) \in \Gamma$ . We endow  $\bar{A}_{\Gamma;J}$  with the quasi-norm

$$\|a\|_{\bar{A}_{\Gamma;J}} = \|a\|_{(A_0, A_1)_{\Gamma;J}} = \inf \left\{ \| (J(2^m, u_m)) \|_{\Gamma} : a = \sum_{m=-\infty}^\infty u_m \right\}$$

(see [37]).

Next, we give a description of  $(A_0, A_1)_{0,q;b}$  by means of the  $J$ -functional.

**Theorem 2.1** *Let  $\bar{A} = (A_0, A_1)$  be a mutually closed  $p$ -Banach couple ( $0 < p \leq 1$ ). Let  $0 < q \leq \infty$  and let  $b \in SV(0, \infty)$  satisfying (2.8) and (2.12). Put  $\Lambda = (\ell_p, \ell_p(2^{-m}))_{0,q;b}$ . Then we have with equivalent quasi-norms*

$$(A_0, A_1)_{0,q;b} = (A_0, A_1)_{\Lambda;J}.$$

**Proof** Let  $a \in (A_0, A_1)_{0,q;b}$ . By the assumption on  $b$ , we know that  $(A_0, A_1)_{0,q;b} \subseteq (A_0 + A_1)^\circ$ . Hence, according to [37, Theorem 3.2], there exists  $(u_m) \subseteq A_0 \cap A_1$  such that  $a = \sum_{m=-\infty}^\infty u_m$  (in  $A_0 + A_1$ ) and

$$\left( \sum_{m=-\infty}^{\infty} \min(1, 2^{k-m})^p J(2^m, u_m)^p \right)^{1/p} \leq cK(2^k, a), \quad k \in \mathbb{Z},$$

where  $c$  only depends on  $p$ . Whence

$$\begin{aligned} \|a\|_{(A_0, A_1)_{\Lambda; J}} &\leq \|(J(2^m, u_m))\|_{(\ell_p, \ell_p(2^{-m}))_{0, q; b}} \\ &\leq \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k) K_p(2^k, ((J(2^m, u_m)); \ell_p, \ell_p(2^{-m}))) \right]^q \right)^{1/q} \\ &\leq \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k) \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{k-m})^p J(2^m, u_m)^p \right)^{1/p} \right]^q \right)^{1/q} \\ &\leq c \left( \sum_{k=-\infty}^{\infty} [b(2^k) K(2^k, a)]^q \right)^{1/q} \\ &= c \|a\|_{(A_0, A_1)_{0, q; b}}. \end{aligned}$$

Conversely, take any  $a \in (A_0, A_1)_{\Lambda; J}$ . We can find a  $J$ -representation  $a = \sum_{m=-\infty}^{\infty} u_m$  with  $\|(J(2^m, u_m))\|_{\Lambda} \leq 2 \|a\|_{(A_0, A_1)_{\Lambda; J}}$ . Since

$$K_p(2^k, a) \leq \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{k-m})^p J(2^m, u_m)^p \right)^{1/p}, \quad k \in \mathbb{Z},$$

we obtain that

$$\begin{aligned} \|a\|_{(A_0, A_1)_{0, q; b}} &\leq \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k) \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{k-m})^p J(2^m, u_m)^p \right)^{1/p} \right]^q \right)^{1/q} \\ &\leq 2^{1/p-1} \|(J(2^m, u_m))\|_{(\ell_p, \ell_p(2^{-m}))_{0, q; b}} \\ &\leq 2^{1/p} \|a\|_{(A_0, A_1)_{\Lambda; J}}. \end{aligned}$$

□

In Theorem 2.1, the sequence space that defines  $(A_0, A_1)_{0, q; b}$  as a  $J$ -space is not explicitly described, it appears as the interpolation space  $\Lambda = (\ell_p, \ell_p(2^{-m}))_{0, q; b}$  instead, what is enough for our aims here. Assuming extra conditions on the couple  $(A_0, A_1)$  and on the function  $b$ , there are several papers in the literature where the sequence space  $\Lambda$  is explicitly described. More precisely, in the case of logarithmic interpolation spaces, explicit descriptions as  $J$ -spaces have been obtained by Cobos and Kühn [16] for the case of ordered Banach couples, by Cobos and Segurado [20] and Besoy, Cobos and Fernández-Cabrera [7] for general Banach couples and by

Besoy and Cobos [6] for quasi-Banach couples. If  $(A_0, A_1)$  is a Banach couple and  $1 \leq q \leq \infty$ , an explicit description of  $(A_0, A_1)_{0,q;b}$  as a  $J$ -space has been recently established by Grover and Opic [38].

The following estimate for the norm of the shift operator  $\tau_k$  on  $\Lambda$  will be useful later.

**Lemma 2.2** *Let  $0 < p \leq 1, 0 < q \leq \infty$  and let  $b \in SV(0, \infty)$  satisfying (2.8). Put  $\Lambda = (\ell_p, \ell_p(2^{-m}))_{0,q;b}$ . Then, for any  $k \in \mathbb{Z}$ , we have*

$$\|\tau_k\|_{\Lambda, \Lambda} \leq 2^{1/p-1} \bar{b}(2^{-k}).$$

**Proof** Given any  $\xi = (\xi_m) \in \Lambda$ , we have

$$\begin{aligned} \|\tau_k \xi\|_{\Lambda} &\leq \left( \sum_{n=-\infty}^{\infty} [b(2^n)K_p(2^n, \tau_k \xi; \ell_p, \ell_p(2^{-m}))]^q \right)^{1/q} \\ &= \left( \sum_{n=-\infty}^{\infty} \left[ b(2^n) \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{n-m})^p |\xi_{m+k}|^p \right)^{1/p} \right]^q \right)^{1/q} \\ &\leq \bar{b}(2^{-k}) \left( \sum_{n=-\infty}^{\infty} \left[ b(2^{n+k}) \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{n+k-m})^p |\xi_m|^p \right)^{1/p} \right]^q \right)^{1/q} \\ &\leq 2^{1/p-1} \bar{b}(2^{-k}) \left( \sum_{n=-\infty}^{\infty} [b(2^n)K(2^n, \xi)]^q \right)^{1/q} \\ &= 2^{1/p-1} \bar{b}(2^{-k}) \|\xi\|_{\Lambda} \end{aligned}$$

where we have used (2.5) in the penultimate inequality. □

### 3 Measure of non-compactness

Let  $A, B$  be quasi-Banach spaces and  $T \in \mathcal{L}(A, B)$ . The (ball) *measure of non-compactness*  $\beta(T) = \beta(T : A \rightarrow B)$  is defined to be the infimum of the set of numbers  $\sigma > 0$  for which there is a finite subset  $\{z_1, \dots, z_n\} \subseteq B$  such that

$$T(U_A) \subseteq \bigcup_{j=1}^n \{z_j + \sigma U_B\}.$$

Here  $U_A, U_B$  are the closed unit balls of  $A$  and  $B$ , respectively. See [24] for details on the measure of non-compactness. Note that  $\beta(T) \leq \|T\|_{A,B}$  and that  $\beta(T) = 0$  if and only if  $T$  is compact. That is,  $\beta(T) = 0$  means that  $T$  transforms each bounded set of  $A$  into a set whose closure is compact in  $B$ .

If  $T_1$  is another operator belonging to  $\mathcal{L}(A, B)$ , then it is not difficult to check that

$$\beta(T + T_1 : A \rightarrow B) \leq c_B(\beta(T : A \rightarrow B) + \beta(T_1 : A \rightarrow B)).$$

If we assume that  $E, X$  are other quasi-Banach spaces and that  $S \in \mathcal{L}(B, E)$  and  $R \in \mathcal{L}(X, A)$ , then we have

$$\beta(STR : X \rightarrow E) \leq \|R\|_{X,A}\beta(T : A \rightarrow B)\|S\|_{B,E}.$$

Furthermore, if  $\|Sb\|_E = \|b\|_B$  for all  $b \in B$ , then

$$\beta(T : A \rightarrow B) \leq 2c_E\beta(ST : A \rightarrow E).$$

If for any  $a \in A$  with  $\|a\|_A < 1$ , there is  $x \in X$  with  $\|x\|_X < 1$  such that  $Rx = a$ , then

$$\beta(T : A \rightarrow B) \leq \beta(TR : X \rightarrow B).$$

We will use freely these properties in our later computations.

Next we establish the main result of the paper. It shows an estimate for the measure of non-compactness of an operator interpolated using parameters  $0, q, b$ .

**Theorem 3.1** *Let  $\bar{A} = (A_0, A_1), \bar{B} = (B_0, B_1)$  be quasi-Banach couples and let  $T \in \mathcal{L}(\bar{A}, \bar{B})$ . Let  $0 < q \leq \infty$  and  $b \in SV(0, \infty)$  satisfying (2.8) and (2.12). Then we have*

- (i)  $\beta(T : \bar{A}_{0,q;b} \rightarrow \bar{B}_{0,q;b}) = 0$  if  $\beta(T : A_0 \rightarrow B_0) = 0$ ,
- (ii)

$$\beta(T : \bar{A}_{0,q;b} \rightarrow \bar{B}_{0,q;b}) \leq C\beta(T : A_0 \rightarrow B_0) \text{ if } 0 \leq \beta(T : A_1 \rightarrow B_1) < \beta(T : A_0 \rightarrow B_0),$$

- (iii)

$$\beta(T : \bar{A}_{0,q;b} \rightarrow \bar{B}_{0,q;b}) \leq C \max \left\{ \beta(T : A_0 \rightarrow B_0), \beta(T : A_0 \rightarrow B_0)\bar{b} \left( \frac{\beta(T : A_0 \rightarrow B_0)}{\beta(T : A_1 \rightarrow B_1)} \right) \right\}$$

$$\text{if } 0 < \beta(T : A_0 \rightarrow B_0) \leq \beta(T : A_1 \rightarrow B_1).$$

Here  $C$  is a constant independent of  $T$ .

**Proof** *Step 1.* Consider the mutually closed quasi-Banach couples  $\bar{A}^\sim = (A_0^\sim, A_1^\sim), \bar{B}^\sim = (B_0^\sim, B_1^\sim)$ . The arguments of [3, Theorem V.1.5] may be modified to give that

$$K(t, a; A_0^{\sim}, A_1^{\sim}) \leq K(t, a; A_0, A_1) \leq \max\{c_{A_0}, c_{A_1}\}K(t, a; A_0^{\sim}, A_1^{\sim}).$$

Therefore,

$$(A_0, A_1)_{0,q;b} = (A_0^{\sim}, A_1^{\sim})_{0,q;b} \quad \text{and} \quad (B_0, B_1)_{0,q;b} = (B_0^{\sim}, B_1^{\sim})_{0,q;b}.$$

Besides,  $T \in \mathcal{L}(\overline{A^{\sim}}, \overline{B^{\sim}})$  and, according to [5, Lemma 3.1], we have

$$\beta(T : A_j^{\sim} \rightarrow B_j^{\sim}) \leq \max\{c_{B_0}, c_{B_1}\}\beta(T : A_j \rightarrow B_j), \quad j = 0, 1.$$

Consequently, without loss of generality we may assume in the following that the couples  $\overline{A} = (A_0, A_1)$  and  $\overline{B} = (B_0, B_1)$  are mutually closed. We may also assume that the spaces  $A_0, A_1, B_0, B_1$  are  $p$ -Banach for some  $0 < p \leq 1$ . Therefore, we can use Theorem 2.1.

*Step 2.* In this step we will introduce vector-valued sequence spaces and projections which will allow to split the operator  $T$ .

Let  $\Lambda = (\ell_p, \ell_p(2^{-m}))_{0,q;b}$ . By Theorem 2.1, we know that  $(A_0, A_1)_{0,q;b} = (A_0, A_1)_{\Lambda;J}$ . Consider the vector-valued sequence space  $\Lambda(G_m)$  where  $G_m = (A_0 \cap A_1, J(2^m, \cdot))$ ,  $m \in \mathbb{Z}$ . Let  $\pi : \Lambda(G_m) \rightarrow (A_0, A_1)_{\Lambda;J}$  be the linear operator defined by  $\pi(u_m) = \sum_{m=-\infty}^{\infty} u_m$  (convergence in  $A_0 + A_1$ ). Then  $\pi$  is surjective and induces the quasi-norm of  $(A_0, A_1)_{\Lambda;J}$ . Note also that  $\pi \in \mathcal{L}(\ell_p(2^{-mj}G_m), A_j)$ ,  $j = 0, 1$ , and its norm is less than or equal to 1.

Put  $\overline{\ell_p(G)} = (\ell_p(G_m), \ell_p(2^{-m}G_m))$ . The following projections will be useful. For  $n \in \mathbb{N}$  and  $u = (u_m) \in \ell_p(G_m) + \ell_p(2^{-m}G_m)$  let

$$\begin{aligned} P_n u &= (\dots, 0, 0, u_{-n}, u_{-n+1}, \dots, u_{n-1}, u_n, 0, 0, \dots), \\ P_n^+ u &= (\dots, 0, 0, u_{n+1}, u_{n+2}, u_{n+3}, \dots), \\ P_n^- u &= (\dots, u_{-n-3}, u_{-n-2}, u_{-n-1}, 0, 0, \dots). \end{aligned}$$

Then the identity operator  $I$  on  $\ell_p(G_m) + \ell_p(2^{-m}G_m)$  can be decomposed as  $I = P_n + P_n^+ + P_n^-$ ,  $n \in \mathbb{N}$ . These projections are bounded from  $\ell_p(2^{-mj}G_m)$  into  $\ell_p(2^{-mj}G_m)$  with norm less than or equal to 1 for  $j = 0, 1$ , and the same happens on  $\Lambda(G_m)$ . Furthermore,

$$\|P_n^+\|_{\ell_p(G_m), \ell_p(2^{-m}G_m)} = 2^{-(n+1)} = \|P_n^-\|_{\ell_p(2^{-m}G_m), \ell_p(G_m)}. \tag{3.1}$$

Write  $F_m = (B_0 + B_1, K(2^m, \cdot))$ ,  $m \in \mathbb{Z}$ . Then the linear operator  $\iota b = (\dots, b, b, b, \dots)$  is a metric injection from  $(B_0, B_1)_{0,q;b}$  into  $\ell_q(b(2^m)F_m)$ . Consider the couple  $\overline{\ell_{\infty}(F)} = (\ell_{\infty}(F_m), \ell_{\infty}(2^{-m}F_m))$ . Note that  $\iota : B_j \rightarrow \ell_{\infty}(2^{-mj}F_m)$  is bounded with norm less than or equal to 1. On  $\overline{\ell_{\infty}(F)}$  we can consider the corresponding sequences of projections that we denote by  $(Q_n), (Q_n^+), (Q_n^-)$ . They enjoy analogous properties as  $(P_n), (P_n^+)$  and  $(P_n^-)$ . In particular, we have

$$\|Q_n^+\|_{\ell_{\infty}(F_m), \ell_{\infty}(2^{-m}F_m)} = 2^{-(n+1)} = \|Q_n^-\|_{\ell_{\infty}(2^{-m}F_m), \ell_{\infty}(F_m)}. \tag{3.2}$$

The following diagram illustrates the situation

$$\begin{array}{ccccc} \ell_p(G_m) & \xrightarrow{\pi} & A_0 & \xrightarrow{T} & B_0 & \xrightarrow{\iota} & \ell_\infty(F_m) \\ \ell_p(2^{-m}G_m) & \xrightarrow{\pi} & A_1 & \xrightarrow{T} & B_1 & \xrightarrow{\iota} & \ell_\infty(2^{-m}F_m) \\ \hline \Lambda(G_m) & \xrightarrow{\pi} & \bar{A}_{0,q;b} & \xrightarrow{T} & \bar{B}_{0,q;b} & \xrightarrow{\iota} & \ell_q(b(2^m)F_m) \end{array}$$

In this diagram, the first three spaces of the last line are obtained by interpolation of the couple above and the fourth space contains the corresponding interpolation space. That is to say, we have

$$\begin{aligned} (\ell_p(G_m), \ell_p(2^{-m}G_m))_{0,q;b} &= \Lambda(G_m) \text{ and} \\ (\ell_\infty(F_m), \ell_\infty(2^{-m}(F_m)))_{0,q;b} &\hookrightarrow \ell_q(b(2^m)F_m). \end{aligned} \tag{3.3}$$

To establish the first formula we proceed as in the case when  $b(t) = (1 + |\log t|)^{\mathbb{A}}$  (see [5, Lemma 3.2]). Take any  $u = (u_m) \in (\ell_p(G_m), \ell_p(2^{-m}G_m))_{0,q;b}$ . For any  $k \in \mathbb{Z}$  and  $0 < \varepsilon < 1$ , there are  $u^j = (u_{j,m}) \in \ell_p(2^{-mj}G_m)$  such that  $u = u^0 + u^1$  and

$$\|u^0\|_{\ell_p(G_m)} + 2^k \|u^1\|_{\ell_p(2^{-m}G_m)} \leq (1 + \varepsilon)K(2^k, u; \ell_p(G_m), \ell_p(2^{-m}G_m)).$$

Then

$$\begin{aligned} \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{k-m})^p \|u_m\|_{G_m}^p \right)^{1/p} &\leq (\|u^0\|_{\ell_p(G_m)}^p + 2^{kp} \|u^1\|_{\ell_p(2^{-m}G_m)}^p)^{1/p} \\ &\leq 2^{1/p-1} (1 + \varepsilon)K(2^k, u; \ell_p(G_m), \ell_p(2^{-m}G_m)) \end{aligned}$$

and thus

$$\begin{aligned} \|u\|_{\Lambda(G_m)} &= \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k)K(2^k, ( \|u_m\|_{G_m} ); \ell_p, \ell_p(2^{-m})) \right]^q \right)^{1/q} \\ &\leq \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k) \left( \sum_{m=-\infty}^{\infty} (\min(1, 2^{k-m}) \|u_m\|_{G_m})^p \right)^{1/p} \right]^q \right)^{1/q} \\ &\leq 2^{1/p-1} (1 + \varepsilon) \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k)K(2^k, u; \ell_p(G_m), \ell_p(2^{-m}G_m)) \right]^q \right)^{1/q} \\ &\leq 2^{1/p} \|u\|_{(\ell_p(G_m), \ell_p(2^{-m}G_m))_{0,q;b}}. \end{aligned}$$

Reciprocally, if  $u = (u_m) \in \Lambda(G_m)$ , given any  $k \in \mathbb{Z}$  we can decompose  $u = u^0 + u^1$  with

$$u_{0,m} = \begin{cases} u_m & \text{if } m \leq k, \\ 0 & \text{if } m > k, \end{cases}, \quad u_{1,m} = \begin{cases} 0 & \text{if } m \leq k, \\ u_m & \text{if } m > k. \end{cases}$$

Then  $u^0 \in \ell_p(G_m)$ ,  $u^1 \in \ell_p(2^{-m}G_m)$  and we have

$$\begin{aligned} & K(2^k, u; \ell_p(G_m), \ell_p(2^{-m}G_m)) \\ & \leq \left( \sum_{m=-\infty}^k \|u_m\|_{G_m}^p \right)^{1/p} + 2^k \left( \sum_{m=k+1}^{\infty} (2^{-m}\|u_m\|_{G_m})^p \right)^{1/p} \\ & \leq 2 \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{k-m})^p \|u_m\|_{G_m}^p \right)^{1/p}. \end{aligned}$$

Consequently,

$$\begin{aligned} & \|u\|_{(\ell_p(G_m), \ell_p(2^{-m}G_m))_{0,q;b}} \\ & = \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k) K(2^k, u; \ell_p(G_m), \ell_p(2^{-m}G_m)) \right]^q \right)^{1/q} \\ & \leq 2 \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k) \left( \sum_{m=-\infty}^{\infty} \min(1, 2^{k-m})^p \|u_m\|_{G_m}^p \right)^{1/p} \right]^q \right)^{1/q} \\ & = 2 \left( \sum_{k=-\infty}^{\infty} \left[ b(2^k) K_p(2^k, (\|u_m\|_{G_m}); \ell_p, \ell_p(2^{-m})) \right]^q \right)^{1/q} \\ & \leq 2^{1/p} \|u\|_{\Lambda(G_m)}. \end{aligned}$$

To establish the second embedding in (3.3), take any  $x = (x_m) \in (\ell_\infty(F_m), \ell_\infty(2^{-m}F_m))_{0,q;b}$ . Give any decomposition  $x = x^0 + x^1$  with  $x^0 = (x_{0,m}) \in \ell_\infty(F_m)$  and  $x^1 = (x_{1,m}) \in \ell_\infty(2^{-m}F_m)$ , and any  $k \in \mathbb{Z}$ , we obtain

$$\begin{aligned} \|x_k\|_{F_k} & \leq c(\|x_k^0\|_{F_k} + \|x_k^1\|_{F_k}) \\ & \leq c(\|x^0\|_{\ell_\infty(F_m)} + 2^k \|x^1\|_{\ell_\infty(2^{-m}F_m)}). \end{aligned}$$

It follows that

$$\|x_k\|_{F_k} \leq cK(2^k, x; \ell_\infty(F_m), \ell_\infty(2^{-m}F_m)), \quad k \in \mathbb{Z}.$$

Therefore,  $\|x\|_{\ell_q(b(2^m)F_m)} \leq c\|x\|_{(\ell_\infty(F_m), \ell_\infty(2^{-m}F_m))_{0,q;b}}$  as we wanted.

Put  $\hat{T} = \iota T \pi$ . Since

$$\begin{aligned} \beta(T : \bar{A}_{0,q;b} \rightarrow \bar{B}_{0,q;b}) & \leq c_1 \beta(\iota T : \bar{A}_{0,q;b} \rightarrow \ell_q(b(2^m)F_m)) \\ & \leq c_2 \beta(\hat{T} : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)), \end{aligned}$$

it suffices to estimate the measure of non-compactness of  $\hat{T}$  acting between the vector-valued sequence spaces. With this aim, for  $n \in \mathbb{N}$  we decompose  $\hat{T}$  as

$$\begin{aligned} \hat{T} &= \hat{T}P_n + \hat{T}(P_n^+ + P_n^-) \\ &= \hat{T}P_n + Q_n\hat{T}(P_n^+ + P_n^-) + Q_n^-\hat{T}P_n^+ + Q_n^+\hat{T}P_n^- + Q_n^-\hat{T}P_n^- + Q_n^+\hat{T}P_n^+ \end{aligned}$$

and we proceed to estimate the measure of non-compactness of each of these six operators acting from  $\Lambda(G_m)$  into  $\ell_q(b(2^m)F_m)$ .

*Step 3.* We start with  $Q_n^-\hat{T}P_n^+$ . We are going to show that  $\beta(Q_n^-\hat{T}P_n^+ : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m))$  tends to 0 as  $n \rightarrow \infty$ .

Using the factorization

$$\ell_p(G_m) \xrightarrow{P_n^+} \ell_p(2^{-m}G_m) \xrightarrow{\hat{T}} \ell_\infty(2^{-m}F_m) \xrightarrow{Q_n^-} \ell_\infty(F_m)$$

and (3.1) and (3.2), we get

$$\|Q_n^-\hat{T}P_n^+ : \ell_p(G_m) \rightarrow \ell_\infty(F_m)\| \leq 2^{-(n+1)} \|T : A_1 \rightarrow B_1\| 2^{-(n+1)} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

In addition, the factorization

$$\ell_p(2^{-m}G_m) \xrightarrow{P_n^+} \ell_p(2^{-m}G_m) \xrightarrow{\hat{T}} \ell_\infty(2^{-m}F_m) \xrightarrow{Q_n^-} \ell_\infty(2^{-m}F_m)$$


yields that

$$\|Q_n^-\hat{T}P_n^+ : \ell_p(2^{-m}G_m) \rightarrow \ell_\infty(2^{-m}F_m)\| \leq \|T : A_1 \rightarrow B_1\| \text{ for any } n \in \mathbb{N}.$$

Therefore, by formulae (3.3) and (2.10), we obtain

$$\begin{aligned} &\lim_{n \rightarrow \infty} \beta(Q_n^-\hat{T}P_n^+ : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) \\ &\leq c_1 \lim_{n \rightarrow \infty} \|Q_n^-\hat{T}P_n^+\|_{\overline{\ell_p(G)_{0,q;b}}, \overline{\ell_\infty(F)_{0,q;b}}} = 0. \end{aligned}$$

*Step 4.* Consider  $Q_n^+\hat{T}P_n^-$ . Using the factorizations



and having in mind estimates (3.1), (3.2) and formulae (3.3) we get that  $\|P_n^- : \Lambda(G_m) \rightarrow \ell_p(G_m)\| \leq c_2$  and  $\|Q_n^+ : \ell_\infty(F_m) \rightarrow \ell_q(b(2^m)F_m)\| \leq c_3$ . Hence, with the help of the diagram

$$\begin{array}{ccc} \Lambda(G_m) & \xrightarrow{Q_n^+ \hat{T} P_n^-} & \ell_q(b(2^m)F_m) \\ \downarrow P_n^- & & \uparrow Q_n^+ \\ \ell_p(G_m) & \xrightarrow{\hat{T}} & \ell_\infty(F_m) \end{array}$$

we derive

$$\begin{aligned} \beta(Q_n^+ \hat{T} P_n^- : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) &\leq c_2 c_3 \beta(\hat{T} : \ell_p(G_m) \rightarrow \ell_\infty(F_m)) \\ &\leq c_2 c_3 \beta(T : A_0 \rightarrow B_0). \end{aligned}$$

*Step 5.* Now we proceed with  $Q_n^- \hat{T} P_n^-$ . Take any  $\sigma_j > \beta(T : A_j \rightarrow B_j)$ ,  $j = 0, 1$ . First we are going to compare  $\|Q_n^- \hat{T} P_n^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)}$  with  $\sigma_1$ . We have

$$\|Q_n^- \hat{T} P_n^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)} \leq \|\hat{T} P_n^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)}$$

and

$$\|\hat{T} P_1^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)} \geq \|\hat{T} P_2^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)} \geq \dots \geq 0.$$

Therefore, the sequence  $(\|\hat{T} P_n^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)})$  is convergent, say, to  $\tau \geq 0$ . Let  $(v_n) \subseteq U_{\ell_p(2^{-m}G_m)}$  such that  $\lim_{n \rightarrow \infty} \|\hat{T} P_n^- v_n\|_{\ell_\infty(2^{-m}F_m)} = \tau$ . To relate  $\tau$  and  $\sigma_1$ , let  $\{z_1, \dots, z_r\} \subseteq B_1$  such that

$$T\pi(U_{\ell_p(2^{-m}G_m)}) \subseteq \bigcup_{k=1}^r \{z_k + \sigma_1 U_{B_1}\}.$$

We can find a subsequence  $(v_{n'})$  of  $(v_n)$  and some  $1 \leq k \leq r$  such that  $\|T\pi P_{n'}^- v_{n'} - z_k\|_{B_1} \leq \sigma_1$  for all  $n'$ . Then, for any  $s \in \mathbb{Z}$ , we have

$$\begin{aligned} K(2^s, z_k) &\leq \|T\pi P_{n'}^- v_{n'}\|_{B_0} + 2^s \|z_k - T\pi P_{n'}^- v_{n'}\|_{B_1} \\ &\leq \|P_{n'}^- v_{n'}\|_{\ell_p(G_m)} \|T\|_{A_0, B_0} + 2^s \sigma_1 \\ &\leq 2^{-n'} \|T\|_{A_0, B_0} + 2^s \sigma_1 \rightarrow 2^s \sigma_1 \text{ as } n' \rightarrow \infty. \end{aligned}$$

It follows that

$$\|tz_k\|_{\ell_\infty(2^{-m}F_m)} = \sup_{s \in \mathbb{Z}} \{2^{-s} K(2^s, z_k)\} \leq \sigma_1.$$

Hence,

$$\begin{aligned} \tau &= \lim_{n' \rightarrow \infty} \|\hat{T} P_{n'}^- v_{n'}\|_{\ell_\infty(2^{-m}F_m)} \\ &\leq \max\{c_{B_0}, c_{B_1}\} \sup_{n'} \{\|\hat{T} P_{n'}^- v_{n'} - tz_k\|_{\ell_\infty(2^{-m}F_m)} + \|tz_k\|_{\ell_\infty(2^{-m}F_m)}\} \end{aligned}$$

$$\begin{aligned} &\leq \max\{c_{B_0}, c_{B_1}\} \sup_{n'} \{\|T\pi P_{n'}^- v_{n'} - z_k\|_{B_1} + \sigma_1\} \\ &\leq 2 \max\{c_{B_0}, c_{B_1}\} \sigma_1. \end{aligned}$$

Since the sequence  $(\|\hat{T} P_n^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)})$  is decreasing, we conclude that there exists  $N_1 \in \mathbb{N}$  such that if  $n \geq N_1$  then

$$\begin{aligned} \|\mathcal{Q}_n^- \hat{T} P_n^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)} &\leq \|\hat{T} P_n^-\|_{\ell_p(2^{-m}G_m), \ell_\infty(2^{-m}F_m)} \\ &\leq 3 \max\{c_{B_0}, c_{B_1}\} \sigma_1. \end{aligned}$$

Next we compare  $\|\mathcal{Q}_n^- \hat{T} P_n^-\|_{\ell_p(G_m), \ell_\infty(F_m)}$  with  $\sigma_0$ . Since sequences having a finite number of coordinates different from 0 are dense in  $\ell_p(G_m)$ , we can find  $\{d_1, \dots, d_s\} \subseteq U_{\ell_p(G_m)}$  such that each  $d_k$  has a finite number of coordinates different from 0 and with

$$\hat{T}(U_{\ell_p(G_m)}) \subseteq \bigcup_{k=1}^s \{\hat{T}d_k + c_4\sigma_0 U_{\ell_\infty(F_m)}\}$$

where  $c_4 = 3 \max\{c_{B_0}, c_{B_1}\}^2$ . We can also find  $N_2 \in \mathbb{N}$  such that if  $n \geq N_2$  we have

$$\|\mathcal{Q}_n^- \hat{T}d_k\|_{\ell_\infty(F_m)} \leq 2^{-(n+1)} \|\hat{T}d_k\|_{\ell_\infty(2^{-m}F_m)} \leq \sigma_0 \text{ for any } 1 \leq k \leq s.$$

Take any  $n \geq N_2$  and any  $u \in U_{\ell_p(G_m)}$ . Then  $P_n^- u \in U_{\ell_p(G_m)}$  and so there is  $1 \leq k \leq s$  such that  $\|\hat{T} P_n^- u - \hat{T}d_k\|_{\ell_\infty(F_m)} \leq c_4\sigma_0$ . Therefore,  $\|\mathcal{Q}_n^- \hat{T} P_n^- u\|_{\ell_\infty(F_m)} \leq \|\mathcal{Q}_n^- \hat{T} P_n^- u - \mathcal{Q}_n^- \hat{T}d_k\|_{\ell_\infty(F_m)} + \|\mathcal{Q}_n^- \hat{T}d_k\|_{\ell_\infty(F_m)} \leq 2c_4\sigma_0$ .

Finally, using (3.3) and (2.9), we derive that there is  $N \in \mathbb{N}$  such that if  $n \geq N$  then

$$\begin{aligned} \beta(\mathcal{Q}_n^- \hat{T} P_n^- : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) &\leq c_5 \|\mathcal{Q}_n^- \hat{T} P_n^-\|_{\overline{\ell_p(G)}_{0,q;b}, \overline{\ell_\infty(F)}_{0,q;b}} \\ &\leq \begin{cases} c_6\sigma_0 & \text{if } \sigma_1 \leq \sigma_0, \\ c_6\sigma_0 \bar{b} \left(\frac{\sigma_0}{\sigma_1}\right) & \text{if } \sigma_0 < \sigma_1. \end{cases} \end{aligned}$$

With similar arguments one can show that there is a constant  $c_7 > 0$  such that

$$\beta(\mathcal{Q}_n^+ \hat{T} P_n^+ : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) \leq \begin{cases} c_7\sigma_0 & \text{if } \sigma_1 \leq \sigma_0, \\ c_7\sigma_0 \bar{b} \left(\frac{\sigma_0}{\sigma_1}\right) & \text{if } \sigma_0 < \sigma_1. \end{cases}$$

*Step 6.* Given any quasi-Banach sequence lattice  $\Gamma$ , we can define a quasi-norm  $\|\cdot\|_{\tilde{\Gamma}}$  in  $\mathbb{R}^{2n+1}$  by  $\|x\|_{\tilde{\Gamma}} = \|\tilde{x}\|_{\Gamma}$ , where  $x = (x_k)_{-n \leq k \leq n} \in \mathbb{R}^{2n+1}$ ,  $\tilde{x} = \sum_{k=-n}^n x_k e_k$ ,  $e_k = (\delta_m^k)_{m \in \mathbb{Z}}$  and  $\delta_m^k$  is the Kronecker delta. Compactness of the unit ball  $U = U_{(\mathbb{R}^{2n+1}, \|\cdot\|_{\tilde{\Gamma}})}$  in  $(\mathbb{R}^{2n+1}, \|\cdot\|_{\tilde{\Gamma}})$  will be useful to estimate the measure of non-compactness of the remaining operators.

Let  $\sigma_j > \beta(T : A_j \rightarrow B_j), j = 0, 1$ . We can find finite sets  $\Sigma_j = \{h_1^j, \dots, h_{L_j}^j\} \subseteq B_j$  such that

$$T(U_{A_j}) \subseteq \bigcup_{l=1}^{L_j} \{h_l^j + \sigma_j U_{B_j}\}, \quad j = 0, 1. \tag{3.4}$$

Let  $N \in \mathbb{N}$  such that  $2^{N-1} \leq \sigma_1/\sigma_0 < 2^N$  if  $\sigma_0 \leq \sigma_1$  and let  $N = 0$  if  $\sigma_1 < \sigma_0$ .

As for  $\hat{T}P_n$ , consider the quasi-norm  $\|\cdot\|_{\tilde{\Lambda}}$  on  $\mathbb{R}^{2n+1}$  and let  $\eta = \left\| \sum_{k=-n}^n \frac{e_k}{\|e_k\|_{\Lambda}} \right\|_{\Lambda}^{-1}$ . By compactness of  $U = U_{(\mathbb{R}^{2n+1}, \|\cdot\|_{\tilde{\Lambda}})}$ , we can find a finite set  $\Upsilon = \{\lambda^1, \dots, \lambda^s\} \subseteq U$  such that

$$U \subseteq \bigcup_{d=1}^s \{\lambda^d + \eta U\}.$$

We associate to each  $\lambda^d = (\lambda_k^d)_{-n \leq k \leq n}$  the numbers

$$\varphi_k^j = \varphi_{k,\lambda^d}^j = \left( \frac{\eta}{\|e_k\|_{\Lambda}} + |\lambda_k^d| \right) 2^{-kj}, \quad j = 0, 1.$$

Next, for  $-n \leq k \leq n, \lambda^d \in \Upsilon, h_l^0 \in \Sigma_0$  and  $h_y^1 \in \Sigma_1$  in (3.4), pick any  $g_k$  in the intersection  $(\varphi_k^0 h_l^0 + \varphi_k^0 \sigma_0 U_{B_0}) \cap (\varphi_k^1 h_y^1 + \varphi_k^1 \sigma_1 U_{B_1})$  provided it is non-empty and let  $g_k = 0$  otherwise. Consider the finite set  $\Phi$  formed by all sums  $\sum_{k=-n}^n g_k$ . We look at  $\overline{B_{0,q;b}}$  as a  $J$ -space. We have

$$\beta(\hat{T}P_n : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) \leq c_1 \beta(T\pi P_n : \Lambda(G_m) \rightarrow \overline{B_{\Lambda;J}}).$$

We are going to estimate the last term with the help of  $\Phi$ .

For any  $u = (u_m) \in U_{\Lambda(G_m)}$ , we can find  $\lambda^d \in \Upsilon$  such that

$$|J(2^k, u_k) - \lambda_k^d| \|e_k\|_{\Lambda} \leq \|(J(2^k, u_k) - \lambda_k^d)\|_{\tilde{\Lambda}} \leq \eta, \quad -n \leq k \leq n.$$

It follows that  $|J(2^k, u_k)| \leq \frac{\eta}{\|e_k\|_{\Lambda}} + |\lambda_k^d| = 2^{kj} \varphi_k^j$ . This yields that  $\|u_k\|_{A_j} \leq \varphi_k^j, -n \leq k \leq n, j = 0, 1$ . By (3.4), there are  $h_l^0 \in \Sigma_0$  and  $h_y^1 \in \Sigma_1$  such that

$$\|Tu_k - \varphi_k^0 h_l^0\|_{B_0} \leq \varphi_k^0 \sigma_0$$

and

$$\|Tu_k - \varphi_k^1 h_y^1\|_{B_1} \leq \varphi_k^1 \sigma_1.$$

Hence, the intersection  $(\varphi_k^0 h_l^0 + \varphi_k^0 \sigma_0 U_{B_0}) \cap (\varphi_k^1 h_y^1 + \varphi_k^1 \sigma_1 U_{B_1})$  is not empty and for the  $g_k$  corresponding to that intersection we have

$$\begin{aligned} J(2^{k-N}, Tu_k - g_k) &\leq \max\{\|Tu_k - \varphi_k^0 h_l^0\|_{B_0}^p + \|\varphi_k^0 h_l^0 - g_k\|_{B_0}^p, \\ &\quad 2^{(k-N)p}(\|Tu_k - \varphi_k^1 h_y^1\|_{B_1}^p + \|\varphi_k^1 h_y^1 - g_k\|_{B_1}^p)\}^{1/p} \\ &\leq 2^{1/p} \max\{\sigma_0, 2^{-N}\sigma_1\} \varphi_k^0. \end{aligned}$$

Then,  $g = \sum_{k=-n}^n g_k$  belongs to  $\Phi$  and

$$\begin{aligned} \|T\pi P_n u - g\|_{\bar{B}_{\Lambda;J}} &= \left\| \sum_{k=-n}^n (Tu_k - g_k) \right\|_{\bar{B}_{\Lambda;J}} \\ &\leq \|\tau_N(\dots, 0, J(2^{-n-N}, Tu_{-n} - g_{-n}), \dots, J(2^{n-N}, Tu_n - g_n), 0, 0, \dots)\|_{\Lambda} \\ &\leq 2^{1/p} \|\tau_N\|_{\Lambda, \Lambda} \max\{\sigma_0, 2^{-N}\sigma_1\} \|(\dots, 0, 0, \varphi_{-n}^0, \dots, \varphi_n^0, 0, 0, \dots)\|_{\Lambda} \\ &\leq c_2 \bar{b}(2^{-N}) \max\{\sigma_0, 2^{-N}\sigma_1\} \end{aligned}$$

where we have used Lemma 2.2 and definition of  $\varphi_k^0$  in the last inequality. Whence, according to the choice of  $N$  and (2.3), we obtain that

$$\begin{aligned} \beta(\hat{T}P_n : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) &\leq c_1 \beta(T\pi P_n : \Lambda(G_m) \rightarrow \bar{B}_{\Lambda;J}) \\ &\leq c_3 \bar{b}(2^{-N}) \max\{\sigma_0, 2^{-N}\sigma_1\} \\ &\leq \begin{cases} c_4 \sigma_0 & \text{if } \sigma_1 \leq \sigma_0, \\ c_4 \sigma_0 \bar{b} \left(\frac{\sigma_0}{\sigma_1}\right) & \text{if } \sigma_0 < \sigma_1. \end{cases} \end{aligned}$$

Next we consider  $Q_n \hat{T}(P_n^+ + P_n^-)$ . This time we work with  $\bar{A}_{0,q;b}$  and  $\bar{B}_{0,q;b}$  realized as  $K$ -spaces. We put  $\Delta = \ell_q(b(2^m))$ . We have

$$\beta(Q_n \hat{T}(P_n^+ + P_n^-) : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) \leq c_5 \beta(Q_n \hat{T} : \bar{A}_{0,q;b} \rightarrow \Delta(F_m)).$$

Let now  $\eta = \left\| \sum_{k=-n}^n \frac{e_k}{\|e_k\|_{\Delta}} \right\|_{\Delta}^{-1}$  and consider on  $\mathbb{R}^{2n+1}$  the quasi-norm  $\|\cdot\|_{\tilde{\Delta}}$ . There is a finite set  $\Psi = \{\mu^1, \dots, \mu^s\} \subseteq U = U_{(\mathbb{R}^{2n+1}, \|\cdot\|_{\tilde{\Delta}})}$  such that

$$U \subseteq \bigcup_{d=1}^s \{\mu^d + \eta U\}.$$

Starting from  $\mu^d = (\mu_k^d)_{-n \leq k \leq n}$  we define the numbers

$$\psi_k^j = \psi_{k, \mu^d}^j = \bar{b}(2^{-N}) \left( \frac{\eta}{\|e_k\|_{\Delta}} + |\mu_k^d| \right) 2^{-(k+N)j}, \quad j = 0, 1,$$

where  $N$  was defined in the following line to (3.4). Let  $\Omega$  be the finite subset of  $\Delta(F_m)$  formed by all vectors  $z^{d,l,y} = (z_m^{d,l,y})_{m \in \mathbb{Z}}$  where

$$z_m^{d,l,y} = \begin{cases} 0 & \text{if } m \notin [-n, n] \\ \psi_m^0 h_l^0 + \psi_m^1 h_y^1 & \text{if } -n \leq m \leq n, \end{cases}$$

where  $h_l^0 \in \Sigma_0$  and  $h_y^1 \in \Sigma_1$  are the vectors of (3.4). We refer to  $z^{d,l,y}$  as the element of  $\Omega$  associated to  $\mu^d, h_l^0$  and  $h_y^1$ .

Given any  $a \in U_{\bar{A}_{0,q;b}}^-$ , using the shift operator  $\tau_N$  and (2.4), we have

$$\|(K(2^{m+N}, a))\|_{\Delta} \leq \|\tau_N\|_{\Delta} \|a\|_{\bar{A}_{0,q;b}} \leq \bar{b}(2^{-N}).$$

Therefore, there is  $\mu^d \in \Psi$  such that

$$\|(K(2^{m+N}, a) - \bar{b}(2^{-N})\mu_m^d)_{-n \leq m \leq n}\|_{\tilde{\Delta}} < \eta \bar{b}(2^{-N}).$$

Hence

$$\|K(2^{m+N}, a) - \bar{b}(2^{-N})\mu_m^d\|_{\Delta} < \eta \bar{b}(2^{-N}), \quad -n \leq m \leq n,$$

and so  $K(2^{m+N}, a) < \psi_m^0$  for  $-n \leq m \leq n$ . It follows that we can decompose  $a = a_{0,m} + a_{1,m}$  with  $a_{j,m}$  belonging to  $A_j$  and such that  $\|a_{0,m}\|_{A_0} + 2^{m+N} \|a_{1,m}\|_{A_1} < \psi_m^0$ . Therefore, there are  $h_l^0 \in \Sigma_0$  and  $h_y^1 \in \Sigma_1$  such that

$$\|Ta_{0,m} - \psi_m^0 h_l^0\|_{B_0} \leq \psi_m^0 \sigma_0$$

and

$$\|Ta_{1,m} - \psi_m^1 h_y^1\|_{B_1} \leq \psi_m^1 \sigma_1, \quad -n \leq m \leq n.$$

If we take  $z = z^{d,l,y}$  the element of  $\Omega$  associated to  $\mu^d, h_l^0$ , and  $h_y^1$ , then we have

$$\begin{aligned} \|Q_n \iota T a - z\|_{\Delta(F_m)} &= \|(K(2^m, T a - z_m^{d,l,y}))_{-n \leq m \leq n}\|_{\tilde{\Delta}} \\ &\leq \|(\|Ta_{0,m} - \psi_m^0 h_l^0\|_{B_0} + 2^m \|Ta_{1,m} - \psi_m^1 h_y^1\|_{B_1})_{-n \leq m \leq n}\|_{\tilde{\Delta}} \\ &\leq \|(\psi_m^0 \sigma_0 + 2^m \psi_m^1 \sigma_1)_{-n \leq m \leq n}\|_{\tilde{\Delta}} \\ &\leq c_6 \bar{b}(2^{-N})(\sigma_0 + 2^{-N} \sigma_1). \end{aligned}$$

Consequently,

$$\begin{aligned} \beta(Q_n \hat{T}(P_n^+ + P_n^-) : \Lambda(G_m) \rightarrow \ell_q(b(2^m)F_m)) &\leq c_5 \beta(Q_n \iota T : \bar{A}_{0,q;b} \rightarrow \Delta(F_m)) \\ &\leq c_7 \bar{b}(2^{-N})(\sigma_0 + 2^{-N} \sigma_1) \end{aligned}$$

$$\leq \begin{cases} c_8\sigma_0 & \text{if } \sigma_1 \leq \sigma_0, \\ c_8\sigma_0\bar{b}\left(\frac{\sigma_0}{\sigma_1}\right) & \text{if } \sigma_0 < \sigma_1, \end{cases}$$

where we have used the value of  $N$  and (2.3) in the last inequality.

*Step 7.* Collecting the estimates of the Steps 3 to 6, we conclude that there is a constant  $C > 0$  independent of  $T$  such that if we split the operator as in the Step 2 and we take a suitable  $n$ , then for  $\sigma_j > \beta(T : A_j \rightarrow B_j)$ , we have

$$\beta(\hat{T} : \Lambda(G_m) \rightarrow \ell_q(2^m F_m)) \leq \begin{cases} C\sigma_0 & \text{if } \sigma_1 \leq \sigma_0, \\ C \max \left\{ \sigma_0, \sigma_0\bar{b}\left(\frac{\sigma_0}{\sigma_1}\right) \right\} & \text{if } \sigma_0 < \sigma_1. \end{cases}$$

Then, if  $\beta(T : A_0 \rightarrow B_0) = 0$ , letting  $\sigma_0 \rightarrow 0$  and using (2.2) we obtain case (i) of the statement. If  $0 \leq \beta(T : A_1 \rightarrow B_1) < \beta(T : A_0 \rightarrow B_0)$ , letting  $\sigma_0 \rightarrow \beta(T : A_0 \rightarrow B_0)$  we get the case (ii). Finally, if  $0 < \beta(T : A_0 \rightarrow B_0) \leq \beta(T : A_1 \rightarrow B_1)$ , taking  $\sigma_j = (1 + \varepsilon)\beta(T : A_j \rightarrow B_j)$  and letting  $\varepsilon$  goes to 0 we derive the case (iii). This finishes the proof.  $\square$

**Remark 3.2** On the contrary to the case of the real method (see [17, 22]), if  $T \in \mathcal{L}(\bar{A}, \bar{B})$  and  $T : A_1 \rightarrow B_1$  is compact, then  $T : \bar{A}_{0,q;b} \rightarrow \bar{B}_{0,q;b}$  might not be compact. A counterexample can be found in [13, Remark 2.4].

For limiting methods with  $\theta = 1$  we have the following direct consequence of (2.7) and Theorem 3.1.

**Theorem 3.3** Let  $\bar{A} = (A_0, A_1)$ ,  $\bar{B} = (B_0, B_1)$  be quasi-Banach couples and let  $T \in \mathcal{L}(\bar{A}, \bar{B})$ . Let  $0 < q \leq \infty$  and  $v \in SV(0, \infty)$  satisfying

$$\left( \int_0^1 v(t)^q \frac{dt}{t} \right)^{1/q} < \infty, \text{ and also that}$$

$$\left( \int_1^\infty v(t)^q \frac{dt}{t} \right)^{1/q} = \infty \text{ if } q < \infty \text{ and } \lim_{t \rightarrow \infty} v(t) = \infty \text{ if } q = \infty.$$

Then we have

- (i)  $\beta(T : \bar{A}_{1,q;v} \rightarrow \bar{B}_{1,q;v}) = 0$  if  $\beta(T : A_1 \rightarrow B_1) = 0$ ,
- (ii)  $\beta(T : \bar{A}_{1,q;v} \rightarrow \bar{B}_{1,q;v}) \leq C\beta(T : A_1 \rightarrow B_1)$  if  $0 \leq \beta(T : A_0 \rightarrow B_0) < \beta(T : A_1 \rightarrow B_1)$ ,
- (iii)

$$\beta(T : \bar{A}_{1,q;v} \rightarrow \bar{B}_{1,q;v}) \leq C \max \left\{ \beta(T : A_1 \rightarrow B_1), \right.$$

$$\beta(T : A_1 \rightarrow B_1) \bar{v} \left( \frac{\beta(T : A_1 \rightarrow B_1)}{\beta(T : A_0 \rightarrow B_0)} \right) \Big\}$$

$$\text{if } 0 < \beta(T : A_1 \rightarrow B_1) \leq \beta(T : A_0 \rightarrow B_0).$$

Here  $C$  is a constant independent of  $T$ .

### 4 Applications

Let  $(R, \mu)$  be a non-atomic  $\sigma$ -finite measure space. For  $0 < p, q \leq \infty$  and  $b \in SV(0, \infty)$ , the *Lorentz–Karamata space*  $L_{p,q;b}(R)$  is formed by all (equivalent classes of) measurable functions  $f$  on  $R$  which have a finite quasi-norm

$$\|f\|_{L_{p,q;b}(R)} = \left( \int_0^\infty [t^{1/p} b(t) f^*(t)]^q \frac{dt}{t} \right)^{1/q}$$

(the integral should be replaced by the supremum if  $q = \infty$ ). Here  $f^*$  stands for the non-increasing rearrangement of  $f$  defined by

$$f^*(t) = \inf\{s > 0 : \mu\{x \in R : |f(x)| > s\} \leq t\}.$$

We refer to [25] and [29] for properties of Lorentz–Karamata spaces. Note that if  $b(t) = (1 + |\log t|)^a$  we get the *Lorentz–Zygmund spaces*  $L_{p,q}(\log L)_a$  (see [2, 3]). If  $\mathbb{A} = (\alpha_0, \alpha_\infty) \in \mathbb{R}^2$  and

$$b(t) = \ell^{\mathbb{A}}(t) = \begin{cases} (1 + |\log t|)^{\alpha_0} & \text{for } 0 < t \leq 1, \\ (1 + |\log t|)^{\alpha_\infty} & \text{for } 1 < t < \infty, \end{cases}$$

then we obtain the *generalized Lorentz–Zygmund spaces*  $L_{p,q}(\log L)_{\mathbb{A}}(R)$  (see [39]). If  $b \equiv 1$  then we obtain the *Lorentz spaces*  $L_{p,q}(R)$  (see [4, 8, 43]) and if, in addition,  $p = q$  then we get the *Lebesgue spaces*  $L_p(R)$ .

In what follows, we work with couples of Lebesgue spaces  $(L_{p_0}(R), L_{p_1}(R))$ ,  $(L_{q_0}(S), L_{q_1}(S))$  and operators  $T \in \mathcal{L}((L_{p_0}(R), L_{p_1}(R)), (L_{q_0}(S), L_{q_1}(S)))$ . We put

$$\beta(T_j) = \beta(T : L_{p_j}(R) \rightarrow L_{q_j}(S)), \quad j = 0, 1.$$

It is shown in [29, Corollary 5.3] that

$$L_{p,q;b}(R) = (L_1(R), L_\infty(R))_{\theta,q;b} \tag{4.1}$$

provided that  $1 < p < \infty$ ,  $0 < \theta < 1$ ,  $1/p = 1 - \theta$ ,  $0 < q \leq \infty$  and  $b \in SV(0, \infty)$ .

As a consequence of Theorem 3.1 we can establish the following result for Lorentz–Karamata spaces.

**Theorem 4.1** *Let  $(R, \mu)$  and  $(S, \nu)$  be non-atomic  $\sigma$ -finite measure spaces. Let  $1 < p_0 < p_1 < \infty$ ,  $1 < q_0 < q_1 < \infty$ ,  $0 < q < \infty$  and let  $b \in SV(0, \infty)$  satisfying (2.8) and (2.12). Put*

$$b_0(t) = b(t^{1/p_0-1/p_1}) \left( \frac{1}{b(t^{1/p_0-1/p_1})^q} \int_t^\infty b(s^{1/p_0-1/p_1})^q \frac{ds}{s} \right)^{1/\min\{p_0, q\}}$$

and

$$b_1(t) = b(t^{1/q_0-1/q_1}) \left( \frac{1}{b(t^{1/q_0-1/q_1})^q} \int_t^\infty b(s^{1/q_0-1/q_1})^q \frac{ds}{s} \right)^{1/\max\{q_0, q\}}.$$

If  $T \in \mathcal{L}((L_{p_0}(R), L_{p_1}(R)), (L_{q_0}(S), L_{q_1}(S)))$  then

$$T : L_{p_0, q; b_0}(R) \rightarrow L_{q_0, q; b_1}(S) \text{ boundedly.}$$

Moreover, for  $\beta(T) = \beta(T : L_{p_0, q; b_0}(R) \rightarrow L_{q_0, q; b_1}(S))$  we have

- (a)  $\beta(T) = 0$  if  $\beta(T_0) = 0$ ,
- (b)  $\beta(T) \leq C\beta(T_0)$  if  $0 \leq \beta(T_1) < \beta(T_0)$ ,
- (c)  $\beta(T) \leq C \max \{ \beta(T_0), \beta(T_0)\bar{b}(\beta(T_0)/\beta(T_1)) \}$  if  $0 < \beta(T_0) \leq \beta(T_1)$ .

Here  $C > 0$  is a constant independent of  $T$ .

**Proof** Let  $0 < \theta_0 < \theta_1 < 1$  such that  $1/p_j = 1 - \theta_j$ ,  $j = 0, 1$ . We have  $L_{p_j}(R) = (L_1(R), L_\infty(R))_{\theta_j, p_j}$ . Hence, we can use the reiteration formula of [29, Theorem 3.2] to work with the space  $(L_{p_0}(R), L_{p_1}(R))_{0, q; b}$ . Then, according to [36, Theorem 4.10] and (4.1), we obtain

$$L_{p_0, q; b_0}(R) = (L_1(R), L_\infty(R))_{\theta_0, q; b_0} \hookrightarrow (L_{p_0}(R), L_{p_1}(R))_{0, q; b}.$$

Similarly, but using now [36, Theorem 4.8] with  $\eta_j = 1 - 1/q_j$ ,  $j = 0, 1$ , we get

$$(L_{q_0}(S), L_{q_1}(S))_{0, q; b} \hookrightarrow (L_1(S), L_\infty(S))_{\eta_0, q; b_1} = L_{q_0, q; b_1}(S).$$

Therefore, the result follows interpolating with parameters  $0, q, b$  the couples  $(L_{p_0}(R), L_{p_1}(R)), (L_{q_0}(S), L_{q_1}(S))$ , applying Theorem 3.1 and having in mind the embeddings pointed out above.  $\square$

Subsequently, for  $\tau \in \mathbb{R}$  and  $\mathbb{A} = (\alpha_0, \alpha_\infty) \in \mathbb{R}^2$ , we put  $\mathbb{A} + \tau = (\alpha_0 + \tau, \alpha_\infty + \tau)$ . Recall that  $\alpha^+ = \max\{\alpha, 0\}$  for  $\alpha \in \mathbb{R}$ .

**Remark 4.2** Let  $\mathbb{A} = (\alpha_0, \alpha_\infty) \in \mathbb{R}^2$  such that  $\alpha_\infty + 1/q < 0 < \alpha_0 + 1/q$  and let  $b(t) = (1 + |\log t|)^{\mathbb{A}}$ . Then for the function  $b_0$  in Theorem 4.1 we obtain

$$b_0(t) \approx b(t) \left( \frac{1}{b(t)^q} \int_t^\infty b(s)^q \frac{ds}{s} \right)^{1/\min\{p_0, q\}}$$

$$\begin{aligned} &\approx (1 + |\log t|)^{\mathbb{A}} (1 + |\log t|)^{1/\min\{p_0, q\}} \\ &= (1 + |\log t|)^{\mathbb{A} + \frac{1}{\min\{p_0, q\}}}. \end{aligned}$$

Similarly,

$$b_1(t) \approx (1 + |\log t|)^{\mathbb{A} + \frac{1}{\max\{q_0, q\}}}.$$

Hence, we have that

$$L_{p_0, q; b_0}(R) = L_{p_0, q}(\log L)_{\mathbb{A} + \frac{1}{\min\{p_0, q\}}}(R)$$

and

$$L_{q_0, q; b_1}(S) = L_{q_0, q}(\log L)_{\mathbb{A} + \frac{1}{\max\{q_0, q\}}}(S).$$

Moreover, by [14, Lemma 2.1] and [5, (2.6)], we have

$$\bar{b}(t) \leq (1 + |\log t|)^{(\alpha_0^+ - \alpha_\infty, (-\alpha_0)^+)}.$$

Consequently, writing down Theorem 4.1 for this choice of  $b$  we recover a result of Besoy and Cobos (see [5, Corollary 3.13]), which is a quantitative version of a compactness result of Edmunds and Opic (see [26, Corollary 4] and also [20, Corollary 4.5]).

The following result refers to the case  $1 < p_1 < p_0 < \infty$ .

**Theorem 4.3** *Let  $(R, \mu)$  and  $(S, \nu)$  be non-atomic  $\sigma$ -finite measure spaces. Let  $1 < p_1 < p_0 < \infty$ ,  $1 < q_0 < q_1 < \infty$ ,  $0 < q < \infty$  and let  $b \in SV(0, \infty)$  satisfying (2.8) and (2.12). Put*

$$\tilde{b}_0(t) = b(t^{1/p_0-1/p_1}) \left( \frac{1}{b(t^{1/p_0-1/p_1})^q} \int_0^t b(s^{1/p_0-1/p_1})^q \frac{ds}{s} \right)^{1/\min\{p_0, q\}}$$

and

$$b_1(t) = b(t^{1/q_0-1/q_1}) \left( \frac{1}{b(t^{1/q_0-1/q_1})^q} \int_t^\infty b(s^{1/q_0-1/q_1})^q \frac{ds}{s} \right)^{1/\max\{q_0, q\}}.$$

If  $T \in \mathcal{L}((L_{p_0}(R), L_{p_1}(R)), (L_{q_0}(S), L_{q_1}(S)))$  then

$$T : L_{p_0, q; \tilde{b}_0}(R) \rightarrow L_{q_0, q; b_1}(S) \text{ boundedly.}$$

Moreover, for  $\beta(T) = \beta(T : L_{p_0, q; \tilde{b}_0}(R) \rightarrow L_{q_0, q; b_1}(S))$  we have

(a)  $\beta(T) = 0$  if  $\beta(T_0) = 0$ ,

- (b)  $\beta(T) \leq C\beta(T_0)$  if  $0 \leq \beta(T_1) < \beta(T_0)$ ,
- (c)  $\beta(T) \leq C \max \{ \beta(T_0), \beta(T_0)\bar{b}(\beta(T_0)/\beta(T_1)) \}$  if  $0 < \beta(T_0) \leq \beta(T_1)$ .

Here  $C > 0$  is a constant independent of  $T$ .

**Proof** Consider the couple  $(L_\infty(R), L_1(R))$ . We have

$$L_{p_j}(R) = (L_\infty(R), L_1(R))_{\tilde{\theta}_j, p_j} \text{ where } \tilde{\theta}_j = \frac{1}{p_j}, j = 0, 1.$$

So  $0 < \tilde{\theta}_0 < \tilde{\theta}_1 < 1$  and we still can use [29, Theorem 3.2] and [36, Theorem 4.10] to get that

$$(L_\infty(R), L_1(R))_{\tilde{\theta}_0, q; u} \hookrightarrow (L_{p_0}(R), L_{p_1}(R))_{0, q; b}$$

where

$$u(t) = b(t^{1/p_1-1/p_0}) \left( \frac{1}{b(t^{1/p_1-1/p_0})^q} \int_t^\infty b(s^{1/p_1-1/p_0})^q \frac{ds}{s} \right)^{1/\min\{p_0, q\}}.$$

According to the relationship between the  $K$ -functionals of  $(L_\infty(R), L_1(R))$  and  $(L_1(R), L_\infty(R))$ , making a change of variables and using (4.1), we obtain

$$\begin{aligned} \|f\|_{(L_\infty(R), L_1(R))_{\tilde{\theta}_0, q; u}} &= \left( \int_0^\infty [t^{-\tilde{\theta}_0} u(t) K(t, f; L_\infty(R), L_1(R))]^q \frac{dt}{t} \right)^{1/q} \\ &= \left( \int_0^\infty [t^{1-\tilde{\theta}_0} \tilde{b}_0(t^{-1}) K(t^{-1}, f; L_1(R), L_\infty(R))]^q \frac{dt}{t} \right)^{1/q} \\ &= \left( \int_0^\infty [t^{\tilde{\theta}_0-1} \tilde{b}_0(t) K(t, f; L_1(R), L_\infty(R))]^q \frac{dt}{t} \right)^{1/q} \\ &= \|f\|_{(L_1(R), L_\infty(R))_{1-\tilde{\theta}_0, q; \tilde{b}_0}} \approx \|f\|_{L_{p_0, q; \tilde{b}_0}(R)}. \end{aligned}$$

Therefore

$$L_{p_0, q; \tilde{b}_0}(R) \hookrightarrow (L_{p_0}(R), L_{p_1}(R))_{0, q; b}.$$

Since the embedding

$$(L_{q_0}(S), L_{q_1}(S))_{0, q; b} \hookrightarrow L_{q_0, q; b_1}(S)$$

has been established in Theorem 4.1, we can conclude the result by interpolating with parameters  $0, q, b$  and applying Theorem 3.1.  $\square$

**Remark 4.4** If  $b(t) = (1 + |\log t|)^\mathbb{A}$  with  $\mathbb{A} = (\alpha_0, \alpha_\infty)$  and  $\alpha_\infty + 1/q < 0 < \alpha_0 + 1/q$ , then

$$\tilde{b}_0(t) \approx (1 + |\log t|)^{\mathbb{A} + \frac{1}{\min\{p_0, q\}}}$$

where  $\tilde{\mathbb{A}} = (\alpha_\infty, \alpha_0)$  and Theorem 4.3 gives estimates for the measure of non-compactness of

$$T : L_{p_0,q}(\log L)_{\tilde{\mathbb{A}} + \frac{1}{\min\{p_0,q\}}}(R) \rightarrow L_{q_0,q}(\log L)_{\mathbb{A} + \frac{1}{\max\{q_0,q\}}}(R).$$

Proceeding similarly, but using [36, Theorem 4.8], we can derive results for  $1 < q_1 < q_0 < \infty$ .

We finish the paper with some results when the main information on  $T$  refers to the restriction from  $L_{p_1}(R)$  into  $L_{q_1}(S)$ .

**Theorem 4.5** *Let  $(R, \mu)$  and  $(S, \nu)$  be non-atomic  $\sigma$ -finite measure spaces. Let  $1 < p_1 < p_0 < \infty, 1 < q_1 < q_0 < \infty, 0 < q < \infty$  and let  $v \in SV(0, \infty)$  satisfying that*

$$\left(\int_0^1 v(t)^q \frac{dt}{t}\right)^{1/q} < \infty \text{ and } \left(\int_1^\infty v(t)^q \frac{dt}{t}\right)^{1/q} = \infty.$$

Put

$$v_0(t) = v(t^{1/p_0-1/p_1}) \left(\frac{1}{v(t^{1/p_0-1/p_1})^q} \int_t^\infty v(s^{1/p_0-1/p_1})^q \frac{ds}{s}\right)^{1/\min\{p_1,q\}}$$

and

$$v_1(t) = v(t^{1/q_0-1/q_1}) \left(\frac{1}{v(t^{1/q_0-1/q_1})^q} \int_t^\infty v(s^{1/q_0-1/q_1})^q \frac{ds}{s}\right)^{1/\max\{q_1,q\}}.$$

If  $T \in \mathcal{L}((L_{p_0}(R), L_{p_1}(R)), (L_{q_0}(S), L_{q_1}(S)))$  then

$$T : L_{p_1,q;v_0}(R) \rightarrow L_{q_1,q;v_1}(S) \text{ boundedly.}$$

Furthermore, for  $\beta(T) = \beta(T : L_{p_1,q;v_0}(R) \rightarrow L_{q_1,q;v_1}(S))$  we have

- (a)  $\beta(T) = 0$  if  $\beta(T_1) = 0$ ,
- (b)  $\beta(T) \leq C\beta(T_1)$  if  $0 \leq \beta(T_0) < \beta(T_1)$ ,
- (c)  $\beta(T) \leq C \max\{\beta(T_1), \beta(T_1)\bar{\nu}(\beta(T_1)/\beta(T_0))\}$  if  $0 < \beta(T_1) \leq \beta(T_0)$ .

Here  $C > 0$  is a constant independent of  $T$ .

**Proof** According to (2.7), for any quasi-Banach couple  $(A_0, A_1)$  we have  $(A_0, A_1)_{1,q;v} = (A_1, A_0)_{0,q;b}$  where  $b(t) = v(1/t)$ . We also have that

$$T \in \mathcal{L}((L_{p_1}(R), L_{p_0}(R)), (L_{q_1}(S), L_{q_0}(S))).$$

Hence, the wanted result follows by interpolating with parameters  $0, q, b$  and applying Theorem 4.1. □

If  $1 < p_0 < p_1 < \infty$  and/or  $1 < q_0 < q_1 < \infty$  we can obtain similar results.

**Remark 4.6** Let  $\mathbb{A} = (\alpha_0, \alpha_\infty)$  with  $\alpha_0 + 1/q < 0 < \alpha_\infty + 1/q$  and let  $v(t) = (1 + |\log t|)^{\mathbb{A}}$ . So,  $v$  satisfies the assumptions of Theorem 4.5. We have

$$\begin{aligned} v_0(t) &\approx (1 + |\log t|)^{\tilde{\mathbb{A}} + \frac{1}{\min\{p_1, q\}}}, \\ v_1(t) &\approx (1 + |\log t|)^{\tilde{\mathbb{A}} + \frac{1}{\max\{q_1, q\}}}. \end{aligned}$$

Moreover, by [5, (2.6)] we know that  $\bar{v}(t) \leq (1 + |\log t|)^{((-\alpha_\infty)^+, \alpha_\infty^+ - \alpha_0)}$ . Writing down Theorem 4.5 for this choice of the parameters we obtain estimates for the measure of non-compactness of

$$T : L_{p_1, q}(\log L)_{\tilde{\mathbb{A}} + \frac{1}{\min\{p_1, q\}}}(R) \rightarrow L_{q_1, q}(\log L)_{\tilde{\mathbb{A}} + \frac{1}{\max\{q_1, q\}}}(R).$$

**Acknowledgements** Fernando Cobos and Luz M. Fernández-Cabrera have been supported in part by UCM Grant PR3/23-30811. Part of the research of Manvi Grover was done while she visited the Department of Mathematical Analysis and Applied Mathematics at Universidad Complutense de Madrid supported in part by the Grant CZ.02.2.69/0.0/0.0/19 – 073/0016935. She would like to thank the Department for its hospitality. The authors would like to thank the referee for his/her comments.

**Funding** Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

**Data availability** Not applicable.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Beauzamy, B.: Espaces d'interpolation réels: topologie et géométrie. Lecture Notes in Math, vol. 666. Springer, Berlin (1978)
2. Bennett, C., Rudnick, K.: On Lorentz-Zygmund spaces. *Dissertationes Math.* **175**, 1–72 (1980)
3. Bennett, C., Sharpley, R.: *Interpolation of Operators*. Academic Press, Boston (1988)
4. Bergh, J., Löfström, J.: *Interpolation Spaces. An introduction*. Springer, Berlin (1976)
5. Besoy, B.F., Cobos, F.: Logarithmic interpolation methods and measure of non-compactness. *Quart. J. Math.* **71**, 73–95 (2020)
6. Besoy, B.F., Cobos, F.: The equivalence theorem for logarithmic interpolation spaces in the quasi-Banach case. *Z. Anal. Anwendungen* **40**, 1–32 (2021)
7. Besoy, B.F., Cobos, F., Fernández-Cabrera, L.M.: On the structure of a limit class of logarithmic interpolation spaces. *Mediterr. J. Math.* **17**, 168 (2020)
8. Butzer, P.L., Berens, H.: *Semi-Groups of Operators and Approximation*. Springer, Berlin (1967)
9. Cobos, F.: Interpolation theory and compactness. In: Lukeš, J., Pick, L. (eds.) *Function Spaces, Inequalities and Interpolation*, pp. 31–75. Prague, Paseky (2009)
10. Cobos, F., Fernández-Cabrera, L.M., Kühn, T., Ullrich, T.: On an extreme class of real interpolation spaces. *J. Funct. Anal.* **256**, 2321–2366 (2009)
11. Cobos, F., Fernández-Cabrera, L.M., Martínez, A.: Abstract K and J spaces and measure of non-compactness. *Math. Nachr.* **280**, 1698–1708 (2007)

12. Cobos, F., Fernández-Cabrera, L.M., Martínez, A.: Measure of non-compactness of operators interpolated by limiting real methods. In: *Operator Theory: Advances and Applications*, 219, pp. 37–54. Springer, Basel (2012)
13. Cobos, F., Fernández-Cabrera, L.M., Martínez, A.: On a paper of Edmunds and Opic on limiting interpolation of compact operators between  $L_p$  spaces. *Math. Nachr.* **288**, 167–175 (2015)
14. Cobos, F., Fernández-Cabrera, L.M., Martínez, A.: Estimates for the spectrum on logarithmic interpolation spaces. *J. Math. Anal. Appl.* **437**, 292–309 (2016)
15. Cobos, F., Fernández-Martínez, P., Martínez, A.: Interpolation of the measure of non-compactness by the real method. *Studia Math.* **135**, 25–38 (1999)
16. Cobos, F., Kühn, T.: Equivalence of  $K$ - and  $J$ -methods for limiting real interpolation spaces. *J. Funct. Anal.* **261**, 3696–3722 (2011)
17. Cobos, F., Kühn, T., Schonbek, T.: One-sided compactness results for Aronszajn-Gagliardo functors. *J. Funct. Anal.* **106**, 274–313 (1992)
18. Cobos, F., Martínez, A.: Remarks on interpolation properties of the measure of weak non-compactness and ideal variations. *Math. Nachr.* **208**, 93–100 (1999)
19. Cobos, F., Peetre, J.: Interpolation of compactness using Aronszajn-Gagliardo functors. *Israel J. Math.* **68**, 220–240 (1989)
20. Cobos, F., Segurado, A.: Description of logarithmic interpolation spaces by means of the  $J$ -functional and applications. *J. Funct. Anal.* **268**, 2906–2945 (2015)
21. Cordeiro, J.M.: *Interpolación de Ciertas Clases de Operadores por Métodos Multidimensionales*, Ph. D. thesis, Publicaciones del Depto. de Matemática Aplicada, Universidad de Vigo (1999)
22. Cwikel, M.: Real and complex interpolation and extrapolation of compact operators. *Duke Math. J.* **65**, 333–343 (1992)
23. Davis, W.J., Figiel, T., Johnson, W.B., Pelczyński, A.: Factoring weakly compact operators. *J. Funct. Anal.* **17**, 311–327 (1974)
24. Edmunds, D.E., Evans, W.D.: *Spectral Theory and Differential Operators*. Clarendon Press, Oxford (1987)
25. Edmunds, D.E., Evans, W.D.: *Hardy Operators. Function Spaces and Embeddings*. Springer, Berlin (2004)
26. Edmunds, D.E., Opic, B.: Limiting variants of Krasnosel’skiĭ’s compact interpolation theorem. *J. Funct. Anal.* **266**, 3265–3285 (2014)
27. Evans, W.D., Opic, B.: Real interpolation with logarithmic functors and reiteration. *Can. J. Math.* **52**, 920–960 (2000)
28. Evans, W.D., Opic, B., Pick, L.: Real interpolation with logarithmic functors. *J. Inequal. Appl.* **7**, 187–269 (2002)
29. Gogatishvili, A., Opic, B., Trebels, W.: Limiting reiteration for real interpolation with slowly varying functions. *Math. Nachr.* **278**, 86–107 (2005)
30. Gustavsson, J.: A function parameter in connection with interpolation of Banach spaces. *Math. Scand.* **42**, 289–305 (1978)
31. Heinrich, S.: Closed operator ideals and interpolation. *J. Funct. Anal.* **35**, 397–411 (1980)
32. König, H.: *Eigenvalue Distribution of Compact Operators*. Birkhäuser, Basel (1986)
33. Köthe, G.: *Topological Vector Spaces I*. Springer, Berlin (1969)
34. Maligranda, L., Quevedo, A.: Interpolation of weakly compact operators. *Arch. Math.* **55**, 280–284 (1990)
35. Mastysłó, M.: On interpolation of weakly compact operators. *Hokkaido Math. J.* **22**, 105–114 (1993)
36. Neves, J.S., Opic, B.: Optimal local embeddings of Besov spaces involving only slowly varying smoothness. *J. Approx. Theory* **254**, 105393 (2020)
37. Nilsson, P.: Reiteration theorems for real interpolation and approximation spaces. *Ann. Mat. Pura Appl.* **132**, 291–330 (1982)
38. Opic, B., Grover, M.: Description of  $K$ -spaces by means of  $J$ -spaces and the reverse problem in the limiting real interpolation. *Math. Nachr.* **296**, 4002–4031 (2023)
39. Opic, B., Pick, L.: On generalized Lorentz-Zygmund spaces. *Math. Inequal. Appl.* **2**, 391–467 (1999)
40. Persson, L.E.: Interpolation with a parameter function. *Math. Scand.* **59**, 199–222 (1986)
41. Szwedek, R.: Measure of non-compactness of operators interpolated by the real method. *Studia Math.* **175**, 157–174 (2006)
42. Teixeira, M.F., Edmunds, D.E.: Interpolation theory and measures of non-compactness. *Math. Nachr.* **104**, 129–135 (1981)

- 
43. Triebel, H.: Interpolation Theory. Function Spaces, Differential Operators. North-Holland, Amsterdam (1978)
  44. Triebel, H.: Theory of Function Spaces II. Birkhäuser, Basel (1992)