



An Early Jurassic (Sinemurian–Toarcian) stratigraphic framework for the occurrence of Organic Matter Preservation Intervals (OMPIs)

R.L. Silva^{a,b,*}, L.V. Duarte^c, G.D. Wach^b, M. Ruhl^a, D. Sadki^d, J.J. Gómez^e, S.P. Hesselbo^f, Weimu Xu^g, D. O'Connor^b, B. Rodrigues^c, J.G. Mendonça Filho^h

^a Department of Geology, School of Natural Sciences, Irish Centre for Research in Applied Geosciences (iCRAG), Trinity College Dublin, The University of Dublin, Ireland

^b Basin and Reservoir Lab, Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Nova Scotia, Canada

^c University of Coimbra & MARE, Department of Earth Sciences, Faculty of Science, Coimbra, Portugal

^d Department of Geology, Faculty of Science, Moulay Ismail University of Meknes, Morocco

^e Geodynamic, Stratigraphy and Palaeontology Department, Geological Sciences Faculty, University Complutense Madrid, and IGEO (UCM-CSIC), Madrid, Spain

^f Camborne School of Mines & Environment and Sustainability Institute, University of Exeter, Penryn, UK

^g School of Earth Sciences & Irish Centre for Research in Applied Geosciences (iCRAG), University College Dublin, Belfield, Dublin 4, Ireland

^h Geosciences Department & LAFO, Federal Univ Rio de Janeiro, Rio de Janeiro, Brazil

ARTICLE INFO

Keywords:

Organic-rich facies
Stable carbon isotopes
Sinemurian
Pliensbachian
Toarcian Oceanic Anoxic Event
Early Jurassic

ABSTRACT

Lower Jurassic sedimentary successions in the Atlantic margin basins include several organic-rich intervals, some with source rock potential; time-equivalent units are also identified in on- and offshore areas worldwide. Despite decades of research, it is still unclear which mechanisms lead to the deposition of organic-rich sediments during the Early Jurassic. The objectives of this study are to construct a detailed temporal and geographical framework of Sinemurian–Toarcian organic matter preservation intervals (OMPIs; subdivided into local, regional, and super-regional) and roughly constrain the relationship of OMPIs with the Lower Jurassic $\delta^{13}\text{C}$ record. This survey combines an in-depth analysis of literature on the distribution of organic-rich facies in the Sinemurian–Toarcian with new geochemical studies [total organic carbon (TOC) and organic matter pyrolysis] from selected outcrop sections from Portugal, Spain, and Morocco.

Strong local control on OMPIs during most of the Sinemurian is suggested. Regionally widespread organic-rich facies are associated with the most negative $\delta^{13}\text{C}$ values of the broad Sinemurian–Pliensbachian negative carbon isotopic trend recorded in organic matter (including the Sinemurian–Pliensbachian Boundary Event). Pliensbachian OMPIs are expressive in the areas bordering the proto-Atlantic Ocean and are often linked with positive $\delta^{13}\text{C}$ excursions and short-lived warm intervals, but OMPIs are also defined for the Late Pliensbachian cool interval. Early Toarcian superregional OMPIs are associated with some of the most pronounced $\delta^{13}\text{C}$ excursions of the Mesozoic. Toarcian maximum TOC content occurs with the positive $\delta^{13}\text{C}$ (recovery) trend following the $\delta^{13}\text{C}$ negative shift typically linked with the Early Toarcian Oceanic Anoxic Event (T-OAE), supporting the notion that peak carbon sequestration/ocean anoxia post-dated the main phase of carbon input into the atmosphere, as also suggested by recent modelling efforts. However, additional superregional OMPIs predate and postdate the T-OAE, indicating that conditions favouring preservation of organic matter (increased productivity and/or enhanced preservation) during the Early Toarcian were not restricted to the T-OAE interval.

The compilation of Sinemurian–Toarcian OMPIs presented in this paper demonstrates that organic-rich intervals of regional and superregional expression in the Lower Jurassic sedimentary record are ubiquitous and may even be more numerous than in the Cretaceous. Considering the association of some of the Sinemurian, Pliensbachian, and Toarcian (not taking into account the T-OAE related OMPIs) regional and superregional OMPIs with well-defined carbon isotopic excursions, it is here suggested that these hold the same relevance as the secondary OAEs of the Cretaceous, such as the Valanginian OAE (Weissert Event), Hauterivian OAE (Faraonin Event), and Late Aptian–Early Albian OAE (OAE 1b cluster).

* Corresponding author at: Department of Geology, School of Natural Sciences, Museum Building, Trinity College Dublin, 1 Park Ln E, Dublin 2, Ireland.
E-mail address: ricardo.cdv@gmail.com (R.L. Silva).

1. Introduction

The deposition and preservation of organic matter-rich sediments during the Early Jurassic has led to the formation of prolific hydrocarbon source-rocks in many locations worldwide (e.g. Powell, 1978; Hallam and Bradshaw, 1979; Fleet et al., 1987; Jenkyns, 1988; Kodina et al., 1988; Baudin et al., 1990; Baudin, 1995; Bessereau et al., 1995; Scotchman, 2001; van de Schootbrugge et al., 2005b; Duarte et al., 2010; Silva et al., 2011, 2017; Suan et al., 2011; Silva and Duarte, 2015; Ruhl et al., 2016; Gómez et al., 2016a; Xu et al., 2017a; Campana et al., 2017). These organic-rich sedimentary rocks are interpreted to be near age equivalent between sedimentary basins and often associated with significant perturbations to the global carbon cycle and changes in regional/global climates and Earth system processes (e.g. Jenkyns and Clayton, 1986, 1997; Jenkyns, 2010; Jenkyns, 1988; van Buchem et al.,

1995; Hesselbo et al., 2000a, 2007; van de Schootbrugge et al., 2005a, 2013; Suan et al., 2008, 2010, 2011, 2015; Dera et al., 2010; Korte and Hesselbo, 2011; Silva et al., 2011, 2015, 2017; Gómez and Goy, 2011; Jenkyns and Weedon, 2013; Franceschi et al., 2014; Hermoso et al., 2014; Korte et al., 2015; Silva and Duarte, 2015; Percival et al., 2016; Ruhl et al., 2016; Gómez et al., 2016b; Them et al., 2017; Xu et al., 2017b; Xu et al., 2017a, 2018; Ruebsam et al., 2018, 2020; Danisch et al., 2019; Danise et al., 2019; Slater et al., 2019; Mercuzot et al., 2020; Schöllhorn et al., 2020b; Schöllhorn et al., 2020a; Storm et al., 2020) (Figs. 1 and 2).

Some of the most remarkable Early Jurassic intervals of enhanced preservation of organic matter (OM), such as the Early Toarcian Oceanic Anoxic Event (T-OAE), are being intensively investigated to constrain the mechanisms and feedback processes controlling oceanographic, biological, and climatic changes during important past global




Fig. 1. Base map with Early Jurassic continent positions and locations of basins and localities with known Lower Jurassic sediments (modified from Damborenea (2002) and references therein). For simplicity, the West Netherlands Basin, Southern Beaujolais, Sub-Briançonnais Basin, Northern Calcareous Alps, and Midlands shelf are not represented here.




Fig. 2. General temporal and paleogeographical trends of Lower Jurassic organic-rich sedimentary successions. *Western North America (NA) basins*: Stratigraphy - (Asgar-Deen et al., 2003); Organic-rich facies - (Caruthers et al., 2011; Porter et al., 2014b; Porter et al., 2014a; Them et al., 2017); *Arctic Basins*: Stratigraphy - (Shurygin et al., 2011); Organic-rich facies - NE (Suan et al., 2011); NW (Kontorovich et al., 1997); *Hebrides Basin (Scotland)*: Stratigraphy - (Hesselbo et al., 1998; Hesselbo and Coe, 2000; Morton, 2004); Organic-rich facies - (Thrasher, 1992; Trueblood, 1992; Scotchman, 2001); *Cleveland Basin (UK)*: Stratigraphy - (Hesselbo et al., 2000b; Simms et al., 2004; Korte and Hesselbo, 2011); Organic-rich facies - (Jenkyns and Clayton, 1997; Scotchman, 2001; Kemp et al., 2005; Wignall et al., 2005; McArthur et al., 2008; Littler et al., 2010; Caswell and Coe, 2012; Salem, 2013; Porter et al., 2014a; Song et al., 2015); *Slyne Basin (offshore Ireland)*: Stratigraphy - (Dancer et al., 2005); Organic-rich facies - (Trueblood, 1992; Scotchman, 2001; Silva et al., 2017); *Wessex Basin (UK)*: Stratigraphy - (Simms, 2004; Gallois, 2008; Jenkyns and Weedon, 2013); Organic-rich facies - (Jenkyns and Clayton, 1997; Weedon and Jenkyns, 1999; Scotchman, 2001; Deconinck et al., 2003; Ruhl et al., 2010; Jenkyns and Weedon, 2013; Price et al., 2016; Xu et al., 2017a); *Paris Basin (France)*: Stratigraphy - (Mouterde et al., 1980); Organic-rich facies - (Bessereau et al., 1995; van Breugel et al., 2006; Hermoso et al., 2012, 2013, 2014; Lézin et al., 2013; Song et al., 2014); *SW Germany*: Stratigraphy - (Menning and Hendrich, 2002; Pieńkowski et al., 2008; Feist-Burkhardt and Pross, 2010); Organic-rich facies - (Röhl et al., 2001; Schmid-Röhl et al., 2002; Röhl and Schmid-Röhl, 2005); data in (McArthur et al., 2008); (Song et al., 2015); *Asturian Basin (Spain)*: Stratigraphy - (Aurell et al., 2003; Comas-Rengifo and Goy, 2010); Organic-rich facies - (Borrego et al., 1996; Gómez et al., 2008, 2016a; Gómez and Goy, 2011); this study; *Basque-Cantabrian Basin (Spain)*: Stratigraphy - (Aurell et al., 2003); Organic-rich facies - (Quesada et al., 1997, 2005; Gómez and Goy, 2011); *Lusitanian Basin (Portugal)*: Stratigraphy - (Duarte and Soares, 2002; Duarte et al., 2010, 2014; Silva et al., 2011, 2015); Organic-rich facies - (de Oliveira et al., 2006; Duarte et al., 2010, 2012; Silva et al., 2010, 2012; Silva et al., 2011, 2012; Poças Ribeiro et al., 2013; Silva, 2015; Silva and Duarte, 2015); this study; *Middle Atlas (Morocco)* Stratigraphy - (Sadki et al., 2014); Organic-rich facies - (Rachidi et al., 2009; Sachse et al., 2012); this study; *High Atlas (Morocco)* Stratigraphy - (Sadki et al., 2014); Organic-rich facies - (Bodin et al., 2010, 2011, 2016); this study; *Tunisia*: Stratigraphy - (Soussi, 2003); Organic-rich facies - (Sousa, 2014); *Umbria Marche Basin (Italy)*: Stratigraphy - (Monaco et al., 1994; Parisi et al., 1996; Rodríguez-Tovar et al., 2016); Organic-rich facies - (Monaco et al., 1994; Parisi et al., 1996; Bucefalo Palliani et al., 1998; Pancost et al., 2004; Sabatino et al., 2009).

environmental perturbations (e.g. Küspert, 1982; Jenkyns and Clayton, 1986; Jenkyns, 1988, 2010; Hesselbo et al., 2000a, 2007; Röhl et al., 2001; Jenkyns et al., 2002; Cohen et al., 2004; Röhl and Schmid-Röhl, 2005; Hermoso et al., 2012, 2013; Hermoso et al., 2009; Al-Suwaidi et al., 2010, 2016; Bodin et al., 2010; Littler et al., 2010; Bodin et al., 2016; Suan et al., 2010, 2011, 2015; Lézin et al., 2013; Sabatino et al., 2013; Kafousia et al., 2014; Kemp and Izumi, 2014; Reolid et al., 2014; Caruthers et al., 2014; Pieńkowski et al., 2016; Martinez et al., 2017; Silva et al., 2017; Them et al., 2017; Xu et al., 2017b, 2018; Ruvalcaba Baroni et al., 2018; Fantasia et al., 2018, 2019b; Ruesam et al., 2019;

Danise et al., 2019; Remírez and Algeo, 2020; Schöllhorn et al., 2020a; Storm et al., 2020; Rodrigues et al., 2021). However, there is increasing evidence that organic carbon burial and preservation, even when associated with widespread carbon sequestration events such as the T-OAE, were modulated by local geography, bathymetry, climate, ocean chemistry, nutrient supply, biological communities and, consequently, the local depositional environment (e.g. Duarte et al., 2010, 2014; Silva et al., 2012, 2013, 2015, 2017, 2021; Reolid et al., 2014; Suan et al., 2015; Rodrigues et al., 2016, 2019, 2020b; Bruneau et al., 2017; Them et al., 2017; Martindale et al., 2017; Rodríguez-Tovar et al., 2017;

Ruvalcaba Baroni et al., 2018; Danise et al., 2019) (Fig. 2).

With over 40 years of research on the understanding of the processes leading to Early Jurassic OM enrichment in marine and terrestrial environments, there are still large uncertainties concerning the impact of local versus global factors leading to OM enrichment. Uncertainty arises from the (1) temporal and spatial ambiguity when correlating most Lower Jurassic organic-rich sedimentary successions, (2) limited constraints on the amount, chemical composition, and thermal maturity of OM, and (3) incomplete understanding of environmental and depositional constraints on OM production and preservation at a wide range of geological time intervals. Understanding the link between OM production and preservation/sequestration and climate change is vital to advancing knowledge of processes and feedback mechanisms in the Earth Systems that drive global change at a wide range of time scales and to constrain the relative importance of the many factors (including natural and anthropogenic) contributing to modern-day climate change.

This study focuses on the temporal and spatial occurrence of organic-rich intervals through the Sinemurian–Toarcian interval, comprising most of the Lower Jurassic. The two objectives of this paper are to (1) construct a detailed temporal and geographical framework of Sinemurian–Toarcian organic-rich facies occurrences at a global scale, defined here as organic matter preservation intervals (OMPIs) and (2) approximately constrain the relationship of OMPIs with the Lower Jurassic $\delta^{13}\text{C}$ record. This study combines an in-depth analysis of literature on the distribution and characteristics of OM (quantity of carbon) in the Sinemurian–Toarcian with new geochemical studies from selected outcrop sections from Portugal, Spain, and Morocco.

1.1. Production and preservation of organic matter and its impact on the global carbon cycle

For convenience, the carbon cycle is usually divided into two cycles, the short- and long-term carbon cycles (Berner, 1999). Carbon in the short-term carbon cycle is rapidly exchanged within the surficial reservoirs, consisting of the oceans, atmosphere, biosphere, and soil, whereas carbon in the long-term cycle is slowly exchanged between the geosphere and the surficial systems (Berner, 1999; Ciais et al., 2013). The short-term carbon cycle is the dominant control on atmospheric $p\text{CO}_2$ over millennia, whereas the long-term carbon cycle controls atmospheric CO_2 and O_2 over millions of years (Berner, 1999, 2006).

Equilibrium and kinetic fractionation among OM, CO_2 , and HCO_3^- control the carbon isotope ratios of atmospheric CO_2 and oceanic inorganic and organic carbon (e.g. Emerson and Hedges, 2008; Silva et al., 2020a). The main processes that affect the $^{13}\text{C}/^{12}\text{C}$ ratio of the “superficial” carbon reservoirs at geological time intervals are the (1) equilibrium between dissolved inorganic carbon (DIC) and atmospheric CO_2 , (2) fractionation between DIC and carbonate minerals, (3) fixation of CO_2 and production of biomass via photosynthesis, and (4) respiration, remineralisation, preservation, and sedimentary reworking and resuspension of sedimentary OM (Hayes et al., 1999).

There is little isotopic discrimination effect during respiration. However, through photosynthesis in the superficial layers of the ocean, ^{12}C is taken up preferentially over ^{13}C , leaving the superficial DIC pool enriched in ^{13}C . As OM sinks into intermediate and deep waters, “sinking” organic carbon is remineralised, and the deeper ocean DIC pool becomes enriched in ^{12}C . At geological time scales, removal of isotopically “light” OM into the sedimentary OM reservoir generates a relative enrichment in ^{13}C of dissolved CO_2 in seawater and continuation of this process leads to an overall increase in ^{13}C of marine carbonate, atmospheric CO_2 , and OM (e.g. Jenkyns and Clayton, 1986; Wefer and Berger, 1991; Marshall, 1992; Patterson and Walter, 1994; Mitchell et al., 1996; Weissert et al., 1998; Holmden et al., 1998; Kump and Arthur, 1999; Hesselbo et al., 2000b; Berner, 2003; Immenhauser et al., 2003; Swart and Eberli, 2005; Panchuk et al., 2006; Newton and Bottrell, 2007; Emerson and Hedges, 2008; Silva et al., 2011, 2020a).

Removal of isotopically light carbon from the oceans through OM

sequestration into the geological archive is recorded as an increase in $\delta^{13}\text{C}$ of contemporaneous carbonates and marine and terrestrial OM (Berner, 2003). Positive $\delta^{13}\text{C}$ excursions are often interpreted to represent enhanced burial and sequestration of OM into the sedimentary realm (e.g. Schlanger and Jenkyns, 1976; Jenkyns, 1988, 2010; Arthur et al., 1990; Kump and Arthur, 1999; Silva et al., 2011, 2020a). In particular cases, such as the T-OAE, the $\delta^{13}\text{C}$ record may respond in a more complex way if deposition of OM is accompanied by large-scale carbon release from internal or external sources with different isotopic composition (e.g. Küspert, 1982; Jenkyns and Clayton, 1986; Jenkyns, 1988, 2010; Hesselbo et al., 2000a, 2007; Pálffy and Smith, 2000; Röhl et al., 2001; Jenkyns et al., 2002; Cohen et al., 2004; Röhl and Schmid-Röhl, 2005; Hermoso et al., 2012, 2013; Hermoso et al., 2009; Al-Suwaidi et al., 2010, 2016; Bodin et al., 2010; Littler et al., 2010; Bodin et al., 2016; Suan et al., 2010, 2011, 2015; Lézini et al., 2013; Sabatino et al., 2013; Kafousia et al., 2014; Kemp and Izumi, 2014; Reolid et al., 2014; Caruthers et al., 2014; Silva and Duarte, 2015; Pieńkowski et al., 2016; Martinez et al., 2017; Silva et al., 2017, 2020a; Them et al., 2017; Xu et al., 2017b, 2018; Fantasia et al., 2018, 2019b; Rodrigues et al., 2019, 2020b; Ruebsam et al., 2019; Danise et al., 2019; Storm et al., 2020; Ullmann et al., 2020).

Two primary debates exist today regarding the processes and mechanisms leading to the occurrence and preservation of OM in sediments (Forsman and Hunt, 1958; Huc and Durand, 1974; Pedersen and Calvert, 1990; Largeau and Derenne, 1993; Tyson, 1995; van Buchem et al., 1995; Vandenbroucke and Largeau, 2007; Zonneveld et al., 2010; Jenkyns, 2010; Suárez-Ruiz et al., 2012; Silva et al., 2012, 2013, 2020a; Hemingway et al., 2019; Hülse et al., 2019). The first debate concerns the role of primary productivity and associated carbon flux from surface waters to the sediment-water interface (Pedersen and Calvert, 1990) vs marine anoxia and associated enhanced OM preservation (Demaison and Moore, 1980; Tyson, 1995) in controlling the stratigraphic and geographic occurrence of organic-rich facies; the second relates to the mechanisms leading to the preservation and transformation of OM at geological time scales: degradation/recondensation (Huc and Durand, 1974; Tissot and Welte, 1978, 1984); selective preservation (Philip and Calvin, 1976; Tegelaar et al., 1989; Largeau and Derenne, 1993; Silva et al., 2012); sorptive mineral protection (Bishop et al., 1992; Mayer, 1994; Hedges and Keil, 1995; Kennedy et al., 2002; Burdige, 2007; Kennedy and Wagner, 2011), OM sulfurization (Sinninghe Damsté et al., 1988; Adam et al., 1998), and OM-mineral interaction (van Buchem et al., 1995; Kennedy et al., 2002; Hemingway et al., 2019). It is widely acknowledged that anoxia and enhanced productivity can both drive the occurrence of organic-rich facies. However, organic-rich sedimentary rocks characterized by lipid-rich OM with elevated Hydrogen Index (HI) seems to require low availability of O_2 in bottom waters (e.g. Espitalié et al., 1977, 1985; Demaison and Moore, 1980; Tissot and Welte, 1984; Kohnen et al., 1990, 1992; Goodarzi et al., 1993; Schouten et al., 1994; Sinninghe Damsté et al., 1995; van Buchem et al., 1995; Rodger Harvey et al., 1995; Sun et al., 2002; Farrimond et al., 2003; Lee et al., 2004; Bowden et al., 2006; Zonneveld et al., 2010; Silva et al., 2012).

2. Methodology: a stratigraphic framework for Lower Jurassic organic-rich facies

The temporal and spatial occurrence of organic-rich sedimentary successions of Sinemurian–Toarcian age is reviewed by combining published organic geochemical data with new biostratigraphic and geochemical data (total organic carbon, TOC, and organic matter pyrolysis) from Lower Jurassic outcrops from Portugal, Spain, and Morocco (Appendixes 1–4). Some geographic areas are underrepresented in terms of data availability and warrant further studies to better constrain the extent and duration of Early Jurassic organic-rich deposition; a brief review of these locations is presented in Appendix 1 (A1.4. Uncertain data and data underrepresentation). A reference framework for the temporal and spatial occurrence of individual Early Jurassic

Organic Matter Preservation Intervals (OMPIs) is presented in Figs. 3 and 4 and Appendixes 2–4.

2.1. Organic-rich sedimentary rocks

Organic-rich sedimentary rocks are here defined by present-day TOC values equal or higher than 2%. This cut-off value was chosen because:




Fig. 3. Geographic distribution of basins with record of organic-rich facies per OMPI. See Fig. 1 for basin identification and Table 1 and Appendix 2 for OMPI identification.




Fig. 3. (continued).

a) Sedimentary rocks with TOC > 2% may appear slightly darker compared to low-TOC rocks, making them more easily recognisable in the field or in hand samples (see also Hunt, 1995) and, therefore, from literature where geochemical data are not available and rocks are only described as organic-rich or black shales (there are,

however, other factors that may also result in dark mudstones, for example, mineralogical composition). Black shale is a common classification in the literature and, in its broadest definition, corresponds to a dark coloured, usually laminated mudstone, calcareous




Fig. 3. (continued).

 mudstone, or marl with TOC ranging from 1 to 30 wt% (e.g. Swanson, 1961; Weissert, 1981).

 Welte, 1984; Jones, 1987; Lewan, 1987; Chinn, 1991; Katz, 1995; Law, 1999; Peters et al., 2005).

b) This value roughly corresponds to the average TOC of “productive” hydrocarbon source rocks (see discussion in Ronov, 1958; Tissot and




Fig. 4. $\delta^{13}\text{C}_{\text{TOC}}$, TOC, and projection of OMPs for the Sinemurian–Toarcian at the reference Llanbedr (Mochras Farm) core, Cardigan Bay Basin (modified from Xu et al., 2018; Storm et al., 2020). The dark grey bars refer to OMPs observed at Mochras whereas the light grey bars reflect the approximate position of OMPs against the $\delta^{13}\text{C}_{\text{TOC}}$ curve. Blue line = seven-point moving average. Black squares = samples taken from core slabs; white squares = samples taken from reserve bags (~1.4-m intervals) of broken core. Orange circles = $\delta^{13}\text{C}_{\text{wood}}$. Depth of the samples from reserve bags refers to midpoint of the sample interval. Ammonite biostratigraphy after (Ivimey-Cook, 1971; Copestake and Johnson, 2013). Loc – Local; reg – regional; superregional; S – Sinemurian; P – Pliensbachian; LP – Late Pliensbachian; sp – spinatum; P-TOA – Pliensbachian–Toarcian; T-OAE – Toarcian oceanic anoxic Event; T – Toarcian; pCIE – positive carbon isotopic excursion; nCIE – negative carbon isotopic excursion; OMPi – organic matter preservation interval; F – fault.

2.2. Organic Matter Preservation Interval (OMPI)

An Organic Matter Preservation Interval (OMPI) corresponds to a constrained time interval (at the Subchron level) characterized by the deposition of organic-rich sediments (organic-rich sedimentary rocks with present day TOC > 2%) at local, regional, or superregional scales (as defined in subsection 2.1). OMPs are defined irrespective of kerogen type and classification, location, and processes of OM production, accumulation, and diagenesis/maturation. They are classified as local, regional, or superregional (Table 1, Figs. 3 and 4 and Appendix 2–4):

a) Local OMPI: Organic-rich sedimentary rocks observed in one section.

b) Regional OMPI: Time-equivalent (at Subchronozone scale) organic-rich sedimentary rocks observed in two or more sections separated by more than ~1000 km (arbitrarily defined) or located in two “non-contiguous” basins.

c) Superregional OMPI: Time-equivalent organic-rich sedimentary rocks in two or more non-contiguous paleogeographical domains. Examples of paleogeographical domains are Northern Europe, Tethys, Mediterranean, Central Atlantic, Arctic, Pacific North America, Pacific South America, Asia.

OMPis are described by first indicating the Age (Subage) where these are defined, followed by Chron, Subchron, and current understanding of

Table 1
Early Jurassic OMPIs - abbreviations and geographic distribution (see Appendix 2 for references).

| Abbreviation | OMPI | Known geographic distribution |
|---------------|---|-------------------------------|
| OMPI T10 | Late Toarcian (<i>levesquei-aalensis</i>) | Local (regional?) |
| OMPI T9 | Late Toarcian (<i>thouarsense</i> ,?) | Local |
| OMPI T8 | Middle Toarcian (<i>variabilis</i> ,?) | Local (regional?) |
| OMPI T7 | Middle Toarcian (<i>bifrons, fibulatum</i>) | Regional (superregional?) |
| OMPI T6 | Middle Toarcian (<i>bifrons, commune</i>) | Superregional |
| OMPI T5 | Early Toarcian (<i>falciferum, elegans-falciferum</i>) | Superregional |
| OMPI T4 (OAE) | Early T-OAE + nCIE (positive trend of the T-OAE nCIE) | Superregional |
| OMPI T3 (OAE) | Early T-OAE -nCIE (negative trend of the TOAE nCIE) | Superregional |
| OMPI T2 (OAE) | Early T-OAE pre nCIE | Superregional |
| OMPI T1 | Early Toarcian (<i>tenuicostatum, paltum-tenuicostatum</i>) | Superregional |
| OMPI P-Toa | Late Pliensbachian-Toarcian Event | Superregional |
| OMPI P11 | Late Pliensbachian (<i>spinatum, hawskerense</i>) | Superregional |
| OMPI P10 | Late Pliensbachian (<i>margaritatus, gibbosus</i>) | Local (superregional?) |
| OMPI P9 | Late Pliensbachian (<i>margaritatus, subnodosus/gibbosus</i>) | Regional (superregional?) |
| OMPI P8 | Late Pliensbachian (<i>margaritatus, subnodosus</i>) | Regional (superregional?) |
| OMPI P7 | Late Pliensbachian (<i>margaritatus, stokesi</i>) | Regional (superregional?) |
| OMPI P6 | Early-Late Pliensbachian (<i>davoei, figulinum-margaritatus, stokesi</i>) | regional (superregional?) |
| OMPI P5 | Early Pliensbachian (<i>davoei, maculatum-capricornus</i>) | Regional (superregional?) |
| OMPI P4 | Early Pliensbachian (<i>ibex, luridum</i>) | Local (superregional?) |
| OMPI P3 | Early Pliensbachian (<i>ibex, valdani</i>) | Regional (superregional?) |
| OMPI P2 | Early Pliensbachian (<i>jamesoni, jamesoni-ibex, masseanum</i>) | Regional (superregional?) |
| OMPI P1 | Early Pliensbachian (<i>jamesoni, brevespina</i>) | Regional (superregional?) |
| OMPI S-P | Sinemurian-Pliensbachian (<i>raricostatum, raricostatoides-jamesoni, polymorphus</i>) (inc. S-PBE cf. Korte and Hesselbo, 2011) | Regional (superregional?) |
| OMPI S7 | Late Sinemurian (base <i>raricostatum, densinodulum</i> ?) | Regional |
| OMPI S6 | Late Sinemurian (<i>oxynotum, oxynotum</i>) | Local (regional?) |
| OMPI S5 | Late Sinemurian (<i>obtusum, obtusum-stellare</i>) | Regional (superregional?) |
| OMPI S4 | Early Sinemurian (<i>turneri, birchii</i>) | Local (regional?) |
| OMPI S3 | Early Sinemurian (<i>semicostatum, resupinatum-turneri, birchii</i>) | Local (superregional?) |
| OMPI S2 | Early Sinemurian (<i>bucklandi/semicostatum</i>) | Local |
| OMPI S1 | Early Sinemurian (<i>bucklandi, rotiforme</i>) | Local |

the extent of the associated organic-rich facies, e.g. Early Pliensbachian (*davoei, maculatum/capricornus*) regional OMPI (see Table 1 and Appendix 2 for abbreviations for each OMPI). Uncertainty regarding the spatial representation of an OMPI is noted first by indicating the confirmed and then possible spatial classification, e.g., regional (superregional?). In particular cases, we adopt a historical denomination of a recognized feature of the geological record intimately associated with the OMPI, such as the Toarcian Oceanic Anoxic Event [e.g., OMPI T2 (OAE)]. The minimum requirement to define an OMPI is a single sample, representative of a continuous bed at outcrop scale, with present day TOC > 2% and a confident stratigraphic age assignment (using, for example, bio- or chemostratigraphy) to a Subchronozone. Organic-rich sedimentary rocks that cannot be confidently assigned to a Subchronozone cannot be used to define an OMPI. However, such loosely dated occurrences may support observations from other better-dated sections, as is the case for the Sinemurian successions of Western North American basins.

3. OMPIs: temporal and paleogeographical trends

Because $\delta^{13}\text{C}$ records exist for many of the Lower Jurassic sedimentary successions showing organic matter enrichment, below we briefly discuss the stratigraphic occurrence of OMPIs within the context of the Lower Jurassic $\delta^{13}\text{C}$ record (Table 1, Figs. 1–4, and Appendixes 1–4). A detailed discussion on processes leading to Early Jurassic short- and long-term global carbon cycle perturbations is beyond the scope of this study.

3.1. Early Sinemurian–Late Pliensbachian

3.1.1. Sinemurian–Early Pliensbachian: OMPIs S1–S7, OMPI S-P and OMPIs P1–P6

The Early Sinemurian to Early Pliensbachian interval is marked by multiple shifts of up to around 5‰ in the carbon isotopic composition of organic and inorganic substrates from marine and continental environments (e.g. Jenkyns et al., 2002; van de Schootbrugge et al., 2005a; Suan et al., 2010; Silva et al., 2011; Jenkyns and Weedon, 2013; Duarte et al., 2014; Price et al., 2016; Ruhl et al., 2016; Xu et al., 2017a; Hesselbo et al., 2020; Schöhlhorn et al., 2020b; Storm et al., 2020). Organic matter deposition and preservation in the Sinemurian seems to be strongly dependent on local depositional and tectonic conditions (Chadwick, 1986; Stapel et al., 1996; Rasmussen et al., 1998; Alves et al., 2002; Duarte et al., 2010, 2012, 2014; Poças Ribeiro et al., 2013; Silva et al., 2013), controlling, for example, relative sea-level and organic productivity (Bessereau et al., 1995; Jenkyns and Weedon, 2013; Schöhlhorn et al., 2020b). It was suggested that deposition of organic-rich facies and black shales in the Sinemurian appears to be restricted to basins and time-intervals marked by extensional faulting and formations of depocenters with a geometry favourable to water mass stratification and organic matter accumulation and preservation (e.g. the Lusitanian and Wessex basins, Fleet et al., 1987).

A strong cyclicity in TOC distribution in the Wessex Basin is observed in the Lower Pliensbachian (Weedon and Jenkyns, 1990, 1999). It was suggested that these cycles resulted from high-frequency climatic variations that controlled the degree of oxygenation at the seafloor (van Buchem et al., 1995). Changes in the clay mineral assemblages in the Cardigan Bay Basin (Deconinck et al., 2019) reinforce the view that high-frequency climatic variations may have played a pivotal role in the OMPIs associated with the broad Late Sinemurian–Pliensbachian negative carbon isotopic excursion (nCIE). It was recently suggested that the stratigraphic occurrence of positive and negative (0.5–2‰ in TOC) CIEs in Sinemurian and Pliensbachian marine sedimentary archives reflect orbital (astronomical) forcing of the global exogenic carbon cycle, with a ~ 405 kyr eccentricity periodicity (Storm et al., 2020).

3.1.1.1. Sinemurian pCIE. The Sinemurian positive carbon isotopic excursion (pCIE positive and negative trend, ~2.5‰ in TOC) spans from the mid-*semicostatum* to the *obtusum* (*obtusum/stellare* subchronozone) chronozones (Storm et al., 2020) (Fig. 4). Organic-rich facies of this age are geochemically identified by TOC data and temporally constrained by detailed biostratigraphy in the Wessex Basin (Jenkyns and Weedon, 2013; Schöhlhorn et al., 2020b), Lusitanian Basin (Poças Ribeiro et al., 2013; Brito et al., 2017; Duarte et al., 2017), and Cardigan Bay Basin (Storm et al., 2020), and additionally observed, for example, in the Cadwallader Terrane (Porter et al., 2014b) (Appendixes 1–3).

In the Wessex Basin, the organic-rich sedimentary rocks of the local (superregional?) OMPI S3 are associated with relatively more negative $\delta^{13}\text{C}_{\text{org}}$ values (Jenkyns and Weedon, 2013; Schöhlhorn et al., 2020b). A similar situation is observed in the Last Creek 2 section from the Cadwallader Terrane (Porter et al., 2014b), where an interval tentatively dated to the top of the *leslei* (= *turneri*) Chronozone presents the same relationship between TOC and $\delta^{13}\text{C}_{\text{org}}$. However, a recent study from the Wessex Basin seems to indicate that, when accounting for changes in

type and source of OM, a corrected $\delta^{13}\text{C}$ curve ($\delta^{13}\text{C}\text{-HI}$ index = measured $\delta^{13}\text{C}_{\text{TOC}} - (a * \text{HI})$, a being the slope of the correlation trend between $\delta^{13}\text{C}_{\text{TOC}}$ and HI, Schöllhorn et al., 2020b; Suan et al., 2015) shows an association between OMPs and corrected pCIEs. Further studies are necessary to validate that HI variation truly reflects changes in OM or if, for example, this is a reflection of different degradation/diagenetic processes of similar OM (Silva et al., 2013; Suan et al., 2015; Charbonnier et al., 2020; Storm et al., 2020).

3.1.1.2. Sinemurian *Liasidium* Event. The Early Jurassic Sinemurian *Liasidium* Event, *obtusum* chonozone (*denotatus* Subchronozone)–*oxy-notum* chonozone (*oxynotum* Subchronozone) (Riding et al., 2013; Hesselbo et al., 2020) is characterized by the acme of the *Liasidium variable* dinoflagellate and high abundances of *Classopollis classoides*, suggesting an association with high land and sea temperatures (Riding et al., 2013; Hesselbo et al., 2020; Schöllhorn et al., 2020b) (Fig. 4). Masetti et al. (2017) and Franceschi et al. (2019) tentatively correlated the “Arnioceras time Event” nCIE from the southern Alps with the negative shift identified by Riding et al. (2013). So far, this carbon isotopic disturbance appears not to be stratigraphically associated with widespread deposition of OM. However, a poorly age constrained organic-rich interval (currently not defined as an OMPi due to uncertain age assignment) in the Lusitanian Basin (Portugal) may be contemporaneous with the Sinemurian *Liasidium* Event (Appendixes 2 and 3).

3.1.1.3. Sinemurian–Pliensbachian nCIE. The Sinemurian–Pliensbachian nCIE corresponds to a long-term negative and then positive $\delta^{13}\text{C}$ trend of about 2–5‰ in organic matter, expressing a significant change in the global carbon cycle across the Late Sinemurian (*oxynotum* Chronozone) to the Early Pliensbachian (*ibex* Chronozone, *luridum* Subchronozone) (Hesselbo et al., 2000b; Jenkyns et al., 2002; Rosales et al., 2004; Silva et al., 2011; Korte and Hesselbo, 2011; Jenkyns and Weedon, 2013; Duarte et al., 2014; Franceschi et al., 2014; Price et al., 2016; Ruhl et al., 2016; Gómez et al., 2016b; Peti et al., 2017; Danisch et al., 2019; Schöllhorn et al., 2020b; Schöllhorn et al., 2020a; Storm et al., 2020; Mercuzot et al., 2020), including the Sinemurian–Pliensbachian Boundary Event (S-PBE, *rariocostatum* Chronozone, *aplanatum* Subchronozone–*jamesoni* Chronozone, *taylori* Subchronozone, cf. Korte and Hesselbo, 2011). This long-long term nCIE is marked by numerous episodes of local and regional (superregional?) deposition of OM in several European basins, corresponding to OMPs S6, S7, S-P, P1, P2, P3, and P4 (Table 1, Fig. 3 and Fig. 4, and Appendixes 2 and 3).

Earth system processes leading to the long ranging Sinemurian–Pliensbachian nCIE and climatic and environmental changes are largely unknown but are speculated to relate to either the opening of the Hispanic Corridor or a late phase of enhanced global continental (silicate) weathering induced by the Central Atlantic Margin Province volcanism (see discussion in Korte and Hesselbo, 2011; Porter et al., 2013; Gómez et al., 2016b; Price et al., 2016; Ruhl et al., 2016; Danisch et al., 2019; Franceschi et al., 2019; Schöllhorn et al., 2020b, Schöllhorn et al., 2020a; Storm et al., 2020). Considering the extremely limited stratigraphic data outside the northern western Tethys realm (see A1.4), significant accumulation of OM (high TOC) during the Sinemurian–Pliensbachian nCIE appears to be most expressive in the Wessex and Lusitanian basins; the exception is the regional (superregional?) OMPi S-P where sedimentary OM enrichment was more widespread (Fig. 3). The regional (superregional?) OMPi S-P is associated with the most negative $\delta^{13}\text{C}$ values of the broad Sinemurian–Pliensbachian nCIE [including the Sinemurian–Pliensbachian boundary event (S-PBE) cf. Korte and Hesselbo, 2011], and coincides with the final stages of the Late Sinemurian warming event and transition to subsequent cooling (Korte and Hesselbo, 2011; Gómez et al., 2016b). The Trento platform (Italy) is marked by platform sedimentary deposits with relatively elevated TOC, but below 2%, which are interpreted to be associated with the S-PBE, suggesting relative OM enrichment even in shallow areas at this time

(Franceschi et al., 2014, 2019).

3.1.2. Late Pliensbachian pCIE: OMPs P7–P10

The Late Pliensbachian is marked by a relatively low amplitude long-term pCIE of about 2–4‰ in carbonate, peaking in the *margaritatus* Chronozone (circa *gibbosus* Subchronozone) (Fig. 4 and Appendix 3). The regional (superregional?) OMPs P7, P8, and P9 are concomitant with superimposed smaller-scale $\delta^{13}\text{C}$ positive excursions (~0.5–2‰ in carbonate and TOC) observed in multiple basins (see Silva et al., 2011), possibly expressing long (~405 kyr) eccentricity modulation of the global carbon cycle and climatic and environmental conditions at those times (Jenkyns et al., 2002; Silva et al., 2011, 2015; Caruthers et al., 2014; Silva and Duarte, 2015; Mercuzot et al., 2020; Storm et al., 2020).

Despite the recognized gaps in data coverage, organic-rich facies associated with the Late Pliensbachian pCIE are (apparently) concentrated mainly along the areas bordering the proto-Atlantic Ocean and possibly the Boreal Sea (Fig. 3). During the Late Pliensbachian, organic productivity and preservation were enhanced during major transgressive episodes (Hallam, 1981; de Oliveira et al., 2006; Duarte et al., 2010; Silva et al., 2011, 2012, 2015; Silva and Duarte, 2015). Rosales et al. (2006) suggested a link between second-order relative sea-level changes in the Basque–Cantabrian Basin and variations in seawater geochemistry during the Early Jurassic, noting a coincidence between transgressions and increasing $\delta^{13}\text{C}$, TOC and Hydrogen Index (and vice-versa for regressions). It was also suggested that the Late Pliensbachian black shales were associated with warm temperatures (hyperthermals?) within the relatively cooler Late Pliensbachian (Silva and Duarte, 2015; Peti and Thibault, 2017; Schöllhorn et al., 2020b).

3.1.3. *Spinatum* nCIE: OMPi P11

The *spinatum* Chronozone is generally regarded as a cool interval (Hinnov and Park, 1999; Price, 1999; Dera et al., 2009b; Dera et al., 2009a; Suan et al., 2010; Korte and Hesselbo, 2011; Silva and Duarte, 2015; Gómez et al., 2016b; Ruebsam et al., 2019; Deconinck et al., 2020; Storm et al., 2020). It has been suggested that geological storage of OM associated with the Late Pliensbachian pCIE resulted in decreased atmospheric CO_2 , triggering and/or amplifying cooling in the *spinatum* Chron (e.g. Suan et al., 2010; Silva et al., 2011; Silva and Duarte, 2015; Storm et al., 2020). It was also speculated that a potentially early onset of North Sea doming may have caused the development of regressive facies in the Late Pliensbachian of the North Sea region, which may have promoted the regional shift from a warm to a cooler climate mode (Korte et al., 2015) (Fig. 2). More recently, De Lena et al. (2019) suggested the degassing of volcanic S-species (SO_2) via volcanism was an unlikely driving mechanism of the cool and dry climate of the late Pliensbachian.

The later stages of the *spinatum* nCIE (cf. Storm et al., 2020) were associated with organic-rich rocks defining the superregional OMPi P11 (Figs. 3 and 4).

3.2. Latest Pliensbachian–Toarcian

The latest Pliensbachian–Toarcian is characterized by large amplitude variations in the $\delta^{13}\text{C}$ record. The Early Toarcian is marked by one of the most significant carbon cycle perturbations of the past 200 million years, expressed by a large amplitude nCIE, up to 7‰ in TOC (e.g. Hesselbo et al., 2000a; Röhl et al., 2001; Kemp et al., 2005; Suan et al., 2015), up to 8‰ in fossil wood (e.g. Hesselbo et al., 2007), up to 5‰ in bulk carbonate (e.g. Hermoso et al., 2009; Kafousia et al., 2011), and up to 4‰ in macrofossil carbonate (e.g. brachiopods (Suan et al., 2010; Ullmann et al., 2020) superimposed on a broader Early Toarcian pCIE, which encompasses the *tenuicostatum*–*bifrons* time interval (Fig. 4) (Küspert, 1982; Jenkyns and Clayton, 1986; Jenkyns, 1988, 2010; Hesselbo et al., 2000a, 2007; Röhl et al., 2001; Jenkyns et al., 2002; Cohen et al., 2004; Röhl and Schmid-Röhl, 2005; Woodfine et al., 2008; Hermoso et al., 2009, 2012, 2013; Al-Suwaidi et al., 2010, 2016; Bodin et al., 2010; Littler et al., 2010; Bodin et al., 2016; Suan et al., 2010,

2011, 2015; Lézin et al., 2013; Sabatino et al., 2013; Kafousia et al., 2014; Kemp and Izumi, 2014; Pittet et al., 2014; Reolid et al., 2014; Caruthers et al., 2014; Pieńkowski et al., 2016; Martinez et al., 2017; Silva et al., 2017; Them et al., 2017; Xu et al., 2017b, 2018; Fantasia et al., 2018, 2019b; Ruebsam et al., 2019, 2020; Boulila et al., 2019; Danise et al., 2019; Jin et al., 2020; Ruebsam and Al-Husseini, 2020).

As previously observed, Lower Toarcian organic-rich facies are described in many areas of the northern West Tethys Shelf, Boreal and Panthalassic margins (e.g. Baudin et al., 1990; Röhl et al., 2001; van de Schootbrugge et al., 2005b; Gómez et al., 2008; McArthur et al., 2008; Al-Suwaidi et al., 2010; Jenkyns, 2010; Suan et al., 2011, 2015; Hermoso et al., 2012; Kemp and Izumi, 2014; Pieńkowski et al., 2016; Silva et al., 2017; Them et al., 2017; Ruvalcaba Baroni et al., 2018; Rodrigues et al., 2020c). These are particularly expressive in the central and northern European epicontinental areas, where deposition took place in anoxic basins under dominantly warm and humid conditions and associated with an intensification of the hydrological cycle and fluvial runoff (Jenkyns, 1988, 2010; Rees et al., 1999; Cohen et al., 2004; Pearce et al., 2008; Dera et al., 2009a, 2009b; Brazier et al., 2015; Them et al., 2017; Fantasia et al., 2018; Rodrigues et al., 2021; Rodrigues et al., 2020b). In the southern areas of the West Tethys Shelf, such as western and southern Iberian, and northern Gondwana margins, organic-rich intervals are spatially and temporally restricted (e.g. Hesselbo et al., 2007; Bodin et al., 2010; Gómez and Goy, 2011; Reolid et al., 2012, 2014; Rodrigues et al., 2020b, 2021; Rodrigues et al., 2016, 2019, 2020a; Ait-Itto et al., 2017; Fantasia et al., 2019a; Ruebsam et al., 2020; Silva et al., 2021) and the absence of significant Lower Toarcian OM enrichment is interpreted as a consequence of oxic conditions and low productivity associated with a semi-arid climate and general circulation patterns (e.g. van de Schootbrugge et al., 2005b; Reolid et al., 2014, 2020; Ruvalcaba Baroni et al., 2018; Fantasia et al., 2019b; Rodrigues et al., 2019, 2020a, 2020b; Ruebsam et al., 2020) (Fig. 3 and Appendix 2).

3.2.1. P-Toa nCIE: OMPI P-Toa

A smaller-scale (~2–3‰ in TOC) nCIE is observed at the base of the Toarcian *polymorphum* Chronozone, the Pliensbachian–Toarcian event (P–Toa Event), coincidental with the superregional OMPI P–Toa (Hesselbo et al., 2007; Littler et al., 2010). It was suggested that the P–Toa event resulted from a carbon cycle perturbation analogous to the one marking the T-OAE (Littler et al., 2010), and it is stratigraphically associated with an extinction phase in marine benthos at the base of the Toarcian (e.g. Little and Benton, 1995; Littler et al., 2010; Caruthers et al., 2013) and significant continental weathering and volcanic activity (Percival et al., 2015, 2016).

3.2.2. The broad Lower Toarcian pCIE and the T-OAE nCIE: OMPIs T1, T2 (OAE)–T4(OAE), and T 5–7

The superregional OMPI T1 spans the *paltum*–*tenuicostatum* subchrons and is associated with the onset of a broad Lower Toarcian pCIE onto which is superimposed the T-OAE nCIE (Fig. 4). The T-OAE is characterized by the geographically widespread deposition of organic-rich sediments (Fig. 4), ocean anoxia - when geographically widespread areas of the seafloor and oceans become oxygen depleted or anoxic (Schlanger and Jenkyns, 1976; Küspert, 1982; Jenkyns and Clayton, 1986; Jenkyns, 1988; Dickson et al., 2017), global warming (Bailey et al., 2003; Dera et al., 2009b; Suan et al., 2010; Danise et al., 2013; Ullmann et al., 2020), ocean acidification (e.g. Suan et al., 2010; Müller et al., 2020), a carbonate productivity crisis (e.g. Mattioli et al., 2009; Han et al., 2018), and a 2nd-order mass extinction event in benthic and pelagic groups (e.g. Aberhan and Baumiller, 2003; Arias, 2006; Mattioli et al., 2009; Caswell et al., 2009; Comas-Rengifo et al., 2010; Gómez and Arias, 2010; Gómez and Goy, 2011; García Joral et al., 2011; Fraguas et al., 2012; Danise et al., 2013; Caswell and Frid, 2017; Piazza et al., 2019; Cabral et al., 2020). Several mechanisms have been suggested to have initiated the T-OAE nCIE and carbon cycle perturbation: (1) carbon degassing from the Karoo-Ferrar Large Igneous Province

(Pálffy and Smith, 2000, 2) rapid release of methane from gas hydrates from continental-margin and slope sediments (Hesselbo et al., 2000a), (3) thermogenic methane release from subsurface organic-rich shales intruded by dyke and sills associated with Karoo-Ferrar emplacement (McElwain et al., 2005; Svensen et al., 2007, 4) increased CO₂ release from the decomposition and oxidation of terrestrial organic matter associated with increased fungal activity (Pieńkowski et al., 2016), 5) elevated methane release from terrestrial environments (Them et al., 2017), and (6) destabilization of labile cryospheric carbon reservoirs (Silva and Duarte, 2015; Krencker et al., 2019; Ruebsam et al., 2019).

The superregional OMPIs T2(OAE), T3(OAE), and T4(OAE) are associated with widespread deposition of organic-rich facies with high TOC, including black shales. Maximum TOC contents are frequently observed in the superregional OMPI T4(OAE), associated with the positive $\delta^{13}\text{C}$ trend of the T-OAE nCIE (the positive carbon isotopic trend of the T-OAE nCIE is here defined as the interval between minimum $\delta^{13}\text{C}$ and maximum $\delta^{13}\text{C}$ that marks the end of the T-OAE interval) (Fig. 4). The superregional OMPI T4(OAE) is observed even in areas not characterized by significant Lower Toarcian organic-rich deposition, such as the Lusitanian Basin (de Oliveira et al., 2006; Hesselbo et al., 2007; Rodrigues et al., 2016, 2020a).

Carbon sequestration during the T-OAE seems to be broadly associated with a peak in $^{187}\text{Os}/^{188}\text{Os}$ (Cohen et al., 2004; Percival et al., 2016; Them et al., 2017; van Aken et al., 2019) and a minimum in $\delta^{18}\text{O}$ (Suan et al., 2008; Dera et al., 2011; Krencker et al., 2014; Ullmann et al., 2014, 2020; Korte et al., 2015), suggesting that maximum organic carbon sequestration during the T-OAE was concomitant with sustained high temperatures and enhancement of the hydrological cycling and increased weathering rates (Jenkyns, 2010; Xu et al., 2017b; Fantasia et al., 2019a; Rodrigues et al., 2020b; Ullmann et al., 2020). The here demonstrated offset between the T-OAE $\delta^{13}\text{C}$ negative trend and the maximum sedimentary TOC values combined with the widespread nature of the superregional OMPI T4(OAE) agree with recent modelling efforts indicating a lag between peak greenhouse gas input and peak anoxia/carbon sequestration (Ullmann et al., 2020).

Several OMPIs are observed after the T-OAE (Fig. 4 and Appendixes 2 and 3). The superregional OMPIs T5 and T6 and the regional (superregional?) OMPI T7 are associated with the decreasing limb of the broad Early Toarcian $\delta^{13}\text{C}$ positive excursion initiated at the base of the *tenuicostatum* Chronozone (Fig. 4). Oxygen stressed conditions and carbon sequestration were prevalent from the mid-*semicelatum* Subchron (*tenuicostatum* Chron) to the upper *bifrons* Chron in several locations, as recorded in the German sections (Röhl et al., 2001), Cleveland Basin (Thibault et al., 2018), and Slyne Basin (Silva et al., 2017). Post-OAE black shales in the Paris Basin were likely concomitant with third-order sea-level changes (Hermoso et al., 2013; sensu de Graciansky et al., 1998).

4. Comparison with the Cretaceous OAEs

OAEs were initially defined as brief time intervals during which significant portions of the global oceans were extensively deoxygenated (Schlanger and Jenkyns, 1976; Jenkyns, 1980, 2010; Arthur et al., 1990; Leckie et al., 2002). Although a comprehensive model to explain all OAEs is not yet fully constrained, the consensus is that these events were associated with a sudden influx of greenhouse gases into the atmosphere and global warming, leading to an acceleration of the hydrological cycle, increased continental weathering, enhanced nutrient delivery to the oceans, intensified oceanic upwelling, and increased productivity in marine and continental environments (Schlanger and Jenkyns, 1976; Weissert, 1989; Arthur et al., 1990; Hesselbo et al., 2000a; Leckie et al., 2002; Bodin et al., 2010; Jenkyns, 2010; Trabucho-Alexandre et al., 2011; Herrle et al., 2015; Montero-Serrano et al., 2015; Xu et al., 2017b; Fantasia et al., 2018; Silva et al., 2020b).

OAEs are characterized by negative or positive CIEs, interpreted to reflect the sudden influx of isotopically light CO₂ into the atmosphere

(either from, for example, volcanism, metamorphic degassing, or destabilization of gas hydrates) or burial of vast amounts of organic carbon, respectively (Jenkyns, 2010). However, local basinal responses often override global forcing on the occurrence, onset, and termination of organic-rich deposition associated with OAEs (e.g. Trabucho-Alexandre et al., 2011; Fantasia et al., 2019a; Rodrigues et al., 2019, 2020a) and CIEs (reflecting global carbon cycle change) in combination with evidence for regional to superregional bottom-water deoxygenation have become the accepted characteristic features of an OAE, rather than widespread organic-rich deposition alone (Jenkyns, 2010).

The main Mesozoic OAEs comprise the T-OAE, Early Aptian OAE (OAE 1a), and Cenomanian–Turonian OAE (OAE 2). Secondary OAE events, characterized by less prominently expressed CIE and less severe paleoenvironmental perturbations, are the Valanginian OAE (Weissert Event), Hauterivian OAE (Faraoni Event), Late Aptian–Early Albian OAE (OAE 1b cluster), and (more doubtfully) the Late Albian (OAE 1c and OAE 1d) and the Coniacian–Santonian (OAE 3) (Jenkyns, 2010). Main and secondary OAEs are characterized by relatively brief episodes of regional to superregional deposition of organic matter associated with

disruption of the global carbon cycle (Jenkyns, 1980, 2010; Arthur et al., 1990; Leckie et al., 2002) (Fig. 5).

The compilation of Sinemurian–Toarcian OMPIs presented in this paper demonstrates that regional and superregional sedimentary sequestration of OM during the Early Jurassic was more common than previously thought (Fig. 5). Because of the association of OMPIs with well-constrained CIEs (Figs. 3 and 4 and Appendices 3 and 4), we propose that the regional (superregional?) OMPI S-P and the superregional OMPI P11 and OMPI P-Toa should now be considered as secondary OAEs, similarly to the secondary Cretaceous OAEs.

The Early Toarcian superregional OMPIs T1, T5, T6, and the regional (superregional?) OMPI T7 occur in association with the Lower Toarcian pCIE, and their spatial spread supports some degree of equivalency to the secondary Cretaceous OAEs. These OMPIs demonstrate that the establishment of (global?) environmental and depositional conditions leading to widespread preservation of OM in the Late Pliensbachian–Early Toarcian was not exclusively associated with the T-OAE.




Fig. 5. Mesozoic OAEs and comparison with the Early Jurassic OMPIs record (the grey bars correspond to the main Jurassic–Cretaceous OAEs). The Sinemurian–Pliensbachian record was created using the time scale and $\delta^{13}\text{C}$ from the reference Mochras core, Cardigan Bay Basin, UK (Storm et al., 2020). The Toarcian and Cretaceous time scale (Ogg et al., 2016) and $\delta^{13}\text{C}$ record (Herrle et al., 2004; Reichelt, 2005; Föllmi et al., 2006; Jarvis et al., 2006; Hesselbo et al., 2007; Voigt et al., 2012; Martinez et al., 2015; Xu et al., 2018) were generated using TSCreator software (PUBLIC7.4_windows10_latest_06September2019). Cretaceous OAEs are from Arthur et al. (1990); Leckie et al. (2002); Bodin et al. (2007); Mutterlose et al. (2009); Jenkyns (2010); Westermann et al. (2010); Föllmi et al. (2012); Herrle et al. (2015).

5. Limitations and future challenges

The methodology presented here for resolving OMPs applies to any geological interval, is simple and straightforward, and provides a framework for exploratory research on large scale sedimentary organic matter sequestration and its impacts on the global carbon cycle. However, mechanistic understanding of sedimentary OM sequestration at basinal, intrabasinal, and global scales is often hampered by the lack of spatial data coverage, temporal data uncertainty, and sampling/data resolution for many of the intervals studied (see A1.4). In addition, understanding of sedimentation rates and their impact on true rates of organic carbon accumulation, or carbon burial fluxes, is often limited or non-existent; this is particularly important when using a fixed cut-off value of 2% present-day TOC to define an OMP or possible source-rocks. For example, sedimentary successions with low TOC/high sedimentation rate may represent the same organic matter accumulation per time unit as high TOC/low sedimentation rate depositional environments. Furthermore, the intrabasinal spatial extent of an OMP is often unconstrained, even when using the most advanced imaging/modelling tools (e.g. Bruneau et al., 2017). A detailed assessment of sedimentation rates and the translation of present-day TOC values into organic matter accumulation/sequestration rates (or fluxes) for individual OMPs at the outcrop and basin-scale are outside the scope of this study.

Thermal maturation may also lead to an underestimation of the original OM accumulation rate, and possibly even the spatial extent of OMPs, especially if original TOC approximated 2% in now highly thermally mature successions. The magnitude of this effect depends mainly on the level of maturity and the relative proportions of generative and non-generative organic carbon compounds, with type I and type II kerogens having a higher proportion of generative organic carbon which therefore are prone to have depressed original TOC values (Jarvie, 2012). Type III and type IV kerogens are less affected by thermal maturation, and its suppressing effect on TOC, as hydrocarbons generation is limited. Therefore, the loss of carbon via maturation and expulsion may thus have resulted in an underestimation of organic carbon burial throughout geological history, and during Early Jurassic OMPs, when using uncorrected present-day TOC values as a proxy.

Although the present work presents a framework for the stratigraphic distribution of OMPs across the ~20 Myr of the Sinemurian–Toarcian time interval, we emphasize that without a detailed assessment of (1) inter- and intrabasinal bio- and chronostratigraphic correlations, (2) duration and stratigraphic completeness, (3) TOC stratigraphic and basinal variability, and (4) thermal maturation and diagenesis, the study of OMPs as defined here will generally underestimate the spatial and temporal patterns in geological organic carbon storage and, therefore, result in inaccurate estimates on OM burial and as a sink of atmospheric CO₂. Future research should aim to disentangle (1) the noted complexities in estimating original TOC and organic carbon accumulation rates, (2) temporal and spatial variability in environmental or Earth system feedback mechanisms driving sedimentary carbon sequestration, and (3) their combined impact on the global carbon cycle. Further work is expected to define additional Early Jurassic OMPs and to improve current understanding of the spatial and temporal distribution of individual OMPs.

6. Conclusions

The recognition of several Early Jurassic organic matter preservation intervals (OMPs) of regional and superregional extent gives new insight into the role that OM sequestration played in Earth System processes and provides support to current theoretical models for carbon cycle perturbations in the Early Jurassic (e.g., the T-OAE). From this study, we conclude that:

1. Sinemurian regional OMPs were mainly restricted to a small number of European basins. A relatively more widespread OMP was

associated with the most negative $\delta^{13}\text{C}$ interval of the long-term Sinemurian–Pliensbachian $\delta^{13}\text{C}$ excursion. The stratigraphic and causal relationship between OMPs and CIEs (as representative of global carbon cycle perturbations) is not always straightforward.

2. Pliensbachian regional (superregional?) OMPs occur mostly in the basins bordering the proto-Atlantic Ocean, often associated with positive $\delta^{13}\text{C}$ excursions and short-lived warming intervals (hyperthermals?). As suggested previously, it is apparent that significant OM sequestration predates [regional (superregional?) OMPs P7, P8, and P9] and coincides with the onset of the Late Pliensbachian cool episode and the *spinatum* nCIE (superregional OMP P11).
3. As extensively discussed in the existing literature, Early Toarcian OMPs are widespread. The Early Toarcian superregional OMPs T2 (OAE), T3(OAE) and T4(OAE) are associated with the most pronounced $\delta^{13}\text{C}$ negative carbon isotopic excursion in the Mesozoic. Maximum TOC contents [superregional OMP T4(OAE)] occur in association with the positive $\delta^{13}\text{C}$ trend of the Early Toarcian Oceanic Anoxic Event (T-OAE), indicating, and as suggested by recent modelling efforts, that peak OM sequestration and anoxia post-dated the main event of carbon injection into the atmosphere. However, several superregional OMPs pre- and postdated the T-OAE interval. These OMPs demonstrate that the establishment of (global?) environmental and depositional changes led to widespread preservation of OM in the Late Pliensbachian and Early Toarcian and thus not exclusively during the main Early Toarcian OAE.
4. This study demonstrates that organic-rich intervals of regional and superregional expression in the Early Jurassic are more common than previously thought. It is contended that considering the association of the regional (superregional?) OMP S-P (*raricostatum*, *raricostatooides-jamesoni*, *polymorphus*) and the superregional OMP P11 (*spinatum*, *hawskerense*) and OMP P-Toa with well-characterized CIEs, these should now be considered secondary OAEs, similarly to the secondary Cretaceous OAEs. The superregional Early Toarcian OMPs T1 (*tenuicostatum*, *paltum-tenuicostatum*), T5 (*falciferum*, *elegans-falciferum*), T6 (*bifrons*, *commune*), and the regional (superregional?) OMP T7 (*bifrons*, *fibulatum*) occur in association with the broad lower Toarcian pCIE.

Declaration of Competing Interest

None.

Acknowledgements

This research was supported by the Source Rock and Geochemistry of the Central Atlantic Margins consortium (Dalhousie University, Basin and Reservoir Lab, PI-Grant Wach). Ricardo L. Silva was also supported by iCRAG (project: Temporal and spatial variability in Lower Jurassic hydrocarbon source rock quality in Irish off-shore marine basins, PI-Dr Micha Ruhl). Luis V. Duarte was partially supported by the strategic project UID/MAR/04292/2019 granted to the Marine and Environmental Sciences Centre (MARE). Stephen Hesselbo acknowledges funding from NERC (NE/N018508/1). We gratefully acknowledge the Editor Shuhab Khan, and Stephane Bodin and three other anonymous referees for their insightful comments that greatly benefited and improved the manuscript. This study is a contribution to the UNESCO-IUGS projects IGCP 655 Toarcian Oceanic Anoxic Event: Impact on marine carbon cycle and ecosystems and IGCP 739 The Mesozoic–Paleogene hyperthermal events.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.earscirev.2021.103780>.

References

- Aberhan, M., Baumiller, T.K., 2003. Selective extinction among Early Jurassic bivalves: a consequence of anoxia. *Geology* 31, 1077–1080. <https://doi.org/10.1130/G19938.1>.
- Adam, P., Philippe, E., Albrecht, P., 1998. Photochemical sulfurization of sedimentary organic matter: a widespread process occurring at early diagenesis in natural environments? *Geochim. Cosmochim. Acta* 62, 265–271. [https://doi.org/10.1016/S0016-7037\(97\)00332-3](https://doi.org/10.1016/S0016-7037(97)00332-3).
- Ait-Itto, F.-Z., Price, G.D., Ait Addi, A., Chafiki, D., Mannani, I., 2017. Bulk-carbonate and belemnite carbon-isotope records across the Pliensbachian-Toarcian boundary on the northern margin of Gondwana (Issouka, Middle Atlas, Morocco). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 466, 128–136. <https://doi.org/10.1016/j.palaeo.2016.11.014>.
- Al-Suwaidi, A., Angelozzi, G.N., Baudin, F., Damborenea, S.E., Hesselbo, S.P., Jenkyns, H.C., Mancenido, M.O., Riccardi, A.C., 2010. First record of the Early Toarcian Oceanic Anoxic Event from the Southern Hemisphere, Neuquen Basin, Argentina. *J. Geol. Soc. London* 167, 633–636. <https://doi.org/10.1144/0016-76492010-025.SPECIAL>.
- Al-Suwaidi, A.H., Hesselbo, S.P., Damborenea, S.E., Mancenido, M.O., Jenkyns, H.C., Riccardi, A.C., Angelozzi, G.N., Baudin, F., 2016. The Toarcian Oceanic Anoxic Event (Early Jurassic) in the Neuquén Basin, Argentina: a reassessment of age and carbon isotope stratigraphy. *J. Geol.* 124, 171–193. <https://doi.org/10.1086/684831>.
- Alves, T.M., Gawthorpe, R.L., Hunt, D.W., Monteiro, J.H., 2002. Jurassic tectono-sedimentary evolution of the Northern Lusitanian Basin (offshore Portugal). *Mar. Pet. Geol.* 19, 727–754. [https://doi.org/10.1016/S0264-8172\(02\)00036-3](https://doi.org/10.1016/S0264-8172(02)00036-3).
- Arias, C., 2006. Northern and Southern Hemispheres ostracod palaeobiogeography during the Early Jurassic: possible migration routes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 233, 63–95. <https://doi.org/10.1016/j.palaeo.2005.12.013>.
- Arthur, M.A., Brumsack, H.-J., Jenkyns, H.C., Schlanger, S.O., 1990. Stratigraphy, geochemistry, and paleoceanography of organic carbon-rich Cretaceous sequences. In: Ginsburg, R.N., Beaudoin, B. (Eds.), *Cretaceous Resources, Events and Rhythms*. Springer Netherlands, Dordrecht, pp. 75–119. https://doi.org/10.1007/978-94-015-6861-6_6.
- Asgar-Deen, M., Hall, R., Craig, J., Riediger, C., 2003. New biostratigraphic data from the Lower Jurassic Fernie Formation in the subsurface of west-central Alberta and their stratigraphic implications. *Can. J. Earth Sci.* 40, 45–63. <https://doi.org/10.1139/e02-096>.
- Aurell, M., Robles, S., Bádenas, B., Rosales, I., Quesada, S., Meléndez, G., García-Ramos, J.C., 2003. Transgressive-regressive cycles and Jurassic palaeogeography of northeast Iberia. *Sediment. Geol.* 162, 239–271. [https://doi.org/10.1016/S0037-0738\(03\)00154-4](https://doi.org/10.1016/S0037-0738(03)00154-4).
- Bailey, T.R., Rosenthal, Y., McArthur, J.M., van de Schootbrugge, B., Thirlwall, M.F., 2003. Paleocceanographic changes of the Late Pliensbachian-Early Toarcian interval: a possible link to the genesis of an Oceanic Anoxic Event. *Earth Planet. Sci. Lett.* 212, 307–320. [https://doi.org/10.1016/S0012-821X\(03\)00278-4](https://doi.org/10.1016/S0012-821X(03)00278-4).
- Baudin, F., 1995. Depositional controls on mesozoic source rocks in the tethys. In: Huc, A.-Y. (Ed.), *Paleogeography, Paleoclimate, and Source Rock*. The American Association of Petroleum Geologists, Houston, pp. 191–211.
- Baudin, F., Herbin, J.P., Vandenbroucke, M., 1990. Mapping and geochemical characterization of the Toarcian organic matter in the Mediterranean Tethys and Middle East. *Org. Geochem.* 16, 677–687. [https://doi.org/10.1016/0146-6380\(90\)90109-D](https://doi.org/10.1016/0146-6380(90)90109-D).
- Berner, R.A., 1999. A new look at the long-term carbon cycle. *GSA Today* 9, 2–6. <https://doi.org/10.1002/qj.49710443903>.
- Berner, R.A., 2003. The long-term carbon cycle, fossil fuels and atmospheric composition. *Nature*. <https://doi.org/10.1038/nature02131>.
- Berner, R.A., 2006. GEOCARBSULF: a combined model for Phanerozoic atmospheric O₂ and CO₂. *Geochim. Cosmochim. Acta* 70, 5653–5664. <https://doi.org/10.1016/j.gca.2005.11.032>.
- Bessereau, G., Guillocheau, F., Huc, A.-Y., 1995. Source rock occurrence in a sequence stratigraphic framework: the example of the Lias of the Paris Basin. In: Huc, A.-Y. (Ed.), *Paleogeography, Paleoclimate, and Source Rock*. The American Association of Petroleum Geologists, Houston, pp. 273–301.
- Bishop, A.N., Kearsley, A.T., Patience, R.L., 1992. Analysis of sedimentary organic materials by scanning electron microscopy: the application of backscattered electron imagery and light element X-ray microanalysis. *Org. Geochem.* 18, 431–446. [https://doi.org/10.1016/0146-6380\(92\)90106-8](https://doi.org/10.1016/0146-6380(92)90106-8).
- Bodin, S., Godet, A., Matera, V., Steinmann, P., Vermeulen, J., Gardin, S., Adatte, T., Coccioni, R., Föllmi, K.B., 2007. Enrichment of redox-sensitive trace metals (U, V, Mo, As) associated with the late Hauterivian Faraoni oceanic anoxic event. *Int. J. Earth Sci.* 96, 327–341. <https://doi.org/10.1007/s00531-006-0091-9>.
- Bodin, S., Mattioli, E., Fröhlich, S., Marshall, J.D., Boutib, L., Lahsini, S., Redfern, J., 2010. Toarcian carbon isotope shifts and nutrient changes from the Northern margin of Gondwana (High Atlas, Morocco, Jurassic): Palaeoenvironmental implications. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 297, 377–390. <https://doi.org/10.1016/j.palaeo.2010.08.018>.
- Bodin, S., Fröhlich, S., Boutib, L., Lahsini, S., Redfern, J., 2011. Early Toarcian source-rock potential in the central High Atlas Basin (central Morocco): regional distribution and depositional model. *J. Pet. Geol.* 34, 345–363. <https://doi.org/10.1111/j.1747-5457.2011.00509.x>.
- Bodin, S., Krencker, F.-N., Kothe, T., Hoffmann, R., Mattioli, E., Heimhofer, U., Kabiri, L., 2016. Perturbation of the carbon cycle during the late Pliensbachian – early Toarcian: new insight from high-resolution carbon isotope records in Morocco. *J. African Earth Sci.* 116, 89–104. <https://doi.org/10.1016/j.jafrearsci.2015.12.018>.
- Borrego, A.G., Hagemann, H.W., Blanco, C.G., Valenzuela, M., Suárez De Centi, C., 1996. The Pliensbachian (Early Jurassic) “anoxic” event in Asturias, northern Spain: Santa Mera Member, Rodiles formation. *Org. Geochem.* 25, 295–309. [https://doi.org/10.1016/S0146-6380\(96\)00121-0](https://doi.org/10.1016/S0146-6380(96)00121-0).
- Boullila, S., Galbrun, B., Sadki, D., Gardin, S., Bartolini, A., 2019. Constraints on the duration of the early Toarcian T-OAE and evidence for carbon-reservoir change from the High Atlas (Morocco). *Glob. Planet. Change* 175, 113–128. <https://doi.org/10.1016/j.gloplacha.2019.02.005>.
- Bowden, S.A., Farrimond, P., Snape, C.E., Love, G.D., 2006. Compositional differences in biomarker constituents of the hydrocarbon, resin, asphaltene and kerogen fractions: an example from the Jet Rock (Yorkshire, UK). *Org. Geochem.* 37, 369–383. <https://doi.org/10.1016/j.orggeochem.2005.08.024>.
- Brazier, J.M., Suan, G., Tacail, T.T., Simon, L., Martin, J.E., Mattioli, E., Balter, V., 2015. Calcium isotope evidence for dramatic increase of continental weathering during the Toarcian oceanic anoxic event (Early Jurassic). *Earth Planet. Sci. Lett.* 411, 164–176. <https://doi.org/10.1016/j.epsl.2014.11.028>.
- Brito, M., Rodrigues, R., Baptista, R., Duarte, L.V., Azerêdo, A.C., Jones, C.M., 2017. Geochemical characterization of oils and their correlation with Jurassic source rocks from the Lusitanian Basin (Portugal). *Mar. Pet. Geol.* 85, 26. <https://doi.org/10.1016/j.marpetgeo.2017.05.010>.
- Bruneau, B., Chauveau, B., Baudin, F., Moretti, I., 2017. 3D stratigraphic forward numerical modelling approach for prediction of organic-rich deposits and their heterogeneities. *Mar. Pet. Geol.* 82, 1–20. <https://doi.org/10.1016/j.marpetgeo.2017.01.018>.
- Bucefalo Palliani, R., Cirilli, S., Mattioli, E., 1998. Phytoplankton response and geochemical evidence of the lower Toarcian relative sea level rise in the Umbria-Marche basin (Central Italy). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 142, 33–50. [https://doi.org/10.1016/S0031-0182\(97\)00152-1](https://doi.org/10.1016/S0031-0182(97)00152-1).
- Burdige, D.J., 2007. Preservation of organic matter in marine sediments: controls, mechanisms, and an imbalance in sediment organic carbon budgets? *Chem. Rev.* 107, 467–485. <https://doi.org/10.1021/cr050347q>.
- Cabral, M.C., Lord, A.R., Pinto, S., Duarte, L.V., Azerêdo, A.C., 2020. Ostracods of the Toarcian (Jurassic) of Peniche, Portugal: taxonomy and evolution across and beyond the GSSP interval. *Bull. Geosci.* 243–278. <https://doi.org/10.3140/bull.geosci.1778>.
- Campana, F., Fantoni, R., Masetti, D., Scotti, P., 2017. Stratigraphic and naphthogenic characterization of the Sparviero 1 bis Mesozoic succession (southern Adriatic basin, Italy). *J. Mediterr. Earth Sci.* 9, 89–93.
- Caruthers, A.H., Gröcke, D.R., Smith, P.L., 2011. The significance of an Early Jurassic (Toarcian) carbon-isotope excursion in Haida Gwaii (Queen Charlotte Islands), British Columbia, Canada. *Earth Planet. Sci. Lett.* 307, 19–26. <https://doi.org/10.1016/j.epsl.2011.04.013>.
- Caruthers, A.H., Smith, P.L., Gröcke, D.R., 2013. The Pliensbachian-Toarcian (Early Jurassic) extinction, a global multi-phased event. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 386, 104–118. <https://doi.org/10.1016/j.palaeo.2013.05.010>.
- Caruthers, A.H., Smith, P.L., Grocke, D.R., 2014. The Pliensbachian-Toarcian (Early Jurassic) extinction: a North American perspective. In: Keller, G., Kerr, A.C. (Eds.), *Volcanism, Impacts, and Mass Extinctions: Causes and Effects*. Geological Society of America, pp. 225–243. [https://doi.org/10.1130/2014.2505\(11\)](https://doi.org/10.1130/2014.2505(11)).
- Caswell, B.A., Coe, A.L., 2012. A high-resolution shallow marine record of the Toarcian (Early Jurassic) Oceanic Anoxic Event from the East Midlands Shelf, UK. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 365–366, 124–135. <https://doi.org/10.1016/j.palaeo.2012.09.021>.
- Caswell, B.A., Frid, C.L.J., 2017. Marine ecosystem resilience during extreme deoxygenation: the Early Jurassic oceanic anoxic event. *Oecologia* 183, 275–290. <https://doi.org/10.1007/s00442-016-3747-6>.
- Caswell, B.A., Coe, A.L., Cohen, A.S., 2009. New range data for marine invertebrate species across the early Toarcian (Early Jurassic) mass extinction. *J. Geol. Soc. Lond.* 166, 859–872. <https://doi.org/10.1144/0016-76492008-0831>.
- Chadwick, R.A., 1986. Extension tectonics in the Wessex Basin, southern England. *J. Geol. Soc. London* 143, 465–488. <https://doi.org/10.1144/gsjgs.143.3.0465>.
- Charbonnier, G., Adatte, T., Föllmi, K.B., Suan, G., 2020. Effect of intense weathering and postdepositional degradation of organic matter on Hg/TOC proxy in organic-rich sediments and its implications for deep-time investigations. *Geochem. Geophys. Geosyst.* 21. <https://doi.org/10.1029/2019GC008707>.
- Chinn, E.W., 1991. The role of organic geochemistry in petroleum exploration. *Basin Res. Inst. Bull.* 1, 15–23.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R.B., Piao, S., Thornton, P., 2013. Carbon and other biogeochemical cycles. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Midgley, P.M. (Eds.), *Climate Change 2013: The Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 465–570.
- Cohen, A.S., Coe, A.L., Harding, S.M., Schwark, L., 2004. Osmium isotope evidence for the regulation of atmospheric CO₂ by continental weathering. *Geology* 32, 157–160. <https://doi.org/10.1130/G20158.1>.
- Comas-Rengifo, M.J., Goy, A., 2010. Caracterización biocronoestratigráfica del Sinemuriense Superior y el Pliensbachense entre los afloramientos de la Playa Vega y Lastres (Asturias). In: García-Ramos, J.C. (Ed.), *Las Sucesiones Margo-Calcareas Marinas Del Jurásico Inferior y Las Series Fluviales Del Jurásico Superior*. Acanalados de La Playa de Vega (Ribadesella). Guía de Campo (Excursión A), V Congreso Del Jurásico de España. Servitex, Oviedo, Spain, pp. 10–18.
- Comas-Rengifo, M.J., Arias, C., Gómez, J.J., Goy, A., Herrero, C., Ossete, M.L., Palencia, A., 2010. A complementary section for the proposed Toarcian (Lower

- Jurassic) global stratotype: the Almonacid de la Cuba section (Spain). *Stratigr. Geol. Correl.* 18, 133–152. <https://doi.org/10.1134/S0869593810020036>.
- Copestake, P., Johnson, B., 2013. Lower Jurassic Foraminifera from the Llanbedr (Mochras Farm) Borehole, North Wales, UK. *Monogr. Palaeontogr. Soc.* 167, 1–403. <https://doi.org/10.1080/02693445.2013.11963952>.
- Damborenea, S.E., 2002. Jurassic evolution of Southern Hemisphere marine palaeobiogeographic units based on benthonic bivalves. *Geobios* 35, 51–71. [https://doi.org/10.1016/S0016-6995\(02\)00048-7](https://doi.org/10.1016/S0016-6995(02)00048-7).
- Dancer, P.N., Kenyon-Roberts, S.M., Downey, J.W., Baillie, J.M., Meadows, N.S., Maguire, K., 2005. The Corrib gas field, offshore west of Ireland. In: Doré, A.G., Vining, B.A. (Eds.), *Petroleum Geology: North-West Europe and Global Perspectives*, Petroleum Geology Conference Series 6. The Geological Society, London, pp. 1035–1046.
- Danisch, J., Kabiri, L., Nutz, A., Bodin, S., 2019. Chemostratigraphy of Late Sinemurian–Early Pliensbachian shallow to deep-water deposits of the Central High Atlas Basin: palaeoenvironmental implications. *J. Afr. Earth Sci.* 153, 239–249. <https://doi.org/10.1016/j.jafrearsci.2019.03.003>.
- Danise, S., Twitchett, R.J., Little, C.T.S., Clémence, M.-E., 2013. The impact of global warming and anoxia on marine benthic community dynamics: an example from the Toarcian (Early Jurassic). *PLoS One* 8, e56255. <https://doi.org/10.1371/journal.pone.0056255>.
- Danise, S., Clémence, M.-E., Price, G.D., Murphy, D.P., Gómez, J.J., Twitchett, R.J., 2019. Stratigraphic and environmental control on marine benthic community change through the early Toarcian extinction event (Iberian Range, Spain). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 524, 183–200. <https://doi.org/10.1016/j.palaeo.2019.03.039>.
- de Grazioplene, J.C., Hardenbol, J., Jacquin, T., Vail, P.R. (Eds.), 1998. *Mesozoic and Cenozoic Sequence Stratigraphy of European Basins*. SEPM Society for Sedimentary Geology, Tulsa, Oklahoma, U.S.A. <https://doi.org/10.2110/pec.98.02>
- De Lena, L.F., Taylor, D., Guex, J., Bartolini, A., Adatte, T., van Acken, D., Spangenberg, J.E., Samankassou, E., Vennemann, T., Schaltegger, U., 2019. The driving mechanisms of the carbon cycle perturbations in the late Pliensbachian (Early Jurassic). *Sci. Rep.* 9, 1–12. <https://doi.org/10.1038/s41598-019-54593-1>.
- de Oliveira, L.C.V., Rodrigues, R., Duarte, L.V., Lemos, V.B., 2006. Oil generation potential assessment and paleoenvironmental interpretation based on biomarkers and stable carbon isotopes of the Pliensbachian - lower Toarcian (Lower Jurassic) of the Peniche region (Lusitanian Basin, Portugal). *Bull. Geosci.* 14, 207–234.
- Deconinck, J.-F., Hesselbo, S.P., Debussier, N., Averbuch, O., Baudin, F., Bessa, J., 2003. Environmental controls on clay mineralogy of an Early Jurassic mudrock (Blue Lias Formation, southern England). *Int. J. Earth Sci.* 92, 255–266. <https://doi.org/10.1007/s00531-003-0318-y>.
- Deconinck, J., Hesselbo, S.P., Pellenard, P., 2019. Climatic and sea-level control of Jurassic (Pliensbachian) clay mineral sedimentation in the Cardigan Bay Basin, Llanbedr (Mochras Farm) borehole, Wales. *Sedimentology* 66, 2769–2783. <https://doi.org/10.1111/sed.12610>.
- Deconinck, J.-F., Gómez, J.J., Baudin, F., Biscay, H., Bruneau, L., Cocquerez, T., Mathieu, O., Pellenard, P., Santoni, A.-L., 2020. Diagenetic and environmental control of the clay mineralogy, organic matter and stable isotopes (C, O) of Jurassic (Pliensbachian-lowermost Toarcian) sediments of the Rodiles section (Asturian Basin, Northern Spain). *Mar. Pet. Geol.* 115, 104286. <https://doi.org/10.1016/j.marpetgeo.2020.104286>.
- Demaison, G.J., Moore, G.T., 1980. Anoxic environments and oil source bed genesis. *Org. Geochem.* 2, 9–31. [https://doi.org/10.1016/0146-6380\(80\)90017-0](https://doi.org/10.1016/0146-6380(80)90017-0).
- Dera, G., Pellenard, P., Neige, P., Deconinck, J.F., Pucéat, E., Dommergues, J.L., 2009a. Distribution of clay minerals in Early Jurassic Peritethyan seas: Palaeoclimatic significance inferred from multiproxy comparisons. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 271, 39–51. <https://doi.org/10.1016/j.palaeo.2008.09.010>.
- Dera, G., Pucéat, E., Pellenard, P., Neige, P., Delsate, D., Joachimski, M.M., Reisberg, L., Martinez, M., 2009b. Water mass exchange and variations in seawater temperature in the NW Tethys during the Early Jurassic: Evidence from neodymium and oxygen isotopes of fish teeth and belemnites. *Earth Planet. Sci. Lett.* 286, 198–207. <https://doi.org/10.1016/j.epsl.2009.06.027>.
- Dera, G., Neige, P., Dommergues, J.-L., Fara, E., Laffont, R., Pellenard, P., 2010. High-resolution dynamics of Early Jurassic marine extinctions: the case of Pliensbachian–Toarcian ammonites (Cephalopoda). *J. Geol. Soc. London.* 167, 21–33. <https://doi.org/10.1144/0016-76492009-068>.
- Dera, G., Brigaud, B., Monna, F., Laffont, R., Pucéat, E., Deconinck, J.-F., Pellenard, P., Joachimski, M.M., Durllet, C., 2011. Climatic ups and downs in a disturbed Jurassic world. *Geology* 39, 215–218. <https://doi.org/10.1130/G31579.1>.
- Dickson, A.J., Gill, B.C., Ruhl, M., Jenkyns, H.C., Porcelli, D., Idiz, E., Lyons, T.W., van den Boorn, S.H.J.M., 2017. Molybdenum-isotope chemostratigraphy and paleoceanography of the Toarcian Oceanic Anoxic Event (Early Jurassic). *Paleoceanography* 32, 813–829. <https://doi.org/10.1002/2016PA003048>.
- Duarte, L.V., Soares, A.F., 2002. Litostratigrafia das séries margo-calciárias do Jurássico inferior da Bacia Lusitânica (Portugal). *Com. Inst. Geol. Min.* 89, 135–154.
- Duarte, L.V., Silva, R.L., Oliveira, L.C.V., Comas-Rengifo, M.J., Silva, F., 2010. Organic-rich facies in the Sinemurian and Pliensbachian of the Lusitanian basin, Portugal: total organic carbon distribution and relation to transgressive-regressive facies cycles. *Geol. Acta* 8, 325–340. <https://doi.org/10.1344/105.000001536>.
- Duarte, L.V., Silva, R.L., Mendonça Filho, J.G., Poças Ribeiro, N.P., Chagas, R.B.A., 2012. High-resolution stratigraphy, palynofacies and source rock potential of the Água de Madeiros Formation (Lower Jurassic), Lusitanian Basin, Portugal. *J. Pet. Geol.* 35, 105–126. <https://doi.org/10.1111/j.1747-5457.2012.00522.x>.
- Duarte, L.V., Comas-Rengifo, M.J., Silva, R.L., Paredes, R., Goy, A., 2014. Carbon isotope stratigraphy and ammonite biostratigraphy across the Sinemurian–Pliensbachian boundary in the western Iberian margin. *Bull. Geosci.* 89, 719–736. <https://doi.org/10.3140/bull.geosci.1476>.
- Duarte, L.V., Silva, R.L., Mendonça Filho, J.G., Azerêdo, A.C., Paredes, R., 2017. Total organic carbon content and carbon stable isotopes in the Sinemurian shallow-water carbonates (Coimbra Formation) of the Lusitanian Basin, Portugal. In: IAS (Ed.), *33rd International Meeting of Sedimentology, Book of Abstracts*, Toulouse, p. 255.
- Emerson, S., Hedges, J., 2008. *Chemical Oceanography and the Marine Carbon Cycle*. Cambridge University Press.
- Espitalié, J., Laporte, J.L., Madec, M., Marquis, F., Leplat, P., Paulet, J., Boutefeu, A., 1977. Méthode rapide de caractérisation des roches mères, de leur potentiel pétrolier et de leur degré d'évolution. *Rev. l'Institut Français du Pétrole* 32, 23–42. <https://doi.org/10.2516/ogst:1977002>.
- Espitalié, J., Deroo, G., Marquis, F., 1985. La pyrolyse Rock-Eval et ses applications, Deuxième partie. *Rev. l'Institut Fr. du Pet.* 40, 755–784.
- Fantasia, A., Föllmi, K.B., Adatte, T., Bernárdez, E., Spangenberg, J.E., Mattioli, E., 2018. The Toarcian Oceanic Anoxic Event in southwestern Gondwana: an example from the Andean Basin, northern Chile. *J. Geol. Soc. Lond.* 175, 883–902. <https://doi.org/10.1144/jgs2018-008>.
- Fantasia, A., Adatte, T., Spangenberg, J.E., Font, E., Duarte, L.V., Föllmi, K.B., 2019a. Global versus local processes during the Pliensbachian–Toarcian transition at the Peniche GSSP, Portugal: a multi-proxy record. *Earth Sci. Rev.* 198, 102932. <https://doi.org/10.1016/j.earscirev.2019.102932>.
- Fantasia, A., Föllmi, K.B., Adatte, T., Spangenberg, J.E., Mattioli, E., 2019b. Expression of the Toarcian Oceanic Anoxic Event: new insights from a Swiss transect. *Sedimentology* 66, 262–284. <https://doi.org/10.1111/sed.12527>.
- Farrimond, P., Love, G.D., Bishop, A.N., Innes, H.E., Watson, D.F., Snape, C.E., 2003. Evidence for the rapid incorporation of hopanoids into kerogen. *Geochim. Cosmochim. Acta* 67, 1383–1394. [https://doi.org/10.1016/S0016-7037\(02\)01287-5](https://doi.org/10.1016/S0016-7037(02)01287-5).
- Feist-Burkhardt, S., Pross, J., 2010. Dinoflagellate cyst biostratigraphy of the opalinuston formation (middle Jurassic) in the Aalenian type area in southwest Germany and north Switzerland. *Lethaia* 43, 10–31. <https://doi.org/10.1111/j.1502-3931.2009.00170.x>.
- Fleet, A.J., Clayton, C.J., Jenkyns, H.C., Parkinson, D.N., 1987. Liassic source rock deposition in western Europe. In: Brooks, T., Glennie, K. (Eds.), *Petroleum Geology of North West Europe*. Graham & Trotman, London, pp. 59–70.
- Föllmi, K.B., Godet, A., Bodin, S., Linder, P., 2006. Interactions between environmental change and shallow water carbonate buildup along the northern Tethyan margin and their impact on the Early Cretaceous carbon isotope record. *Paleoceanography* 21. <https://doi.org/10.1029/2006PA001313>.
- Föllmi, K.B., Böle, M., Jammet, N., Froidevaux, P., Godet, A., Bodin, S., Adatte, T., Matera, V., Fleitmann, D., Spangenberg, J.E., 2012. Bridging the Faraoni and Selli oceanic anoxic events: late Hauterivian to early Aptian dysaerobic to anaerobic phases in the Tethys. *Clim. Past* 8, 171–189. <https://doi.org/10.5194/cp-8-171-2012>.
- Forsman, J.P., Hunt, J.M., 1958. Insoluble organic matter (kerogen) in sedimentary rocks. *Geochim. Cosmochim. Acta* 15, 170–182. [https://doi.org/10.1016/0016-7037\(58\)90055-3](https://doi.org/10.1016/0016-7037(58)90055-3).
- Fraguas, A., Comas-Rengifo, M.J., Gómez, J.J., Goy, A., 2012. The calcareous nannofossil crisis in Northern Spain (Asturias province) linked to the Early Toarcian warming-driven mass extinction. *Mar. Micropaleontol.* 94–95, 58–71. <https://doi.org/10.1016/j.marmicro.2012.06.004>.
- Franceschi, M., Dal Corso, J., Posenato, R., Roghi, G., Masetti, D., Jenkyns, H.C., 2014. Early Pliensbachian (Early Jurassic) C-isotope perturbation and the diffusion of the Lithiotis Fauna: insights from the western Tethys. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 410, 255–263. <https://doi.org/10.1016/j.palaeo.2014.05.025>.
- Franceschi, M., Corso, J.D., Cobianchi, M., Roghi, G., Penasa, L., Picotti, V., Preto, N., 2019. Tethyan carbonate platform transformations during the Early Jurassic (Sinemurian–Pliensbachian, Southern Alps): Comparison with the Late Triassic Carnian Pluvial Episode. *GSA Bull.* 131, 1255–1275. <https://doi.org/10.1130/B31765.1>.
- Gallois, R.W., 2008. The lithostratigraphy of the Shales-with-Beef Member of the Charmouth Mudstone Formation, Lower Jurassic. *Geosci. South-West Engl.* 12, 32–40.
- García Joral, F., Gómez, J.J., Goy, A., 2011. Mass extinction and recovery of the Early Toarcian (Early Jurassic) brachiopods linked to climate change in Northern and Central Spain. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 302, 367–380. <https://doi.org/10.1016/j.palaeo.2011.01.023>.
- Gómez, J.J., Arias, C., 2010. Rapid warming and ostracods mass extinction at the Lower Toarcian (Jurassic) of central Spain. *Mar. Micropaleontol.* 74, 119–135. <https://doi.org/10.1016/j.marmicro.2010.02.001>.
- Gómez, J.J., Goy, A., 2011. Warming-driven mass extinction in the Early Toarcian (Early Jurassic) of northern and central Spain. Correlation with other time-equivalent European sections. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 306, 176–195. <https://doi.org/10.1016/j.palaeo.2011.04.018>.
- Gómez, J.J., Goy, A., Canales, M.L., 2008. Seawater temperature and carbon isotope variations in belemnites linked to mass extinction during the Toarcian (Early Jurassic) in Central and Northern Spain. Comparison with other European sections. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 258, 28–58. <https://doi.org/10.1016/j.palaeo.2007.11.005>.
- Gómez, J.J., Comas-Rengifo, M.J., Goy, A., 2016a. The hydrocarbon source rocks of the Pliensbachian (Early Jurassic) in the Asturian Basin (Northern Spain): their relationship with the palaeoclimatic oscillations and gamma-ray response. *J. Iber. Geol.* 42, 259–273. <https://doi.org/10.5209/JIGE.53265>.

- Gómez, J.J., Comas-Rengifo, M.J., Goy, A., 2016b. Palaeoclimatic oscillations in the Pliensbachian (Early Jurassic) of the Asturian Basin (Northern Spain). *Clim. Past* 12, 1199–1214. <https://doi.org/10.5194/cp-12-1199-2016>.
- Goodarzi, F., Gentzls, T., Embry, A.F., Osadetz, K.G., Skibo, D.N., Stewart, K.R., 1993. Evaluation of maturity and source rock potential in the Lougheed Island area of the central Sverdrup Basin, Arctic Canada. In: *Vorren, T.O., Bergsager, E., Dahl-Stammes, Ø.A., Holter, E., Johansen, B., Lie, E., Lund, T.B. (Eds.), Arctic Geology and Petroleum Potential*. Norwegian Petroleum Society Special Publications, Elsevier, pp. 147–157. <https://doi.org/10.1016/B978-0-444-88943-0.50014-9>.
- Hallam, A., 1981. A revised sea-level curve for the early Jurassic. *J. Geol. Soc. Lond.* 138, 735–743. <https://doi.org/10.1144/gsjgs.138.6.0735>.
- Hallam, A., Bradshaw, M.J.J., 1979. Bituminous shales and oolitic ironstones as indicators of transgressions and regressions. *J. Geol. Soc. Lond.* 136, 157–164. <https://doi.org/10.1144/gsjgs.136.2.0157>.
- Han, Z., Hu, X., Kemp, D.B., Li, J., 2018. Carbonate-platform response to the Toarcian Oceanic Anoxic Event in the southern hemisphere: implications for climatic change and biotic platform demise. *Earth Planet. Sci. Lett.* 489, 59–71. <https://doi.org/10.1016/j.epsl.2018.02.017>.
- Hayes, J.M., Strauss, H., Kaufman, A.J., 1999. The abundance of ^{13}C in marine organic matter and isotopic fractionation in the global biogeochemical cycle of carbon during the past 800 Ma. *Chem. Geol.* 161, 103–125. [https://doi.org/10.1016/S0009-2541\(99\)00083-2](https://doi.org/10.1016/S0009-2541(99)00083-2).
- Hedges, J.L., Keil, R.G., 1995. Sedimentary organic matter preservation: an assessment and speculative synthesis. *Mar. Chem.* 49, 81–115. [https://doi.org/10.1016/0304-4203\(95\)00008-F](https://doi.org/10.1016/0304-4203(95)00008-F).
- Hemingway, J.D., Rothman, D.H., Grant, K.E., Rosengard, S.Z., Eglinton, T.I., Derry, L.A., Galy, V.V., 2019. Mineral protection regulates long-term global preservation of natural organic carbon. *Nature* 570, 228–231. <https://doi.org/10.1038/s41586-019-1280-6>.
- Hermoso, M., Le Callonnec, L., Minoletti, F., Renard, M., Hesselbo, S.P., 2009. Expression of the Early Toarcian negative carbon-isotope excursion in separated carbonate microfractions (Jurassic, Paris Basin). *Earth Planet. Sci. Lett.* 277, 194–203. <https://doi.org/10.1016/j.epsl.2008.10.013>.
- Hermoso, M., Minoletti, F., Rickaby, R.E.M., Hesselbo, S.P., Baudin, F., Jenkyns, H.C., 2012. Dynamics of a stepped carbon-isotope excursion: ultra high-resolution study of Early Toarcian environmental change. *Earth Planet. Sci. Lett.* 319–320, 45–54. <https://doi.org/10.1016/j.epsl.2011.12.021>.
- Hermoso, M., Minoletti, F., Pellenard, P., 2013. Black shale deposition during Toarcian super-greenhouse driven by sea level. *Clim. Past Discuss.* 9, 4365–4384. <https://doi.org/10.5194/cpd-9-4365-2013>.
- Hermoso, M., Delsate, D., Baudin, F., Le Callonnec, L., Minoletti, F., Renard, M., Faber, A., 2014. Record of Early Toarcian carbon cycle perturbations in a nearshore environment: the Bascharage section (easternmost Paris Basin). *Solid Earth* 5, 793–804. <https://doi.org/10.5194/se-5-793-2014>.
- Herrle, J.O., Köbler, P., Friedrich, O., Erlenkeuser, H., Hemleben, C., 2004. High-resolution carbon isotope records of the Aptian to Lower Albian from SE France and the Mazagan Plateau (DSDP Site 545): a stratigraphic tool for paleoceanographic and paleobiologic reconstruction. *Earth Planet. Sci. Lett.* 218, 149–161. [https://doi.org/10.1016/S0012-821X\(03\)00646-0](https://doi.org/10.1016/S0012-821X(03)00646-0).
- Herrle, J.O., Schröder-Adams, C.J., Davis, W., Pugh, A.T., Galloway, J.M., Fath, J., 2015. Mid-Cretaceous High Arctic stratigraphy, climate, and Oceanic Anoxic Events. *Geology* 43, 403–406. <https://doi.org/10.1130/G36439.1>.
- Hesselbo, S.P., Coe, A.L., 2000. Jurassic Sequences of the Hebrides Basin, Isle of Skye, Scotland, Field Trip Guidebook, International Association of Sedimentologists Regional Meeting, Dublin.
- Hesselbo, S.P., Oates, M.J., Jenkyns, H.C., 1998. The lower Lias Group of the Hebrides Basin. *Scott. J. Geol.* 34, 23–60. <https://doi.org/10.1144/sjg34010023>.
- Hesselbo, S.P., Gröcke, D.R., Jenkyns, H.C., Bjerrum, C.J., Farrimond, P., Morgans Bell, H.S., Green, O.R., 2000a. Massive dissociation of gas hydrate during a Jurassic oceanic anoxic event. *Nature* 406, 392–395. <https://doi.org/10.1038/35019044>.
- Hesselbo, S.P., Meister, C., Gröcke, D.R., 2000b. A potential global stratotype for the Sinemurian–Pliensbachian boundary (Lower Jurassic), Robin Hood’s Bay, UK: ammonite faunas and isotope stratigraphy. *Geol. Mag.* 137, 601–607. <https://doi.org/10.1017/S0016756800004672>.
- Hesselbo, S.P., Jenkyns, H.C., Duarte, L.V., Oliveira, L.C.V., 2007. Carbon-isotope record of the Early Jurassic (Toarcian) Oceanic Anoxic Event from fossil wood and marine carbonate (Lusitanian Basin, Portugal). *Earth Planet. Sci. Lett.* 253, 455–470. <https://doi.org/10.1016/j.epsl.2006.11.009>.
- Hesselbo, S.P., Hudson, A.J.L., Huggett, J.M., Leng, M.J., Riding, J.B., Ullmann, C.V., 2020. Palynological, geochemical, and mineralogical characteristics of the Early Jurassic Liasidium Event in the Cleveland Basin, Yorkshire, UK. *Newslett. Stratigr.* 53, 191–211. <https://doi.org/10.1127/nos/2019/0536>.
- Hinnov, L.A., Park, J.J., 1999. Strategies for assessing Early-Middle (Pliensbachian–Aalenian) Jurassic cyclochronologies. *Philos. Trans. R. Soc. Lond. A* 357, 1831–1859. <https://doi.org/10.1098/rsta.1999.0403>.
- Holmden, C., Creaser, R.A., Muehlenbachs, K., Leslie, S.A., Bergström, S.M., 1998. Isotopic evidence for geochemical decoupling between ancient epicric seas and bordering oceans: implications for secular curves. *Geology* 26, 567–570. [https://doi.org/10.1130/0091-7613\(1998\)026<0567:IEFGDB>2.3.CO;2](https://doi.org/10.1130/0091-7613(1998)026<0567:IEFGDB>2.3.CO;2).
- Huc, A.Y., Durand, B., 1974. Etude des acides humiques et de l’humine de sédiments récents considérés comme précurseurs des kéroènes. In: *Tissot, B., Biener, F. (Eds.), Advances in Organic Geochemistry*. Technip, Paris, pp. 53–72.
- Hülse, D., Arndt, S., Ridgwell, A., 2019. Mitigation of extreme ocean anoxic event conditions by organic matter sulfurization. *Paleoceanogr. Palaeoclimatol.* 34, 476–489. <https://doi.org/10.1029/2018PA003470>.
- Hunt, J.M., 1995. *Petroleum Geochemistry and Geology*, A Series of Books in Geology. Freeman, W. H., New York [https://doi.org/10.1016/S0009-2541\(96\)00131-3](https://doi.org/10.1016/S0009-2541(96)00131-3).
- Immenhauser, A., Della Porta, G., Kenter, J.A.M., Bahamonde, J.R., 2003. An alternative model for positive shifts in shallow-marine carbonate $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$. *Sedimentology* 50, 953–959. <https://doi.org/10.1046/j.1365-3091.2003.00590.x>.
- Ivimey-Cook, H.C., 1971. Stratigraphical palaeontology of the Lower Jurassic of the Llanbedr (Mochras Farm) Borehole. In: *Woodland, W.A. (Ed.), The Llanbedr (Mochras Farm) Borehole*. Rep. No. 71/18. Institute of Geological Sciences, pp. 87–92.
- Jarvie, D.M., 2012. Shale resource systems for oil and gas: part 1—shale-gas resource systems. In: *Breyer, J.A. (Ed.), Shale Reservoirs—Giant Resources for the 21st Century*, AAPG Memoir 97. American Association of Petroleum Geologists, pp. 69–87. <https://doi.org/10.1306/13321446M973489>.
- Jarvis, I., Gale, A.S., Jenkyns, H.C., Pearce, M.A., 2006. Secular variation in Late Cretaceous carbon isotopes: a new $\delta^{13}\text{C}$ carbonate reference curve for the Cenomanian–Campanian (99.6–70.6 Ma). *Geol. Mag.* 143, 561–608. <https://doi.org/10.1017/S0016756806002421>.
- Jenkyns, H.C., 1980. Cretaceous anoxic events: from continents to oceans. *J. Geol. Soc. Lond.* 137, 171–188. <https://doi.org/10.1144/gsjgs.137.2.0171>.
- Jenkyns, H.C., 1988. The early Toarcian (Jurassic) anoxic event; stratigraphic, sedimentary and geochemical evidence. *Am. J. Sci.* 288, 101–151. <https://doi.org/10.1126/science.3.53.32>.
- Jenkyns, H.C., 2010. Geochemistry of oceanic anoxic events. *Geochem. Geophys. Geosyst.* 11, 1–30. <https://doi.org/10.1029/2009GC002788>.
- Jenkyns, H.C., Clayton, C.J., 1986. Black shales and carbon isotopes in pelagic sediments from the Tethyan Lower Jurassic. *Sedimentology* 33, 87–106. <https://doi.org/10.1111/j.1365-3091.1986.tb00746.x>.
- Jenkyns, H., Clayton, C., 1997. Lower Jurassic epicontinental carbonates and mudstones from England and Wales: chemostratigraphic signals and the early Toarcian anoxic event. *Sedimentology* 44, 687–706. <https://doi.org/10.1046/j.1365-3091.1997.d0143.x>.
- Jenkyns, H.C., Weedon, G.P., 2013. Chemostratigraphy (CaCO₃, TOC, $\delta^{13}\text{C}_{\text{org}}$) of Sinemurian (Lower Jurassic) black shales from the Wessex Basin, Dorset and palaeoenvironmental implications. *Newslett. Stratigr.* 46, 1–21. <https://doi.org/10.1127/0078-0421/2013/0029>.
- Jenkyns, H.C., Jones, C., Gröcke, D., Hesselbo, S., Parkinson, D., 2002. Chemostratigraphy of the Jurassic System: applications, limitations and implications for palaeoceanography. *J. Geol. Soc. Lond.* 159, 351–378. <https://doi.org/10.1144/0016-764901-130>.
- Jin, X., Shi, Z., Baranyi, V., Kemp, D.B., Han, Z., Luo, G., Hu, J., He, F., Chen, L., Preto, N., 2020. The Jenkyns Event (early Toarcian OAE) in the Ordos Basin, North China. *Glob. Planet. Change* 193, 103273. <https://doi.org/10.1016/j.gloplacha.2020.103273>.
- Jones, R.W., 1987. Organic facies. In: *Brooks, J., Welte, D. (Eds.), Advances in Petroleum Geochemistry*. Academic Press, London, pp. 78–80.
- Kafousia, N., Karakitsios, V., Jenkyns, H.C., Mattioli, E., 2011. A global event with a regional character: the Early Toarcian Oceanic Anoxic Event in the Pindos Ocean (northern Peloponnese, Greece). *Geol. Mag.* 148, 619–631. <https://doi.org/10.1017/S0016756811000082>.
- Kafousia, N., Karakitsios, V., Mattioli, E., Kenjo, S., Jenkyns, H.C., 2014. The Toarcian Oceanic Anoxic Event in the Ionian zone. *Greece. Palaeogeogr. Palaeoclimatol. Palaeoecol.* 393, 135–145. <https://doi.org/10.1016/j.palaeo.2013.11.013>.
- Katz, B., 1995. Petroleum source rocks — an introductory overview. In: *Katz, B.J. (Ed.), Petroleum Source Rocks, Casebooks in Earth Sciences*. Springer, Berlin, Heidelberg, pp. 1–8. <https://doi.org/10.1007/978-3-642-78911-3>.
- Kemp, D.B., Izumi, K., 2014. Multiproxy geochemical analysis of a Panthalassic margin record of the early Toarcian oceanic anoxic event (Toyora area, Japan). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 414, 332–341. <https://doi.org/10.1016/j.palaeo.2014.09.019>.
- Kemp, D.B., Coe, A.L., Cohen, A.S., Schwark, L., 2005. Astronomical pacing of methane release in the Early Jurassic period. *Nature* 437, 396–399. <https://doi.org/10.1038/nature04361>.
- Kennedy, M.J., Wagner, T., 2011. Clay mineral continental amplifier for marine carbon sequestration in a greenhouse ocean. *Proc. Natl. Acad. Sci.* 108, 9776–9781. <https://doi.org/10.1073/pnas.1018670108>.
- Kennedy, M.J., Pevear, D.R., Hill, R.J., 2002. Mineral surface control of organic carbon in black shale. *Science* (80-). <https://doi.org/10.1126/science.1066611>.
- Kodina, L.A., Bogatcheva, M.P., Lobitzer, H., 1988. An organic geochemical study of austrian bituminous rocks. *Jahrb. der Geol. Bundesanstalt* 131, 291–300.
- Kohnen, M.E.L., Damsté, J.S.S., Kock-van Dalen, A.C., Haven, H.L.T., Rullkötter, J., De Leeuw, J.W., 1990. Origin and diagenetic transformations of C25 and C30 highly branched isoprenoid sulphur compounds: Further evidence for the formation of organically bound sulphur during early diagenesis. *Geochim. Cosmochim. Acta*. [https://doi.org/10.1016/0016-7037\(90\)90121-Z](https://doi.org/10.1016/0016-7037(90)90121-Z).
- Kohnen, M.E.L., Schouten, S., Sinnighe Damsté, J.S., de Leeuw, J.W., Merritt, D., Hayes, J.M., 1992. The combined application of organic sulphur and isotope geochemistry to assess multiple sources of palaeobiochemicals with identical carbon skeletons. *Org. Geochem.* 19, 403–419. [https://doi.org/10.1016/0146-6380\(92\)90008-L](https://doi.org/10.1016/0146-6380(92)90008-L).
- Kontorovich, A.E., Moskvina, V.I., Bostrikov, O.I., Danilova, V.P., Fomin, A.N., Fomichev, A.S., Kostyeva, E.A., Melenevsky, V.N., 1997. Main oil source formations of the West Siberian Basin. *Pet. Geosci.* 3, 343–358. <https://doi.org/10.1144/ptegco.3.4.343>.
- Korte, C., Hesselbo, S.P., 2011. Shallow marine carbon and oxygen isotope and elemental records indicate icehouse-greenhouse cycles during the Early Jurassic. *Paleoceanogr. Palaeoclimatol.* 26, 1–18. <https://doi.org/10.1029/2011PA002160>.

- Korte, C., Hesselbo, S.P., Ullmann, C.V., Dietl, G., Ruhl, M., Schweigert, G., Thibault, N., 2015. Jurassic climate mode governed by ocean gateway. *Nat. Commun.* 6, 10015. <https://doi.org/10.1038/ncomms10015>.
- Krencker, F.N., Bodin, S., Hoffmann, R., Suan, G., Mattioli, E., Kabiri, L., Föllmi, K.B., Immenhauser, A., 2014. The middle Toarcian cold snap: trigger of mass extinction and carbonate factory demise. *Glob. Planet. Change* 117, 64–78. <https://doi.org/10.1016/j.gloplacha.2014.03.008>.
- Krencker, F.-N., Lindström, S., Bodin, S., 2019. A major sea-level drop briefly precedes the Toarcian oceanic anoxic event: implication for Early Jurassic climate and carbon cycle. *Sci. Rep.* 9, 12518. <https://doi.org/10.1038/s41598-019-48956-x>.
- Kump, L.R., Arthur, M.A., 1999. Interpreting carbon-isotope excursions: carbonates and organic matter. *Chem. Geol.* 161, 181–198. [https://doi.org/10.1016/S0009-2541\(99\)00086-8](https://doi.org/10.1016/S0009-2541(99)00086-8).
- Küspert, W., 1982. Environmental changes during oil shale deposition as deduced from stable isotope ratios. In: Einsele, G., Seilacher, A. (Eds.), *Cyclic and Event Stratification*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 482–501. https://doi.org/10.1007/978-3-642-75829-4_36.
- Largeau, C., Derenne, S., 1993. Relative efficiency of the Selective Preservation and Degradation Recondensation pathways in kerogen formation. Source and environment influence on their contributions to type I and II kerogens. *Org. Geochem.* 20, 611–615. [https://doi.org/10.1016/0146-6380\(93\)90027-9](https://doi.org/10.1016/0146-6380(93)90027-9).
- Law, C.A., 1999. Evaluating source rocks. In: Beaumont, E.A., Foster, N.H. (Eds.), *Exploring for Oil and Gas Traps*. American Association of Petroleum Geologists. <https://doi.org/10.1306/TrHbk624C6>, pp. 34 1–3.
- Leckie, R.M., Bralower, T.J., Cashman, R., 2002. Oceanic anoxic events and plankton evolution: Biotic response to tectonic forcing during the mid-Cretaceous. *Paleoceanography* 17. <https://doi.org/10.1029/2001PA000623>, 13-1-13–29.
- Lee, C., Wakeham, S., Arnosti, C., 2004. Particulate organic matter in the sea: the composition conundrum. *AMBIO A J. Hum. Environ.* 33, 565–575. <https://doi.org/10.1579/0044-7447-33.8.565>.
- Lewan, M.D., 1987. Petrographic study of primary petroleum migration in the Woodford Shale and related rock units. In: Doligez, B. (Ed.), *Collection Colloques et Séminaires-Institut Français Du Pétrole*. Editions Technip, Paris, pp. 113–130.
- Lézin, C., Andreu, B., Pellenard, P., Bouchez, J.L., Emmanuel, L., Fauré, P., Landrein, P., 2013. Geochemical disturbance and paleoenvironmental changes during the Early Toarcian in NW Europe. *Chem. Geol.* 341, 1–15. <https://doi.org/10.1016/j.chemgeo.2013.01.003>.
- Little, C.T.S., Benton, M.J., 1995. Early Jurassic mass extinction: a global long-term event. *Geology* 23, 495. [https://doi.org/10.1130/0091-7613\(1995\)023<0495:EJMEAG>2.3.CO;2](https://doi.org/10.1130/0091-7613(1995)023<0495:EJMEAG>2.3.CO;2).
- Littler, K., Hesselbo, S.P., Jenkyns, H.C., 2010. A carbon-isotope perturbation at the Pliensbachian–Toarcian boundary: evidence from the Lias Group, NE England. *Geol. Mag.* 147, 181–192. <https://doi.org/10.1017/S0016756809990458>.
- Marshall, J.D., 1992. Climatic and oceanographic isotopic signals from the carbonate rock record and their preservation. *Geol. Mag.* 129, 143. <https://doi.org/10.1017/S0016756800008244>.
- Martindale, R.C., Them, T.R., Gill, B.C., Marroquín, S.M., Knoll, A.H., 2017. A new Early Jurassic (ca. 183 Ma) fossil Lagerstätte from Ya Ha Tinda, Alberta, Canada. *Geology* 45, 255–258. <https://doi.org/10.1130/G38808.1>.
- Martinez, M., Deconinck, J.-F., Pellenard, P., Riquier, L., Company, M., Reboulet, S., Moiroud, M., 2015. Astrochronology of the Valanginian–Hauterivian stages (Early Cretaceous): Chronological relationships between the Paraná–Etendeka large igneous province and the Weissert and the Faraoni events. *Glob. Planet. Change* 131, 158–173. <https://doi.org/10.1016/j.gloplacha.2015.06.001>.
- Martinez, M., Krencker, F.-N., Mattioli, E., Bodin, S., 2017. Orbital chronology of the Pliensbachian – Toarcian transition from the Central High Atlas Basin (Morocco). *Newslett. Stratigr.* 50, 47–69. <https://doi.org/10.1127/nos/2016/0311>.
- Masetti, D., Figus, B., Jenkyns, Barattolo, F., Mattioli, E., Posenato, R., 2017. Carbon-isotope anomalies and demise of carbonate platforms in the Sinemurian (Early Jurassic) of the Tethyan region: evidence from the Southern Alps (Northern Italy). *Geol. Mag.* 154, 625–650. <https://doi.org/10.1017/S0016756816000273>.
- Mattioli, E., Pittet, B., Petitpierre, L., Mailliot, S., 2009. Dramatic decrease of pelagic carbonate production by nannoplankton across the Early Toarcian anoxic event (T-OAE). *Glob. Planet. Change* 65, 134–145. <https://doi.org/10.1016/j.gloplacha.2008.10.018>.
- Mayer, L.M., 1994. Relationships between mineral surfaces and organic carbon concentrations in soils and sediments. *Chem. Geol.* 114, 347–363. [https://doi.org/10.1016/0009-2541\(94\)90063-9](https://doi.org/10.1016/0009-2541(94)90063-9).
- McArthur, J.M., Algeo, T.J., van de Schootbrugge, B., Li, Q., Howarth, R.J., 2008. Basinal restriction, black shales, Re-Os dating, and the Early Toarcian (Jurassic) oceanic anoxic event. *Paleoceanography* 23, PA4217. <https://doi.org/10.1029/2008PA001607>.
- McElwain, J.C., Wade-Murphy, J., Hesselbo, S.P., 2005. Changes in carbon dioxide during an oceanic anoxic event linked to intrusion into Gondwana coals. *Nature* 435, 479–482. <https://doi.org/10.1038/nature03618>.
- Menning, M., Hendrich, A., 2002. *Stratigraphische Tabelle von Deutschland 2002, Deutsche Stratigraphische Kommission: Stratigraphische Tabelle von Deutschland*.
- Mercuzot, M., Pellenard, P., Durllet, C., Bougeault, C., Meister, C., Dommergues, J.-L., Thibault, N., Baudin, F., Mathieu, O., Bruneau, L., Huret, E., El Hmidi, K., 2020. Carbon-isotope events during the Pliensbachian (Lower Jurassic) on the African and European margins of the NW Tethyan Realm. *Newslett. Stratigr.* 53, 41–69. <https://doi.org/10.1127/nos/2019/0502>.
- Mitchell, S.F., Paul, C.R.C., Gale, A.S., 1996. Carbon isotopes and sequence stratigraphy. In: Howell, J.A., Aitken, J.F. (Eds.), *High Resolution Sequence Stratigraphy: Innovations and Applications*, Geological Society, London, Special Publications 104. Geological Society, London, pp. 11–24.
- Monaco, P., Nocchi, M., Ortega-Huertás, M., Palomo, I., Martínez, F., Chiavini, G., 1994. Depositional trends in the Valdorbia Section (Central Italy) during the Early Jurassic, as revealed by micropaleontology, sedimentology and geochemistry. *Ecolgae Geo. Helv.* 87, 157–223. <https://doi.org/10.1015/0157-67>.
- Montero-Serrano, J.C., Föllmi, K.B., Adatte, T., Spangenberg, J.E., Tribouillard, N., Fantasia, A., Suan, G., 2015. Continental weathering and redox conditions during the early Toarcian Oceanic Anoxic Event in the northwestern Tethys: insight from the Posidonia Shale section in the Swiss Jura Mountains. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 429, 83–99. <https://doi.org/10.1016/j.palaeo.2015.03.043>.
- Morton, N., 2004. The Hebrides Basin. In: Simms, M.J., Chidlaw, N., Morton, N., Page, K. N. (Eds.), *British Lower Jurassic Stratigraphy*. Geological Conservation Review Series, Peterborough, pp. 315–375.
- Mouterde, R., Tintant, H., Allou, J., Gabbily, J., Hanzo, M., Lefavrais, A., Rioult, M., 1980. Lias. In: Mégnien, C., Mégnien, F. (Eds.), *Synthèse Géologique Du Bassin de Paris*. Bureau de Recherches Géologiques et Minières, Paris, pp. 76–121.
- Müller, T., Jurikova, H., Gutjahr, M., Tomašových, A., Schlögl, J., Liebetrau, V., Duarte, L.V., Milovský, R., Suan, G., Mattioli, E., Pittet, B., Eisenhauer, A., 2020. Ocean acidification during the early Toarcian extinction event: evidence from boron isotopes in brachiopods. *Geology* 48, 1184–1188. <https://doi.org/10.1130/G47781.1>.
- Mutterlose, J., Bornemann, A., Herrle, J., 2009. The Aptian – Albian cold snap: evidence for “mid” Cretaceous icehouse interludes. *Neues Jahrb. für Geol. Paläontol. Abhandlungen* 252, 217–225. <https://doi.org/10.1127/0077-7749/2009/0252-0217>.
- Newton, R., Bottrell, S., 2007. Stable isotopes of carbon and sulphur as indicators of environmental change: past and present. *J. Geol. Soc. Lond.* 164, 691–708.
- Ogg, J.G., Ogg, G.M., Gradstein, F.M., 2016. A Concise Geologic Time Scale. <https://doi.org/10.1016/c2009-0-64442-1>.
- Pálffy, J., Smith, P.L., 2000. Synchrony between Early Jurassic extinction, oceanic anoxic event, and the Karoo-Ferrar flood basalt volcanism. *Geology* 28, 747–750. [https://doi.org/10.1130/0091-7613\(2000\)028<0747:SBEJEO>2.3.CO;2](https://doi.org/10.1130/0091-7613(2000)028<0747:SBEJEO>2.3.CO;2).
- Panchuk, K.M., Holmden, C.E., Leslie, S.A., 2006. Local controls on carbon cycling in the ordoevian midcontinent region of North America, with implications for carbon isotope secular curves. *J. Sediment. Res.* 76, 200–211. <https://doi.org/10.2110/jsr.2006.017>.
- Pancost, R.D., Crawford, N., Magness, S., Turner, A., Jenkyns, H.C., Maxwell, J.R., 2004. Further evidence for the development of photic-zone euxinic conditions during Mesozoic oceanic anoxic events. *J. Geol. Soc. Lond.* 161, 353–364. <https://doi.org/10.1144/0016764903-059>.
- Parisi, G., Ortega-Huertás, M., Nocchi, M., Palomo, I., Monaco, P., Martínez, F., 1996. Stratigraphic and geochemical anomalies of the early Toarcian oxygen-poor interval in the Umbria-Marche Apennines (Italy). *Geobios* 29, 469–484. [https://doi.org/10.1016/S0016-6995\(96\)80006-4](https://doi.org/10.1016/S0016-6995(96)80006-4).
- Patterson, W.P., Walter, L.M., 1994. Depletion of ¹³C in seawater ^ΣCO₂ on modern carbonate platforms: significance for the carbon isotopic record of carbonates. *Geology* 22, 885–888. [https://doi.org/10.1130/0091-7613\(1994\)022<0885](https://doi.org/10.1130/0091-7613(1994)022<0885).
- Pearce, C.R., Cohen, A.S., Coe, A.L., Burton, K.W., 2008. Molybdenum isotope evidence for global ocean anoxia coupled with perturbations to the carbon cycle during the early Jurassic. *Geology* 36, 231–234. <https://doi.org/10.1130/G24446A.1>.
- Pedersen, T.F., Calvert, S.E., 1990. Anoxia vs. productivity: what controls the formation of organic-carbon-rich sediments and sedimentary rocks? *Am. Assoc. Pet. Geol. Bull.* 74, 454–466. <https://doi.org/10.1306/0C9B232B-1710-11D7-8645000102C1865D>.
- Percival, L.M.E., Witt, M.L.L., Mather, T.A., Hermoso, M., Jenkyns, H.C., Hesselbo, S.P., Al-Suwaidi, A.H., Storm, M.S., Xu, W., Ruhl, M., 2015. Globally enhanced mercury deposition during the end-Pliensbachian extinction and Toarcian OAE: a link to the Karoo-Ferrar Large Igneous Province. *Earth Planet. Sci. Lett.* 428, 267–280. <https://doi.org/10.1016/j.epsl.2015.06.064>.
- Percival, L.M.E., Cohen, A.S., Davies, M.K., Dickson, A.J., Hesselbo, S.P., Jenkyns, H.C., Leng, M.J., Mather, T.A., Storm, M.S., Xu, W., 2016. Osmium isotope evidence for two pulses of increased continental weathering linked to Early Jurassic volcanism and climate change. *Geology* 44, 759–762. <https://doi.org/10.1130/G37997.1>.
- Peters, K.E., Walters, C.C., Moldowan, J.M., 2005. *The Biomarker Guide*. Cambridge University Press.
- Peti, L., Thibault, N., 2017. Abundance and size changes in the calcareous nannofossil *Schizosphaerella* – Relation to sea-level, the carbonate factory and palaeoenvironmental change from the Sinemurian to earliest Toarcian of the Paris Basin. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 485, 271–282. <https://doi.org/10.1016/j.palaeo.2017.06.019>.
- Peti, L., Thibault, N., Clémence, M.-E., Korte, C., Dommergues, J.-L., Bougeault, C., Pellenard, P., Jelby, M.E., Ullmann, C.V., 2017. Sinemurian–Pliensbachian calcareous nannofossil biostratigraphy and organic carbon isotope stratigraphy in the Paris Basin: calibration to the ammonite biozonation of NW Europe. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 468, 142–161. <https://doi.org/10.1016/j.palaeo.2016.12.004>.
- Phillip, R.P., Calvin, M., 1976. Possible origin for insoluble organic (kerogen) debris in sediments from insoluble cell-wall materials of algae and bacteria. *Nature* 262, 134–136. <https://doi.org/10.1038/262134a0>.
- Piazza, V., Duarte, L.V., Renaudie, J., Aberhan, M., 2019. Reductions in body size of benthic macroinvertebrates as a precursor of the early Toarcian (Early Jurassic) extinction event in the Lusitanian Basin, Portugal. *Paleobiology* 45, 296–316. <https://doi.org/10.1017/pab.2019.11>.
- Pienkowski, G., Schudack, M.E., Bosák, P., Enay, R., Feldman-Olszewska, A., Golonka, J., Gutowski, J., Hergreen, G.F.W., Jordan, P., Krobicki, M., Lathuilière, B., Leinfelder, R.R., Michalik, J., Mönnig, E., Noe-Nygaard, N., Pálffy, J., Pint, A., Rasser, M.W., Reisdorf, A.G., Schmid, D.U., Schweigert, G., Surlyk, F., Wetzal, A., Wong, T.E., 2008. Jurassic. In: McCann, T. (Ed.), *The Geology of Central Europe*

- Volume 2: Mesozoic and Cenozoic. Geological Society of London, London, pp. 823–922. <https://doi.org/10.1144/CEV2P.2>.
- Pienkowski, G., Hodob, M., Ullmann, C.V., 2016. Fungal decomposition of terrestrial organic matter accelerated Early Jurassic climate warming. *Sci. Rep.* 6, 319–330. <https://doi.org/10.1038/srep31930>.
- Pittet, B., Suan, G., Lenoir, F., Duarte, L.V., Mattioli, E., 2014. Carbon isotope evidence for sedimentary discontinuities in the lower Toarcian of the Lusitanian Basin (Portugal): sea level change at the onset of the Oceanic Anoxic Event. *Sediment. Geol.* 303, 1–14. <https://doi.org/10.1016/j.sedgeo.2014.01.001>.
- Poças Ribeiro, N., Mendonça Filho, J.G., Duarte, L.V., Silva, R.L., Mendonça, J.O., Silva, T.F., 2013. Palynofacies and organic geochemistry of the Sinemurian carbonate deposits in the western Lusitanian Basin (Portugal): Coimbra and Água de Madeiros formations. *Int. J. Coal Geol.* 111, 37–52. <https://doi.org/10.1016/j.coal.2012.12.006>.
- Porter, S.J., Selby, D., Suzuki, K., Gröcke, D., 2013. Opening of a trans-Pangaean marine corridor during the Early Jurassic: insights from osmium isotopes across the Sinemurian–Pliensbachian GSSP, Robin Hood’s Bay, UK. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 375, 50–58. <https://doi.org/10.1016/j.palaeo.2013.02.012>.
- Porter, S.J., Selby, D., Cameron, V., 2014a. Characterising the nickel isotopic composition of organic-rich marine sediments. *Chem. Geol.* 387, 12–21. <https://doi.org/10.1016/j.chemgeo.2014.07.017>.
- Porter, S.J., Smith, P.L., Caruthers, A.H., Hou, P., Gröcke, D.R., Selby, D., 2014b. New high resolution geochemistry of Lower Jurassic marine sections in western North America: a global positive carbon isotope excursion in the Sinemurian? *Earth Planet. Sci. Lett.* 397, 19–31. <https://doi.org/10.1016/j.epsl.2014.04.023>.
- Powell, T.G., 1978. An assessment of the hydrocarbon source rock potential of the Canadian Arctic islands. In: *Geological Survey Paper* 78-12. <https://doi.org/10.4095/103606>.
- Price, G., 1999. The evidence and implications of polar ice during the Mesozoic. *Earth Sci. Rev.* 48, 183–210. [https://doi.org/10.1016/S0012-8252\(99\)00048-3](https://doi.org/10.1016/S0012-8252(99)00048-3).
- Price, G.D., Baker, S.J., VanDeVelde, J., Clémence, M.-E., 2016. High-resolution carbon cycle and seawater temperature evolution during the Early Jurassic (Sinemurian–Early Pliensbachian). *Geochim. Geophys. Res.* 17, 3917–3928. <https://doi.org/10.1002/2016GC006541>.
- Quesada, S., Dorransoro, C., Robles, S., Chalor, R., Grimalt, J.O., 1997. Geochemical correlation of oil from the Ayoluengo field to Liassic black shale units in the southwestern Basque-Cantabrian Basin (northern Spain). *Org. Geochem.* 27, 25–40. [https://doi.org/10.1016/S0146-6380\(97\)00045-4](https://doi.org/10.1016/S0146-6380(97)00045-4).
- Quesada, S., Robles, S., Rosales, I., 2005. Depositional architecture and transgressive-regressive cycles within Liassic backstepping carbonate ramps in the Basque-Cantabrian basin, northern Spain. *J. Geol. Soc. Lond.* 162, 531–548. <https://doi.org/10.1144/0016-764903-041>.
- Rachidi, M., Neuweiler, F., Kirkwood, D., 2009. Diagenetic-geochemical patterns and fluid evolution history of a Lower Jurassic petroleum source rock, Middle Atlas, Morocco. *J. Pet. Geol.* 32, 111–128. <https://doi.org/10.1111/j.1747-5457.2009.00439.x>.
- Rasmussen, E.S., Lomholt, S., Andersen, C., Vejbæk, O.V., 1998. Aspects of the structural evolution of the Lusitanian Basin in Portugal and the shelf and slope area offshore Portugal. *Tectonophysics* 300, 199–225. [https://doi.org/10.1016/S0040-1951\(98\)00241-8](https://doi.org/10.1016/S0040-1951(98)00241-8).
- Rees, P.M., Ziegler, A.M., Valdes, P.J., 1999. Jurassic phytogeography and climates: new data and model comparisons. In: Huber, B.T., Macleod, K.G., Wing, S.L. (Eds.), *Warm Climates in Earth History*. Cambridge University Press, Cambridge, pp. 297–318. <https://doi.org/10.1017/CBO9780511564512.011>.
- Reichelt, K., 2005. Late Aptian-Albian of the Vocontian Basin (SE-France) and Albian of NE-Texas: Biostratigraphic and Paleoenvironmental Implications by Planktic Foraminifera Faunas. PhD Thesis. Geowissenschaftlichen Fakultät, Eberhard-Karls-Universität Tübingen.
- Remírez, M.N., Algeo, T.J., 2020. Carbon-cycle changes during the Toarcian (Early Jurassic) and implications for regional versus global drivers of the Toarcian oceanic anoxic event. *Earth Sci. Rev.* 209, 103283. <https://doi.org/10.1016/j.earscirev.2020.103283>.
- Reolid, M., Rodríguez-Tovar, F.J., Marok, A., Sebane, A., 2012. The Toarcian oceanic anoxic event in the Western Saharan Atlas, Algeria (North African paleomargin): role of anoxia and productivity. *Bull. Geol. Soc. Am.* <https://doi.org/10.1130/B30585.1>.
- Reolid, M., Emanuela, M., Nieto, L.M., Rodríguez-Tovar, F.J., 2014. The Early Toarcian Oceanic Anoxic Event in the External Subbetic (Southiberian Palaeomargin, Westernmost Tethys): Geochemistry, nannofossils and ichnology. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 411, 79–94. <https://doi.org/10.1016/j.palaeo.2014.06.023>.
- Reolid, M., Iwanczuk, J., Mattioli, E., Abad, I., 2020. Integration of gamma ray spectrometry, magnetic susceptibility and calcareous nannofossils for interpreting environmental perturbations: an example from the Jenkyns Event (lower Toarcian) from South Iberian Palaeomargin (Median Subbetic, SE Spain). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 560, 110031. <https://doi.org/10.1016/j.palaeo.2020.110031>.
- Riding, J.B., Leng, M.J., Kender, S., Hesselbo, S.P., Feist-Burkhardt, S., 2013. Isotopic and palynological evidence for a new Early Jurassic environmental perturbation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 374, 16–27. <https://doi.org/10.1016/j.palaeo.2012.10.019>.
- Rodger Harvey, H., Tuttle, J.H., Tyler Bell, J., 1995. Kinetics of phytoplankton decay during simulated sedimentation: changes in biochemical composition and microbial activity under oxic and anoxic conditions. *Geochim. Cosmochim. Acta* 59, 3367–3377. [https://doi.org/10.1016/0016-7037\(95\)00217-N](https://doi.org/10.1016/0016-7037(95)00217-N).
- Rodrigues, B., Duarte, L.V., Mendonça Filho, J.G., Santos, L.G., Donizeti de Oliveira, A., 2016. Evidence of terrestrial organic matter deposition across the early Toarcian recorded in the northern Lusitanian Basin, Portugal. *Int. J. Coal Geol.* 168, 35–45. <https://doi.org/10.1016/j.coal.2016.06.016>.
- Rodrigues, B., Silva, R.L., Reolid, M., Mendonça Filho, J.G., Duarte, L.V., 2019. Sedimentary organic matter and $\delta^{13}\text{C}$ Kerogen variation on the southern Iberian palaeomargin (Betic Cordillera, SE Spain) during the latest Pliensbachian–Early Toarcian. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 534, 109342. <https://doi.org/10.1016/j.palaeo.2019.109342>.
- Rodrigues, B., Duarte, L.V., Silva, R.L., Mendonça Filho, J.G., 2020a. Sedimentary organic matter and early Toarcian environmental changes in the Lusitanian Basin (Portugal). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 554, 109781. <https://doi.org/10.1016/j.palaeo.2020.109781>.
- Rodrigues, B., Silva, R.L., Filho, J.G.M., Sadki, D., Mendonça, J.O., Duarte, L.V., 2020b. Late Pliensbachian–Early Toarcian palaeoenvironmental dynamics and the Pliensbachian–Toarcian Event in the Middle Atlas Basin (Morocco). *Int. J. Coal Geol.* 217, 103339. <https://doi.org/10.1016/j.coal.2019.103339>.
- Rodrigues, B., Silva, R.L., Mendonça Filho, J.G., Comas-Rengifo, M.J., Goy, A., Duarte, L.V., 2020c. Kerogen assemblages and $\delta^{13}\text{C}$ Kerogen of the uppermost Pliensbachian–lower Toarcian succession of the Asturian Basin (northern Spain). *Int. J. Coal Geol.* 229, 103573. <https://doi.org/10.1016/j.coal.2020.103573>.
- Rodrigues, B., Silva, R.L., Mendonça Filho, J.G., Reolid, M., Sadki, D., Comas-Rengifo, M.J., Goy, A., Duarte, L.V., 2021. The Phytoclast Group as a tracer of palaeoenvironmental changes in the early Toarcian. *Geol. Soc. London, Spec. Publ.* <https://doi.org/10.1144/SP514-2020-271>. SP514-2020-271.
- Rodríguez-Tovar, F.J., Pardo-Igúzquiza, E., Reolid, M., Bartolini, A., 2016. Spectral analysis of Toarcian sediments from the Valdorbia section (Umbria-Marche Apennines): the astronomical input in the foraminiferal record. *Riv. Ital. Paleontol. Stratigr.* 122, 45–59. <https://doi.org/10.13130/2039-4942/7237>.
- Rodríguez-Tovar, F.J., Miguez-Salas, O., Duarte, L.V., 2017. Toarcian Oceanic Anoxic Event induced unusual behaviour and palaeobiological changes in Thalassinoides tracemakers. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 485, 46–56. <https://doi.org/10.1016/j.palaeo.2017.06.002>.
- Röhl, H., Schmid-Röhl, A., 2005. Lower Toarcian (Upper Liassic) Black Shales of the Central European Epicontinental Basin: a sequence stratigraphic case study from the Sw German Posidonia Shale. In: Harris, N.B. (Ed.), *The Deposition of Organic-Carbon-Rich Sediments: Models, Mechanisms, and Consequences*. Society for Sedimentary Geology, Houston, pp. 165–189. <https://doi.org/10.2110/pec.05.82.0165>.
- Röhl, H.-J., Schmid-Röhl, A., Oschmann, W., Frimmel, A., Schwark, L., 2001. The Posidonia Shale (Lower Toarcian) of SW-Germany: an oxygen-depleted ecosystem controlled by sea level and palaeoclimate. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 169, 273–299. [https://doi.org/10.1016/S0031-0182\(00\)00152-8](https://doi.org/10.1016/S0031-0182(00)00152-8).
- Ronov, A.B., 1958. Organic carbon in sedimentary rocks (in relation to the presence of petroleum). *Geochemistry* 5, 510–536.
- Rosales, I., Quesada, S., Robles, S., 2004. Paleotemperature variations of Early Jurassic seawater recorded in geochemical trends of belemnites from the Basque-Cantabrian basin, northern Spain. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 203, 253–275. [https://doi.org/10.1016/S0031-0182\(03\)00686-2](https://doi.org/10.1016/S0031-0182(03)00686-2).
- Rosales, I., Quesada, S., Robles, S., 2006. Geochemical arguments for identifying second-order sea-level changes in hemipelagic carbonate ramp deposits. *Terra Nova* 18, 233–240. <https://doi.org/10.1111/j.1365-3121.2006.00684.x>.
- Ruebsam, W., Al-Husseini, M., 2020. Calibrating the Early Toarcian (Early Jurassic) with stratigraphic black holes (SBH). *Gondwana Res.* 82, 317–336. <https://doi.org/10.1016/j.gr.2020.01.011>.
- Ruebsam, W., Müller, T., Kovács, J., Pálffy, J., Schwark, L., 2018. Environmental response to the early Toarcian carbon cycle and climate perturbations in the northeastern part of the West Tethys shelf. *Gondwana Res.* 59, 144–158. <https://doi.org/10.1016/j.gr.2018.03.013>.
- Ruebsam, W., Mayer, B., Schwark, L., 2019. Cryosphere carbon dynamics control early Toarcian global warming and sea level evolution. *Glob. Planet. Change* 172, 440–453. <https://doi.org/10.1016/j.gloplacha.2018.11.003>.
- Ruebsam, W., Reolid, M., Schwark, L., 2020. $\delta^{13}\text{C}$ of terrestrial vegetation records Toarcian CO_2 and climate gradients. *Sci. Rep.* 10, 117. <https://doi.org/10.1038/s41598-019-56710-6>.
- Ruhl, M., Deenen, M.H.L., Abels, H.A., Bonis, N.R., Krijgsman, W., Kürschner, W.M., 2010. Astronomical constraints on the duration of the early Jurassic Hettangian stage and recovery rates following the end-Triassic mass extinction (St Audrie’s Bay/East Quantoxhead, UK). *Earth Planet. Sci. Lett.* 295, 262–276. <https://doi.org/10.1016/j.epsl.2010.04.008>.
- Ruhl, M., Hesselbo, S.P., Hinnov, L., Jenkyns, H.C., Xu, W., Riding, J.B., Storm, M., Minisini, D., Ullmann, C.V., Leng, M.J., 2016. Astronomical constraints on the duration of the Early Jurassic Pliensbachian Stage and global climatic fluctuations. *Earth Planet. Sci. Lett.* 455, 149–165. <https://doi.org/10.1016/j.epsl.2016.08.038>.
- Ruvalcaba Baroni, I., Pohl, A., van Helmond, N.A.G.M., Papadomanolaki, N.M., Coe, A.L., Cohen, A.S., van de Schootbrugge, B., Donnadiu, Y., Slomp, C.P., 2018. Ocean circulation in the Toarcian (Early Jurassic): a key control on deoxygenation and carbon burial on the European Shelf. *Paleoceanogr. Paleoclimatol.* 33, 994–1012. <https://doi.org/10.1029/2018PA003394>.
- Sabatino, N., Neri, R., Bellanca, A., Jenkyns, H.C., Baudin, F., Parisi, G., Masetti, D., 2009. Carbon-isotope records of the Early Jurassic (Toarcian) oceanic anoxic event from the Valdorbia (Umbria-Marche Apennines) and Monte Mangart (Julian Alps) sections: palaeoenvironmental and stratigraphic implications. *Sedimentology* 56, 1307–1328. <https://doi.org/10.1111/j.1365-3091.2008.01035.x>.
- Sabatino, N., Vlahovic, I., Jenkyns, H.C., Scopelliti, G., Neri, R., Prtoljan, B., Velic, I., 2013. Carbon-isotope record and palaeoenvironmental changes during the early

- toarcian oceanic anoxic event in shallow-marine carbonates of the adriatic carbonate platform in Croatia. *Geol. Mag.* 150, 1085–1102. <https://doi.org/10.1017/S0016756813000083>.
- Sachse, V.F., Leythaeuser, D., Grobe, A., Rachidi, M., Littke, R., 2012. Organic geochemistry and petrology of a lower Jurassic (Pliensbachian) petroleum source rock from Ait Moussa, Middle Atlas, Morocco. *J. Pet. Geol.* 35, 5–23. <https://doi.org/10.1111/j.1747-5457.2012.00516.x>.
- Sadki, D., Marok, A., Soussi, M., 2014. The Lower and Middle Jurassic of the Western Mediterranean Tethys (Morocco, Algeria, and Tunisia). In: Rocha, R., Pais, J., Kullberg, J.C., Finney, S. (Eds.), *STRATI 2013*. Springer International Publishing, Cham, pp. 699–703. https://doi.org/10.1007/978-3-319-04364-7_132.
- Salem, N.-E., 2013. *Geochemical Characterisation of the Pliensbachian-Toarcian Boundary During the Onset of the Toarcian Oceanic Anoxic Event*. Newcastle University, North Yorkshire, UK. PhD thesis.
- Schlanger, S., Jenkyns, H., 1976. Cretaceous oceanic anoxic events: causes and consequences. *Geol. Mijnb.* 55, 179–184.
- Schmid-Röhl, A., Röhl, H.-J., Oschmann, W., Frimmel, A., Schwark, L., 2002. Palaeoenvironmental reconstruction of Lower Toarcian epicontinental black shales (Posidonia Shale, SW Germany): global versus regional control. *Geobios* 35, 13–20. [https://doi.org/10.1016/S0016-6995\(02\)00005-0](https://doi.org/10.1016/S0016-6995(02)00005-0).
- Schöllhorn, I., Adatte, T., Charbonnier, G., Mattioli, E., Spangenberg, J.E., Föllmi, K.B., 2020a. Pliensbachian environmental perturbations and their potential link with volcanic activity: Swiss and British geochemical records. *Sediment. Geol.* 406, 105665. <https://doi.org/10.1016/j.sedgeo.2020.105665>.
- Schöllhorn, I., Adatte, T., Van de Schootbrugge, B., Houben, A., Charbonnier, G., Janssen, N., Föllmi, K.B., 2020b. Climate and environmental response to the break-up of Pangea during the Early Jurassic (Hettangian-Pliensbachian); the Dorset coast (UK) revisited. *Glob. Planet. Change* 185, 103096. <https://doi.org/10.1016/j.gloplacha.2019.103096>.
- Schouten, S., de Graaf, W., Sinninghe Damsté, J.S., van Driel, G.B., de Leeuw, J.W., 1994. Laboratory simulation of natural sulphurization: II. Reaction of multi-functionalized lipids with inorganic polysulphides at low temperatures. *Org. Geochem.* 22, 825–IN14. [https://doi.org/10.1016/0146-6380\(94\)90142-2](https://doi.org/10.1016/0146-6380(94)90142-2).
- Scotchman, I.C., 2001. Petroleum geochemistry of the Lower and Middle Jurassic in Atlantic margin basins of Ireland and the UK. In: Shannon, E.M., Haughton, P.D.W., Corcoran, D.V. (Eds.), *The Petroleum Exploration of Ireland's Offshore Basins*, Special Publications 188. The Geological Society, London, pp. 31–60. <https://doi.org/10.1144/GSL.SP.2001.188.01.03>.
- Shurygin, B.N., Nikitenko, B.L., Meledina, S.V., Dzyuba, O.S., Knyazev, V.G., 2011. Comprehensive zonal subdivisions of Siberian Jurassic and their significance for Circum-Arctic correlations. *Russ. Geol. Geophys.* 52, 825–844. <https://doi.org/10.1016/j.rgg.2011.07.007>.
- Silva, R.L., 2015. *Jurassic Organic-rich Carbonate Series of the Lusitanian Basin (Portugal): Sedimentology, Geochemistry and Palaeoenvironmental Interpretation*. PhD thesis. University of Coimbra.
- Silva, R.L., Duarte, L.V., 2015. Organic matter production and preservation in the Lusitanian Basin (Portugal) and Pliensbachian climatic hot snaps. *Glob. Planet. Change* 131, 24–34. <https://doi.org/10.1016/j.gloplacha.2015.05.002>.
- Silva, R.L., Duarte, L.V., Comas-Rengifo, M.J., Mendonça Filho, J.G., Azerêdo, A.C., 2011. Update of the carbon and oxygen isotopic records of the Early-Late Pliensbachian (Early Jurassic, ~187Ma): insights from the organic-rich hemipelagic series of the Lusitanian Basin (Portugal). *Chem. Geol.* 283, 177–184. <https://doi.org/10.1016/j.chemgeo.2011.01.010>.
- Silva, R.L., Mendonça Filho, J.G., da Silva, F.S., Duarte, L.V., Silva, T.F., Ferreira, R., Azerêdo, A.C., 2012. Can biogeochemistry aid in the palaeoenvironmental/early diagenesis reconstruction of the ~187 Ma (Pliensbachian) organic-rich hemipelagic series of the Lusitanian Basin (Portugal)? *Bull. Geosci.* 87, 373–382. <https://doi.org/10.3140/bull.geosci.1315>.
- Silva, R.L., Duarte, L.V., Mendonça Filho, J.G., 2013. Optical and geochemical characterization of Lower Sinemurian (Lower Jurassic) fossil wood from the Lusitanian Basin (Portugal). *Geochem. J.* 47, 489–498. <https://doi.org/10.2343/geochemj.2.0270>.
- Silva, R.L., Duarte, L.V., Comas-Rengifo, M.J., 2015. Facies and carbon isotope chemostratigraphy of Lower Jurassic carbonate deposits, Lusitanian Basin (Portugal). In: Ramkumar, M. (Ed.), *Chemostratigraphy. Concepts, Techniques, and Applications*. Elsevier, Amsterdam, pp. 341–371. <https://doi.org/10.1016/B978-0-12-419968-2.00013-3>.
- Silva, R.L., Carlisle, C.A.M., Wach, G., 2017. A new TOC, Rock-Eval, and carbon isotope record of Lower Jurassic source rocks from the Slyne Basin, offshore Ireland. *Mar. Pet. Geol.* 86, 499–511. <https://doi.org/10.1016/j.marpetgeo.2017.06.004>.
- Silva, R.L., Duarte, L.V., Wach, G.D., Morrison, N., Campbell, T., 2020a. Oceanic organic carbon as a possible first-order control on the carbon cycle during the Bathonian–Callovian. *Glob. Planet. Change* 184, 103058. <https://doi.org/10.1016/j.gloplacha.2019.103058>.
- Silva, R.L., Wach, G.D., Hesselbo, S.P., O'Connor, D.E., 2020b. Total organic carbon and pyrolysis analysis of the Lower Cretaceous in Compton Bay and Atherfield, Isle of Wight (England). *Proc. Geol. Assoc.* 131, 51–59. <https://doi.org/10.1016/j.pgeola.2019.11.005>.
- Silva, R.L., Ruhl, M., Barry, C., Reolid, M., Ruebsam, W., 2021. Pacing of late Pliensbachian and early Toarcian carbon cycle perturbations and environmental change in the westernmost Tethys (La Cerradura section, Subbetic Zone of the Betic Cordillera, Spain). *Geol. Soc. London, Spec. Publ.* <https://doi.org/10.1144/SP514-2021-27>. SP514-2021–27.
- Simms, M.J.J., 2004. The Wessex Basin (Dorset and central Somerset). In: Simms, M.J.J., Chidlaw, N., Morton, N., Page, K.N. (Eds.), *British Lower Jurassic Stratigraphy*. Geological Conservation Review Series, Peterborough, pp. 53–109.
- Simms, M.J.J., Chidlaw, N., Morton, N., Page, K.N., 2004. The Cleveland Basin. In: Simms, M.J., Chidlaw, N., Morton, N., Page, K.N. (Eds.), *British Lower Jurassic Stratigraphy*. Geological Conservation Review Series, Peterborough, pp. 237–304.
- Sinninghe Damsté, J.S., Irene, W., Rijpstra, C., de Leeuw, J.W., Schenck, P.A., 1988. Origin of organic sulphur compounds and sulphur-containing high molecular weight substances in sediments and immature crude oils. *Org. Geochem.* 13, 593–606. [https://doi.org/10.1016/0146-6380\(88\)90079-4](https://doi.org/10.1016/0146-6380(88)90079-4).
- Sinninghe Damsté, J.S., Van Duin, A.C.T., Hollander, D., Kohnen, M.E.L., De Leeuw, J. W., 1995. Early diagenesis of bacteriohopanepolyol derivatives: formation of fossil homohopanoids. *Geochim. Cosmochim. Acta* 59, 5141–5157. [https://doi.org/10.1016/0016-7037\(95\)00338-X](https://doi.org/10.1016/0016-7037(95)00338-X).
- Slater, S.M., Twitchett, R.J., Danise, S., Vajda, V., 2019. Substantial vegetation response to Early Jurassic global warming with impacts on oceanic anoxia. *Nat. Geosci.* <https://doi.org/10.1038/s41561-019-0349-z>.
- Song, J., Littke, R., Maquil, R., Weniger, P., 2014. Organic facies variability in the Posidonia Black Shale from Luxembourg: Implications for thermal maturation and depositional environment. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 410, 316–336. <https://doi.org/10.1016/j.palaeo.2014.06.002>.
- Song, J., Littke, R., Weniger, P., Ostertag-Henning, C., Nelskamp, S., 2015. Shale oil potential and thermal maturity of the Lower Toarcian Posidonia Shale in NW Europe. *Int. J. Coal Geol.* 150–151, 127–153. <https://doi.org/10.1016/j.coal.2015.08.011>.
- Soua, M., 2014. A review of Jurassic Oceanic Anoxic Events as recorded in the northern margin of Africa, Tunisia. *J. Geosci. Geomat.* 2, 94–106. <https://doi.org/10.12691/jgg-2-3-4>.
- Soussi, M., 2003. Nouvelle nomenclature lithostratigraphique « événementielle » pour le Jurassique de la Tunisie atlantique. *Geobios* 36, 761–773. <https://doi.org/10.1016/j.geobios.2003.03.001>.
- Stapel, G., Cloetingh, S., Pronk, B., 1996. Quantitative subsidence analysis of the Mesozoic evolution of the Lusitanian basin (western Iberian margin). *Tectonophysics* 266, 493–507. [https://doi.org/10.1016/S0040-1951\(96\)00203-X](https://doi.org/10.1016/S0040-1951(96)00203-X).
- Storm, M.S., Hesselbo, S.P., Jenkyns, H.C., Ruhl, M., Ullmann, C.V., Xu, W., Leng, M.J., Riding, J.B., Gorbanenko, O., 2020. Orbital pacing and secular evolution of the Early Jurassic carbon cycle. *Proc. Natl. Acad. Sci.* 117, 3974–3982. <https://doi.org/10.1073/pnas.1912094117>.
- Suan, G., Pittet, B., Bour, I., Mattioli, E., Duarte, L.V., Mailliot, S., 2008. Duration of the Early Toarcian carbon isotope excursion deduced from spectral analysis: Consequence for its possible causes. *Earth Planet. Sci. Lett.* 267, 666–679. <https://doi.org/10.1016/j.epsl.2007.12.017>.
- Suan, G., Mattioli, E., Pittet, B., Lécuyer, C., Suchéras-Marx, B., Duarte, L.V., Philippe, M., Reggiani, L., Martineau, F., 2010. Secular environmental precursors to Early Toarcian (Jurassic) extreme climate changes. *Earth Planet. Sci. Lett.* 290, 448–458. <https://doi.org/10.1016/j.epsl.2009.12.047>.
- Suan, G., Nikitenko, B.L., Rogov, M.A., Baudin, F., Spangenberg, J.E., Knyazev, V.G., Glinskikh, L.A., Goryacheva, A.A., Adatte, T., Riding, J.B., Föllmi, K.B., Pittet, B., Mattioli, E., Lécuyer, C., 2011. Polar record of Early Jurassic massive carbon injection. *Earth Planet. Sci. Lett.* 312, 102–113. <https://doi.org/10.1016/j.epsl.2011.09.050>.
- Suan, G., van de Schootbrugge, B., Adatte, T., Fiebig, J., Oschmann, W., 2015. Calibrating the magnitude of the Toarcian carbon cycle perturbation. *Paleoceanography* 30, 495–509. <https://doi.org/10.1002/2014PA002758>.
- Suárez-Ruiz, I., Flores, D., Mendonça Filho, J.G., Hackley, P.C., 2012. Review and update of the applications of organic petrology: Part 1, geological applications. *Int. J. Coal Geol.* 99, 54–112. <https://doi.org/10.1016/j.coal.2012.02.004>.
- Sun, M.-Y., Aller, R.C., Lee, C., Wakeham, S.G., 2002. Effects of oxygen and redox oscillation on degradation of cell-associated lipids in surficial marine sediments. *Geochim. Cosmochim. Acta* 66, 2003–2012. [https://doi.org/10.1016/S0016-7037\(02\)00830-X](https://doi.org/10.1016/S0016-7037(02)00830-X).
- Svensen, H., Planke, S., Chevillier, L., Malthé-Sørensen, A., Corfú, F., Jamtveit, B., 2007. Hydrothermal venting of greenhouse gases triggering Early Jurassic global warming. *Earth Planet. Sci. Lett.* 256, 554–566. <https://doi.org/10.1016/j.epsl.2007.02.013>.
- Swanson, V.E., 1961. *Geology and geochemistry of uranium in marine black shales: a review*. In: *USGS Prof. Pap.* 365 (50 pp.).
- Swart, P.K., Eberli, G., 2005. The nature of the $\delta^{13}\text{C}$ of periplatform sediments: implications for stratigraphy and the global carbon cycle. *Sediment. Geol.* 175, 115–129. <https://doi.org/10.1016/j.sedgeo.2004.12.029>.
- Tegelaar, E., de Leeuw, J., Derenne, S., Largeau, C., 1989. A reappraisal of kerogen formation. *Geochim. Cosmochim. Acta* 53, 3103–3106. [https://doi.org/10.1016/0016-7037\(89\)90191-9](https://doi.org/10.1016/0016-7037(89)90191-9).
- Them, T.R., Gill, B.C., Caruthers, A.H., Gröcke, D.R., Tulsy, E.T., Martindale, R.C., Poulton, T.P., Smith, P.L., 2017. High-resolution carbon isotope records of the Toarcian Oceanic Anoxic Event (Early Jurassic) from North America and implications for the global drivers of the Toarcian carbon cycle. *Earth Planet. Sci. Lett.* 459, 118–126. <https://doi.org/10.1016/j.epsl.2016.11.021>.
- Thibault, N., Ruhl, M., Ullmann, C.V., Korte, C., Kemp, D.B., Gröcke, D.R., Hesselbo, S.P., 2018. The wider context of the Lower Jurassic Toarcian oceanic anoxic event in Yorkshire coastal outcrops, UK. *Proc. Geol. Assoc.* 129, 372–391. <https://doi.org/10.1016/j.pgeola.2017.10.007>.
- Thrasher, J., 1992. Thermal effect of the Tertiary Cuillins Intrusive Complex in the Jurassic of the Hebrides: an organic geochemical study. *Geol. Soc. Lond. Spec. Publ.* 62, 35–49. <https://doi.org/10.1144/GSL.SP.1992.062.01.06>.
- Tissot, B.P., Welte, D.H., 1978. *Petroleum Formation and Occurrence: A New Approach to Oil and Gas Exploration*. Springer-Verlag.
- Tissot, B.P., Welte, D.H., 1984. *Petroleum Formation and Occurrence*. Springer, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-642-87813-8>.

- Trabucho-Alexandre, J., Van Gilst, R.I., Rodríguez-López, J.P., De Boer, P.L., 2011. The sedimentary expression of oceanic anoxic event 1b in the North Atlantic. *Sedimentology* 58, 1217–1246. <https://doi.org/10.1111/j.1365-3091.2010.01202.x>.
- Trueblood, S., 1992. Petroleum geology of the Slyne Trough and adjacent basins. In: Parnell, J. (Ed.), *Basins on the Atlantic Seaboard: Petroleum Geology, Sedimentology and Basin Evolution*, Special Publications 62. Geological Society, London, pp. 315–326. <https://doi.org/10.1144/GSL.SP.1992.062.01.24>.
- Tyson, R.V., 1995. *Sedimentary Organic Matter: Organic Facies and Palynofacies*. Springer Netherlands.
- Ullmann, C.V., Thibault, N., Ruhl, M., Hesselbo, S.P., Korte, C., 2014. Effect of a Jurassic oceanic anoxic event on belemnite ecology and evolution. *Proc. Natl. Acad. Sci.* 111, 10073–10076. <https://doi.org/10.1073/pnas.1320156111>.
- Ullmann, C.V., Boyle, R., Duarte, L.V., Hesselbo, S.P., Kasemann, S.A., Klein, T., Lenton, T.M., Piazza, V., Aberhan, M., 2020. Warm afterglow from the Toarcian Oceanic Anoxic Event drives the success of deep-adapted brachiopods. *Sci. Rep.* 10, 6549. <https://doi.org/10.1038/s41598-020-63487-6>.
- van Acken, D., Tütken, T., Daly, J.S., Schmid-Röhl, A., Orr, P.J., 2019. Rhenium-osmium geochronology of the Toarcian Posidonia Shale, SW Germany. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 534, 109294. <https://doi.org/10.1016/j.palaeo.2019.109294>.
- van Breugel, Y., Baas, M., Schouten, S., Mattioli, E., Sinnighe Damsté, J.S., 2006. Isorenieratane record in black shales from the Paris Basin, France: constraints on recycling of respired CO₂ as a mechanism for negative carbon isotope shifts during the Toarcian oceanic anoxic event. *Paleoceanography* 21, 1–8. <https://doi.org/10.1029/2006PA001305>.
- van Buchem, F.S.P., de Boer, P.L., McCave, I.N., Herbin, J.-P., van Buchem, F.S.P., de Boer, P.L., McCave, I.N., Herbin, J.-P., 1995. The organic carbon distribution in Mesozoic Marine sediments and the influence of orbital climatic cycles (England and the Western North Atlantic). In: Huc, A.-Y. (Ed.), *Paleogeography, Paleoclimate, and Source Rock*. The American Association of Petroleum Geologists, Tulsa, Oklahoma, U.S.A., pp. 303–335. <https://doi.org/10.1306/St40595C13>.
- van de Schootbrugge, B., Bailey, T.R., Rosenthal, Y., Katz, M.E., Wright, J.D., Miller, K.G., Feist-Burkhardt, S., Falkowski, P.G., 2005a. Early Jurassic climate change and the radiation of organic-walled phytoplankton in the Tethys Ocean. *Paleobiology* 31, 73–97. [https://doi.org/10.1666/0094-8373\(2005\)031<0073:EJCCAT>2.0.CO;2](https://doi.org/10.1666/0094-8373(2005)031<0073:EJCCAT>2.0.CO;2).
- van de Schootbrugge, B., McArthur, J.M., Bailey, T.R., Rosenthal, Y., Wright, J.D., Miller, K.G., 2005b. Toarcian oceanic anoxic event: an assessment of global causes using belemnite C isotope records. *Paleoceanography* 20, PA3008. <https://doi.org/10.1029/2004PA001102>.
- van de Schootbrugge, B., Bachan, A., Suan, G., Richoz, S., Payne, J.L., 2013. Microbes, mud and methane: Cause and consequence of recurrent early Jurassic anoxia following the end-Triassic mass extinction. *Palaeontology* 56, 685–709. <https://doi.org/10.1111/pala.12034>.
- Vandenbroucke, M., Largeau, C., 2007. Kerogen origin, evolution and structure. *Org. Geochem.* <https://doi.org/10.1016/j.orggeochem.2007.01.001>.
- Voigt, S., Gale, A.S., Jung, C., Jenkyns, H.C., 2012. Global correlation of Upper Campanian - Maastrichtian successions using carbon-isotope stratigraphy: development of a new Maastrichtian timescale. *Newslett. Stratigr.* 45, 25–53. <https://doi.org/10.1127/0078-0421/2012/0016>.
- Weedon, G.P., Jenkyns, H.C., 1990. Regular and irregular climatic cycles and the Belemnite Marls (Pliensbachian, Lower Jurassic, Wessex Basin). *J. Geol. Soc. Lond.* 147, 915–918. <https://doi.org/10.1144/gsjgs.147.6.0915>.
- Weedon, G.P., Jenkyns, H.C., 1999. Cyclostratigraphy and the Early Jurassic timescale: data from the Belemnite Marls, Dorset, southern England. *Bull. Geol. Soc. Am.* 111, 1823–1840. [https://doi.org/10.1130/0016-7606\(1999\)111<1823:CATEJT>2.3.CO;2](https://doi.org/10.1130/0016-7606(1999)111<1823:CATEJT>2.3.CO;2).
- Wefer, G., Berger, W.H., 1991. Isotope paleontology: growth and composition of extant calcareous species. *Mar. Geol.* 100, 207–248. [https://doi.org/10.1016/0025-3227\(91\)90234-U](https://doi.org/10.1016/0025-3227(91)90234-U).
- Weissert, H., 1981. The environment of deposition of black shales in the Early Cretaceous: an ongoing controversy. In: Warme, J.E., Douglas, R.G., Winterer, E.L. (Eds.), *The Deep Sea Drilling Project: A Decade of Progress*. Society for Sedimentary Geology, Tulsa, pp. 547–560. <https://doi.org/10.2110/pec.81.32.0547>.
- Weissert, H., 1989. C-Isotope stratigraphy, a monitor of paleoenvironmental change: a case study from the early cretaceous. *Surv. Geophys.* 10, 1–61. <https://doi.org/10.1007/BF01901664>.
- Weissert, H., Lini, A., Föllmi, K.B., Kuhn, O., 1998. Correlation of Early Cretaceous carbon isotope stratigraphy and platform drowning events: a possible link? *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 137, 189–203. [https://doi.org/10.1016/S0031-0182\(97\)00109-0](https://doi.org/10.1016/S0031-0182(97)00109-0).
- Westermann, S., Föllmi, K.B., Adatte, T., Matera, V., Schnyder, J., Fleitmann, D., Fiet, N., Ploch, I., Duchamp-Alphonse, S., 2010. The Valanginian δ¹³C excursion may not be an expression of a global oceanic anoxic event. *Earth Planet. Sci. Lett.* <https://doi.org/10.1016/j.epsl.2009.12.011>.
- Wignall, P.B., Newton, R.J., Little, C.T.S., 2005. The timing of paleoenvironmental changes and cause-and-effect relationships during the early Jurassic mass extinction in Europe. *Am. J. Sci.* 305, 1014–1032.
- Woodfine, R.G., Jenkyns, H.C., Sarti, M., Baroncini, F., Violante, C., 2008. The response of two Tethyan carbonate platforms to the early Toarcian (Jurassic) oceanic anoxic event: environmental change and differential subsidence. *Sedimentology* 55, 1011–1028. <https://doi.org/10.1111/j.1365-3091.2007.00934.x>.
- Xu, W., Ruhl, M., Hesselbo, S.P., Riding, J.B., Jenkyns, H.C., 2017a. Orbital pacing of the Early Jurassic carbon cycle, black-shale formation and seabed methane seepage. *Sedimentology* 64, 127–149. <https://doi.org/10.1111/sed.12329>.
- Xu, W., Ruhl, M., Jenkyns, H.C., Hesselbo, S.P., Riding, J.B., Selby, D., Naafs, B.D.A., Weijers, J.W.H., Pancost, R.D., Tegelaar, E.W., Idiz, E.F., 2017b. Carbon sequestration in an expanded lake system during the Toarcian oceanic anoxic event. *Nat. Geosci.* 10, 129–134. <https://doi.org/10.1038/ngeo2871>.
- Xu, W., Ruhl, M., Jenkyns, H.C., Leng, M.J., Huggett, J.M., Minisini, D., Ullmann, C.V., Riding, J.B., Weijers, J.W.H., Storm, M.S., Percival, L.M.E., Tosca, N.J., Idiz, E.F., Tegelaar, E.W., Hesselbo, S.P., 2018. Evolution of the Toarcian (Early Jurassic) carbon-cycle and global climatic controls on local sedimentary processes (Cardigan Bay Basin, UK). *Earth Planet. Sci. Lett.* 484, 396–411. <https://doi.org/10.1016/j.epsl.2017.12.037>.
- Zonneveld, K.A.F., Versteegh, G.J.M., Kasten, S., Eglinton, T.I., Emeis, K.-C., Huguet, C., Koch, B.P., de Lange, G.J., de Leeuw, J.W., Middelburg, J.J., Mollenhauer, G., Prah, F.G., Rethemeyer, J., Wakeham, S.G., 2010. Selective preservation of organic matter in marine environments; processes and impact on the sedimentary record. *Biogeosciences* 7, 483–511. <https://doi.org/10.5194/bg-7-483-2010>.