

**Formal proposal for the Global Boundary Stratotype Section and Point (GSSP)
of the Bathonian Stage, at the base of the Zigzag Zone in the Ravin du Bès Section
(Bas-Auran, Subalpine Basin, SE France)**

SIXTO R. FERNÁNDEZ-LÓPEZ, GIULIO PAVIA, ELISABETTA ERBA, MYETTE GUIOMAR,
MARIA H. HENRIQUES, ROBERTO LANZA, CHARLES MANGOLD, DAVIDE OLIVERO AND
DANIELE TIRABOSCHI

Fernández-López, S.R., Pavia, G., Erba, E., Guiomar, M., Henriques, M.H., Lanza, R., Mangold, C., Olivero, D. & Tiraboschi, D. 2007: *Formal proposal for the Global Boundary Stratotype Section and Point (GSSP) of the Bathonian Stage, at the base of the Zigzag Zone in the Ravin du Bès Section (Bas-Auran, Subalpine Basin, SE France)*. International Subcommission on Jurassic Stratigraphy, Bathonian Working Group Ballot 2007. 43 pp.

Sixto Rafael Fernández-López [sixto@geo.ucm.es], Departamento y UEI de Paleontología, Facultad de Ciencias Geológicas (UCM) e Instituto de Geología Económica (CSIC-UCM), 28040-Madrid (Spain);

Giulio Pavia [giulio.pavia@unito.it], Dipartimento di Scienze della Terra, via Valperga Caluso 35, 10125-Torino (Italy);

Elisabetta Erba [elisabetta.erba@unimi.it], Dipartimento di Scienze della Terra “Ardito Desio”, Università degli Studi di Milano, Via Mangiagalli, 34, 20133-Milano (Italy);

Mvette Guiomar [myette.guiomar@libertysurf.fr], Centre de géologie – Parc Saint-Benoît, 04000-Digne les bains (France);

Maria Helena Henriques [hhenriq@dct.uc.pt], Departamento Ciências da Terra e Centro de Geociências, Faculdade de Ciências e Tecnologia, Universidade de Coimbra, 3000-272 Coimbra (Portugal);

Roberto Lanza [roberto.lanza@unito.it], Dipartimento di Scienze della Terra, via Valperga Caluso 35, 10125-Torino (Italy);

Charles Mangold, Université Claude Bernard, Lyon-1, UFR des Sciences de la Terre et CNRS, UMR 5125, 27-43, bd du 11-Novembre-1918, 69622-Villeurbanne cedex (France);

David Olivero [David.Olivero@univ-lyon1.fr], Université Claude Bernard, Lyon-1, UFR des Sciences de la Terre et CNRS, UMR 5125, 27-43, bd du 11-Novembre-1918, 69622-Villeurbanne cedex (France);

Daniele Tiraboschi [daniele.tiraboschi@unimi.it], Dipartimento di Scienze della Terra “Ardito Desio”, Università degli Studi di Milano, Via Mangiagalli, 34, 20133-Milano (Italy).

Abstract

The Global Boundary Stratotype Section and Point (GSSP) for the base of the Bathonian Stage is proposed at the base of limestone bed RB071 (bed 23 in Sturani 1967) in the Ravin du Bès Section (43°57'38"N, 6°18'55"E), Bas-Auran area, in the Chaudon-Norante commune, around 25 km at the South-Southeast of Digne-les-Bains, in the "Alpes de Haute Provence" French department. The Ravin du Bès Section, as formal candidate GSSP for the base of the Bathonian Stage, satisfies most of the requirements recommended by the International Commission on Stratigraphy (ICS): 1) The exposure extends over 13 m in thickness, comprising more than five metres of fossiliferous levels below and above the boundary. The Bathonian basal bed corresponds to the "Marno-calcaires à *Cancellophycus*", is located 7.8 m below the "Terres Noires" Formation and forms part of a transgressive-facies cycle. At the Bajocian-Bathonian transition, no vertical (bio-, ichno- or tapho-) facies changes, stratigraphic gaps or hiatuses have been recorded. There is no evidence of taphonomic condensation (i.e. mixture of fossils of different age or different chronostratigraphic units). Structural complexity, synsedimentary and tectonic disturbances, or important alterations by metamorphism are not relevant constraints in the Bas-Auran area. 2) There is a well-preserved, abundant and diverse fossil record across the boundary interval, with key markers (ammonites and nannofossils) for worldwide correlation of the uppermost Bajocian and Lower Bathonian. The boundary can be characterized by both primary and secondary (auxiliary) biostratigraphic markers. The section appears to be suitable for biostratigraphic study of microfossils, such as foraminifera, but as yet there are no published studies. The base of Bathonian Stage and Zigzag Zone in Bas-Auran corresponds to the renewal of parkinsonids and the first occurrence level of *Gonolkites convergens* Buckman. This level coincides with the first occurrence of *Morphoceras parvum* Wetzel. Calcareous nannofossils, as secondary global marker, are present in all beds and allow characterizing the Bajocian-Bathonian transition. 3) Regional analyses of sequence stratigraphy and manganese chemostratigraphy are available. Spectral gamma-ray data corroborate an Early Bathonian deepening half-cycle of second order. No data are currently available for strontium isotope ($^{87}\text{Sr}/^{86}\text{Sr}$ ratio), oxygen isotope ($\delta^{18}\text{O}$) or carbon isotope ($\delta^{13}\text{C}$) chemostratigraphy. Bajocian and Bathonian deposits have been remagnetized with a steady normal polarity. Volcanogenic deposits suitable for direct radio-isotope dating are not known in the section. 4) The criteria of accessibility, conservation and protection are assured by the "Réserve Naturelle Géologique de Haute Provence", protected under national law as recognised by UNESCO. The Cabo Mondego Section (Portugal), which provides complementary data of the ammonite succession at the Sub-Mediterranean Parvum Subzone and its chronocorrelation with the Northwest European Convergens Subzone, is suggested as the Bathonian auxiliary section and point (ASP) within this GSSP proposal.

Key-words: Bathonian Working Group, chronostratigraphy, biostratigraphy, geochronology.

Introduction

The Bathonian Working Group was established in 1984, during the 1st International Symposium on Jurassic Stratigraphy in Erlangen, in order to improve the geologic data relative to the Bajocian/Bathonian boundary. The Bas-Auran section was first proposed as stratotype in a presentation to the Luxembourg II (1967) Colloquium by Torrens, but not published until 1974 (Morton 1974; Torrens 1974a, b, 1987, 2002). This section was formally proposed as a candidate of the basal boundary stratotype of the Bathonian Stage by Innocenti *et al.* (1990) during the 2nd International Symposium on Jurassic Stratigraphy in Lisbon (1987). Over the following 20 years, several meetings have been organized by the Bathonian Working Group in Digne, La Palud, Budapest, Lyon and Torino. In the Bas-Auran area, the sections of Ravin du Bès, Ravin d'Auran and Ravin des Robines have been remeasured and recollected for taphonomic, sedimentologic and palaeoichnological analysis during 2006 and 2007. Reports from the Bathonian Working Group have been published by Mangold (from 1985 to 1999) and Fernández-López (from 2003 to 2007, as mentioned in Fernández-Lopez 2007b).

The present report, proposing the GSSP for the Bathonian Stage at the base of the Zigzag Zone in the Ravin du Bès Section, has been developed by several specialists and members of the Bathonian Working Group. The proposal is going to be submitted for voting to all members of the Bathonian Working Group (S.R. Fernández-López, convenor) and, later, within the International Subcommission on Jurassic Stratigraphy. The following persons are members of the Bathonian Working Group: Alméras Y. (France), Bardhan S. (India), Bodergat A.M. (France), Callomon J.H. (UK), Cresta S. (Italy), Dietl G. (Germany), Enay R. (France), Fernández-López S.R. (Spain), Galász A. (Hungary), Hall R.L. (Canada), Henriques M.H. (Portugal), Hillebrandt A. von (Germany), Lanza R. (Italy), Mangold C. (France), Matyja B. (Poland), Meléndez G. (Spain), Mitta, V. (Russia), Mönnig, E. (Germany), Morton N. (France), Page K. (UK), Pandey D.K. (India), Pavia G. (Italy), Poulsen N. (Denmark), Poulton T.P. (Canada), Riccardi A.C. (Argentina), Rogov M.A. (Russia), Sandoval J. (Spain), Schlögl J. (Slovak Republic), Schweigert G. (Germany), Seyed-Emami K. (Iran), Wierzbowski A. (Poland), Yin J.-R. (China). This report summarizes relevant results published by specialists, in order to achieve the formal ballot on the selection and proposal of a GSSP for the Bathonian Stage.

Definition of the base of the Bathonian Stage (S.R. Fernández-López)

The Bathonian is the third stage of the Middle Jurassic Series, above the Bajocian and below the Callovian. The name was introduced by d'Halloy (1843) and used as a stage by d'Orbigny (1850, pp. 607–608; 1852, pp. 491–492), derived from the “Bath Oolite”, in the vicinity of the city of Bath in Somerset (England). *Zigzagoceras zigzag* (d'Orbigny 1846, p. 390, pl. 129, figs. 9–10; Arkell 1958, p. 177, text-fig. 60, 1–3) and *Gonolkites convergens* Buckman (1925, pl. 546 A-B; Arkell 1956, pl. 18, fig. 8; pl. 19, figs. 1–2) are the index species, respectively, of the Bathonian basal zone and subzone. The Zigzag Zone was distinguished from the underlying Parkinsoni Zone by Oppel (1857, p. 579), and later assigned to the “Bath-Gruppe” (Oppel 1865, p. 309) in a discussion of the section at “Montagne de Crussol” in the Ardèche (France). The Convergens Subzone was mentioned by Maubeuge (1950, p. 4), based on the “Convergens horizon” that was used in letters by Arkell (1951–59, p.10; 1956, p. 62). The Bajocian/Bathonian boundary established between the Parkinsoni and Zigzag zones was recommended at the two congresses denominated “Colloque du Jurassique” and held in Luxembourg (Rioult 1964; Torrens 1965, 1974a, b). The localities of Bath (England) and the “Montagne de Crussol” (France), however, have been considered quite unsuitable for a typological definition of the Bathonian Stage, because they are condensed sections with discontinuous and lenticular beds (Torrens 1974a, b, 2002; Page 1996b). After the publication of Sturani (1967), the base of bed 23 of the Bas-Auran section, in which *Gonolkites convergens* Buckman, *Parkinsonia pachypleura* Buckman and *Morphoceras parvum* Wetzel first appear, was designated as the type by which to define the base for the Convergens Subzone of the Zigzag Zone and the base of the Bathonian Stage by several authors (Morton 1974; Torrens 1974, 1987; Harland *et al.* 1982). Later, there was general agreement among Bathonian specialists that the Bathonian Stage should start with the Standard Zigzag Zone, whose base is defined by the Convergens Subzone (Horizon 1 of Mangold 1984) followed by the Macrescens Subzone (Sturani 1967). The Parvum Subzone has been proposed by Mangold (1990) to denote the first Bathonian subzone of the Zigzag Zone in the Sub-Mediterranean Province, equivalent to the Convergens Subzone of the Northwest European Province and below the Macrescens Subzone. Analogously, due to palaeobiogeographical changes, the Dimorphitiformis Subzone has been proposed by Sandoval (1983) as the basal Bathonian subzone of the Zigzag Zone in the Mediterranean Province. Therefore, the position of the Bathonian basal boundary became justified by the base of the Northwest European Convergens Subzone of the Zigzag Zone remarkably well recorded in the Bas-Auran area, a place where both Convergens and Parvum subzones can be recognized and denote the rock bodies of the same stratigraphic interval (Fernández-López *et al.* 2007, Pavia *et al.* 2008).

The Ravin du Bès Section (Bas-Auran area)

The Bas-Auran sections are located in southeastern France, in the “Alpes de Haute Provence” French department, in the Chaudon-Norante commune, around 25 km at the South-Southeast of

Digne-les-Bains (Fig. 1). Three sections have been selected in two ravines (Fig. 2). The first, the Ravin du Bès Section (RB) is located near the l'Amata farm (coordinates: 43°57'38"N, 6°18'55"E, altitude 730 m). The second, the Ravin d'Auran Section (RA) is located in front of the Bas-Auran farm (coordinates: 43°57'29"N, 6°19'00"E, altitude 790 m). The third one, the Ravin des Robines Section (RR), is just 400 metres south of the RA section, along the Robines ravine (coordinates: 43°57'09"N, 6°18'50"E, altitude 830 m). They are located on the Castellane sheet of the "Carte géologique détaillée de la France" at the 1:80000 scale (Goguel 1966) on the Digne sheet of the "Carte géologique de la France" at the 1:50000 scale (Graciansky *et al.* 1982) and on the topographic sheet, scale 1:25000, Barrême, n° IGN 3615.

These sections, which are free from significant disconformities, range from the Bomfordi Subzone (Parkinsoni Zone, Upper Bajocian) to the Tenuiplicatus Subzone (Aurigerus Zone, Lower Bathonian) and are over 13 m thick. Structural complexity, synsedimentary and tectonic disturbance, or important alterations by metamorphism, are not relevant constraints in the Bas-Auran area.




Fig. 1. Geographic location of the Bas-Auran area (France).




Fig. 2. Topographic sketch of the Bas-Auran area and location of the stratigraphic sections.

History of research on the Bathonian succession of the "Alpes de Haute Provence" and in particular of the Bas-Auran area (G. Pavia)

The area of Digne-Barrême was remarked by diverse authors as one of the most important in the world for establishing the ammonite zonal succession of the Bathonian Stage (Garnier 1872; Haug 1891, p. 80; Guillaume 1938; Arkell 1956, p.149). The Bas-Auran locality was firstly mentioned by Haug (1891) and later visited by the French Geological Society (Zurcher 1985). Sturani in 1967 published a detailed study of the Bajocian-Bathonian succession with a litho- and biostratigraphical log settled from the whole outcrops of the Bas-Auran area. A partial revision of Sturani's work was produced by Torrens (1987) mainly on the Tenuiplicatus Subzone at the uppermost part of the marly-calcareous succession. Contribution on the lowermost Bathonian beds was finally presented by Innocenti *et al.* (1990) that inserted in Sturani's log the new material derived from field work during ten years. More recent samplings, mainly concentrated on poorly documented and critical intervals, enlarged the Bas-Auran database from the Zigzag Zone, and furnished new and complementary results on the taphonomy of the ammonoid fossil-assemblages (Fernández-López 2007a), as well as on the taxonomy and phylogenetic arrangement of the Bathonian Bigotitinae and the origin of Zigzagiceratinae (Fernández-López *et al.* 2007). In a recently submitted paper (Pavia *et al.* 2008), the

following purposes have been achieved: 1) to describe the successive ammonoid assemblages of the uppermost Bajocian to lowermost Bathonian in the Bas-Auran area, 2) to refine the subzonal biostratigraphic subdivision of the marly-calcareous succession, 3) to characterize the ammonoid content at the very base of the Zigzag Zone, 4) to demonstrate the general continuity of the ammonoid succession and 5) to attest the suitability of one of those sections to be selected as the GSSP of the Bathonian Stage.

Geological setting of the marly-calcareous succession from the Bajocian to Bathonian in the Digne area (D. Olivero & G. Pavia)

The studied area is located in the French Subalpine Basin (FSB), corresponding to a gulf on the northwestern margin of the Tethyan Ocean (Fig. 3).




Fig. 3. The northwestern margin of the Tethyan Ocean, with the location of the French Subalpine Basin (modified from Ziegler 1999).




Fig. 4. The French Subalpine Basin, with the location of the Bas-Auran area (from Olivero 2003, modified).

The basin is bordered by the “Massif Central” on the West and the Alpine Chain on the East (Fig. 4). During Middle Jurassic time, the basin margin was characterized by a network of tilted blocks similar to the present margin of the Atlantic Ocean (Lemoine 1984, 1985). The maximum depth of the central part of the basin probably was about 700–800 metres (Ferry 1990). The region was a transitional area between the epicontinental sea of the Paris Basin and the deep Piedmont oceanic domain. The thrust boundaries figured in the simplified tectonic map (Fig. 5) correspond to the different limits of the various tilted blocks. The Bas-Auran area, located in the middle of one of such block, was thus on the continental slope of the French Subalpine Basin. The studied succession is a cyclic marl-limestone alternation. In most previous works and on the geological map of Digne (Graciansky *et al.* 1982, Olivero and Atrops 1996) it was described as the “Calcaires à *Cancellophycus*” Formation which, in the studied region, ranges from Aalenian to Bathonian and is covered by the “Terres Noires” Formation (Late Bathonian to Oxfordian). The “Calcaires à *Cancellophycus*” Formation should not be mistaken for the partially coeval “Calcaires à *Zoophycos* du Verdon” Formation, Lower Bathonian to Middle Callovian in age, proposed by Olivero and Atrops (1996) in the southernmost transitional area, between the Subalpine Basin and the Provence Platform.




Fig. 5. Simplified tectonic map of the Bas-Auran region (Olivero 2003, modified from Graciansky *et al.* 1982).

Palaeoichnology, taphonomy, sedimentology and sequence stratigraphy of the upper Bajocian to lower Bathonian of the Bas-Auran area (S.R. Fernández-López & D. Olivero)

In the Bas-Auran area, Lower Bathonian deposits comprise black or grey limestone beds alternating with marls usually known as “Marno-calcaires à *Cancellophycus*” (Graciansky *et al.* 1982; Olivero & Atrops 1996). Petrographically and in terms of biofacies, these deposits are relatively uniform mudstones to wackestones, with common ammonoids, scarce sponges and very scarce nautiloids, brachiopods, bivalves, belemnites, echinoids, crinoids and gastropods. As to microfossils, the overall sedimentary facies shows a calcisphere-mudstone texture; the marls contain foraminifers (*Lenticulina*, *Dentalina*), ostracods and molluscs (cephalopods, bivalves, gastropods) along with detrital minerals, quartz, muscovite and biotite (Corbin *et al.* 2000).

Palaeoichnological studies have been carried out by Olivero (1994, 2003). Bioturbation textures are common and bioturbation structures are scarce, indicating dominant softgrounds. *Zoophycos*, *Chondrites* and *Planolites* occur from the bed RB093 to bed RB001. Local concentrations of trace fossils of these ichnotaxa in bed RB039 suggest the development of a soft- to firmground in this stratigraphic level (Fig. 6). Bioturbation structures indicative of firmground (*Thalassinoides*, *Rhizocorallium*, *Zoophycos* and trace fossils related to large *Halimedes*) occur in a more calcareous layer just overlying the top of the bed RB003. Biogenic borings indicative of hardground (*Zapfella*) are common, associated with very scarce encrusting serpulids, on the top of bed RB001, indicating the exceptional development of a stratigraphic discontinuity at the top of the “Marno-calcaires à *Cancellophycus*” on the Bas-Auran area. Sedimentation appears irregular and condensed from bed RB093 towards the top of the Bathonian Zigzag Zone, compared with previous intervals where a more constant and expanded sedimentation is suggested. At the Bajocian-Bathonian transition, however, no stratigraphic gaps or hiatuses have been recorded.




Fig. 6. Ravin du Bès Section and detail of beds around the Bajocian/Bathonian boundary. Limestone bed 071 indicates the base of the Bathonian. Scale bar 1 m.

From a taphonomic point of view (Fig. 7), the occurrence of resedimented and reelaborated ammonoids implies that some form of current flow or winnowing affected the burial of concretionary internal moulds. Ammonoids show the following taphonomic characters at the Bajocian-Bathonian transition: 1) high values of stratigraphic persistence of ammonoid shells, 2) dominance of homogeneous concretionary internal moulds of phragmocones, completely filled with sediment, and 3) dominance of unflattened sedimentary moulds bearing no signs of rounding, bioerosion or dense encrusting by organisms (such as serpulids, bryozoans or oysters). These taphonomic features are indicative of low rate of sedimentation and low rate of accumulation of sediment, associated with sedimentary starving in deep environments (Fernández-López 2007a).

The bed-scale limestone-marl alternation is primary in origin, although accentuated by diagenetic redistribution of carbonate. Lithological differentiation between marly and limestone intervals resulted from alternating episodes of carbonate input and starvation. Both lithologies may contain evidence of sedimentary and taphonomic reworking, associated with scours, which reflect low rate of sedimentation and stratigraphic condensation. There is no evidence, however, of taphonomic condensation (i.e. mixture of fossils of different age or different chronostratigraphic units) in the ammonoid fossil-assemblages, except in level 002. Sedimentological data and sequence-stratigraphy interpretations of the Jurassic deposits in the French Subalpine Basin have been published by Graciansky *et al.* (1993, 1998a, b), Olivero & Atrops 1996, Olivero *et al.* (1997), Hardenbol *et al.* (1998) and Jacquin *et al.* (1998).

Palaeoichnological, taphonomic and sedimentological results confirm, therefore, the development of a deepening phase associated with sedimentary starvation, within 3rd and 2nd order cycles, in the Bas-Auran area, during the Early Bathonian. The maximum deepening of a 2nd-order transgressive/regressive facies cycle (T/R 7, Upper Aalenian–Upper Bathonian, in Graciansky *et al.* 1993, 1998) is at the end of the Early Bathonian, which corresponds to an extensional and deepening phase of the basin. The outcrop successions at Bas-Auran show no obvious signs of non-sequence or discontinuity across the Bajocian/Bathonian boundary interval.




Fig. 7. Ammonoid biostratigraphic data at the Bajocian/Bathonian boundary in the Ravin du Bès and Ravin d'Auran sections, indicating ammonoid preservation states, types of elementary cycles, and system tracts of 3rd and 2nd order cycles (Fernández-López 2007a).

Palaeontological records

The Bomfordi and Convergens subzones in the Bas-Auran area contain an ammonoid succession that displays a maximum value of biostratigraphic and biostratigraphic completeness. Additional macrofossil groups occur in the sections (e. g. sponges, bivalves, brachiopods and belemnites), although they are scarce and have not yet been studied in detail.

Ammonites (S.R. Fernández-López, C. Mangold & G. Pavia)

Biostratigraphic data on ammonoids of the Bas-Auran sections have been published by Sturani (1967), Pavia (1973, 1983a, b, 1984, 1994, 2000, 2007), Torrens (1987), Innocenti *et al.* (1990), Olivero *et al.* (1997) and Joly (2000). New and complementary results from the




Fig. 8. Lower Bathonian ammonites from Bas-Auran area. Specimens have been whitened with magnesium oxide prior to photography. Black asterisk marks the last septum of the phragmocone. Scale bar 1 cm.

1. *Bigotites mondegoensis* Fernández-López *et al.* [M], specimen PU111312, level BA14, Convergents Sbz.
2. *Protozigzagiceras cf. torrensi* (Sturani) [m], specimen PU111573, level RA033 (=BA12), Macrescens Sbz.
- 3ab. *Protozigzagiceras aff. torrensi* (Sturani) [M], specimen PU31694, level BA13, Convergents Sbz.
4. *Bigotites sturanii* Fernández-López *et al.* [M], specimen PU111253, level BA19, Convergents Sbz.
5. *Bigotites diniensis* Sturani [m], specimen PU111243, level BA20, Convergents Sbz.
- 6ab. *Bigotites sturanii* Fernández-López *et al.* [M], specimen PU111233, level BA20, Convergents Sbz.
- 7ab. *Morphoceras parvum* Wetzel [M], specimen PU111564, level BA17, Convergents Sbz.
- 8ab. *Gonolkites convergens* Buckman [M], specimen PU111067, level BA15, Convergents Sbz.

biochronostratigraphic analysis of ammonoid fossil-assemblages at the Bajocian/Bathonian boundary in Bas-Auran are in press (Fig. 8, Fernández-López *et al.* 2007, Pavia *et al.* 2008). In the French Subalpine Basin, the successive ammonoid fossil-assemblages are composed of Mediterranean and Northwest European representatives, associated with some Sub-Mediterranean ones. Upper Bajocian and Lower Bathonian Phylloceratina and Lytoceratina, which represent Mediterranean taxa, are relatively common (up to 25% at subzonal scale, Fig. 9). Northwest European taxa, such as Parkinsoniinae, may surpass 25% at subzonal scale. Lower Bathonian Bigotitinae, endemic and characteristic of the Sub-Mediterranean Province, reach 13%. This complex palaeobiogeographical pattern of the Upper Bajocian and Lower Bathonian ammonoid fossil-assemblages allows recognizing of diverse subzonal schemes and accurate chronocorrelation between the three main provinces of the West Tethyan Subrealm.

The base of Bathonian and Zigzag Zone corresponds to the renewal of parkinsonids and the first occurrence level of *Gonolkites convergens* at the base of limestone bed RB071 (bed 23 in Sturani 1967) in the Ravin du Bès Section. The base of the Bathonian in Bas-Auran sections coincides with the lowest occurrence of *Morphoceras parvum*. The basal ammonite assemblage encloses the following ammonite species (Fig. 10):

Oxycerites limosus (Buckman) [M],
Cadomites deslongchampsii (d'Orbigny) [M+m],
Cadomites crassispinosus Kopik [M+m],
Cadomites stegeus (Buckman) [M+m],
Cadomites psilacanthus (Wermbter) [M+m],
Cadomites gr. *rectelobatus* (Hauer) [M],
Parkinsonia subplanulata Wetzel [m+M],
Gonolkites subgaleatus (Buckman) [M],
Gonolkites convergens Buckman [M],
Morphoceras parvum Wetzel [M].

Ammonite content of the basal Bathonian fossil-assemblage is virtually enlarged by the following taxa, known from below and above but not actually in the basal bed:

Cadomites sturani Galász [M+m],
Polyplectites rozyckii (Kopik) [m],
Parkinsonia cf. *subplanulata* Wetzel [m+M],
Parkinsonia crassa Nicolesco [m+M],
Parkinsonia schloenbachi Schlippe [m+M],
Planisphinctes planilobus Buckman [m],
Phaulozigzag phaulomorphus Buckman [m].

The following species of Phylloceratina and Lytoceratina have been, actually or virtually, identified in the Bathonian basal fossil-assemblage also (Fig. 9):

Phylloceras kudernatschi (Hauer),
Adabofoloceras subobtusum (Kudernatsch),
Adabofoloceras wendti (Sturani),
Phyllopachyceras ebrayi (Ferry),
Calliphylloceras achtalense (Redlich),
Calliphylloceras gr. *disputabile* (Zittel),
Nannolytoceras tripartitum (Raspail),
Lytoceras gr. *eudesianum* (d'Orbigny).

New palaeontological data about the youngest members of Bigotitinae and the oldest members of Zigzagiceratinae are of biochronostratigraphic importance for the subdivision and correlation of the basal Bathonian Zigzag Zone. Three successive biohorizons have been identified and chronocorrelated between the Bas-Auran (French Subalpine Basin) and Cabo Mondego (Lusitanian Basin) successions: Diniensis, Mondegoensis and Protozigzagiceratas biohorizons (Fernández-López *et al.* 2007, Pavia *et al.* 2008). The Diniensis Biohorizon is characterized by the occurrence of *Bigotites diniensis* representatives and it corresponds to the lowest part of the Bathonian Zigzag Zone in the Sub-Mediterranean Province (e.g., Cabo Mondego and Bas-Auran). It encompasses the stratigraphic intervals RA085–RA062 (Fig. 10, levels 23–18 of Sturani 1967) in Ravin d'Auran Section and RB071–RB054 (Fig. 7, levels 23–18 of Sturani 1967) in Ravin du Bès Section. The Mondegoensis

Biohorizon is defined by the lowest occurrence of *Bigotites mondegoensis* representatives. It comprises the stratigraphic intervals RA061–RA044 (Fig. 10, levels 17–14 of Sturani 1967) in Ravin d’Auran Section and RB053–RB034 (Fig. 7, levels 17–14 of Sturani 1967) in Ravin du Bès Section, taking into account the occurrence of *B. mondegoensis* at the level RA061. The Protozigzagiceras Biohorizon is defined by the lowest occurrence of Zigzagiceratinae representatives, in particular *Protozigzagiceras* [M+m] and *Franchia* [M+m]. It encompasses the stratigraphic intervals RA043–RA034 (Fig. 7, level 13 of Sturani 1967) in Ravin d’Auran Section and RB033–RB026 (level 13 of Sturani 1967) in Ravin du Bès Section, taking into account the occurrence of *Protozigzagiceras* from Sturani’s level 13 and *Franchia* from RA035.

According to the results of Pavia *et al.* (2008), the record quality of the ammonoid biostratigraphic succession in the Bas-Auran area can be tested with various palaeontological criteria: the preservation state of fossil-specimens, taphonic populations and fossil-assemblages; abundance, concentration, packing and stratigraphic persistence of fossil-specimens; completeness, constancy and persistence of stratigraphic ranges; completeness and taxonomic diversity of successive fossil-assemblages; biostratigraphic turnover; proportion of virtual and actual palaeontological gaps in successive stratigraphic intervals; proportion of first and last occurrences of taxa; proportion of lowest and highest occurrences of taxa; successive or coincident clustering of last and first occurrences. Values of these twenty-one palaeontological attributes indicate relatively homogeneous and good record quality, gradual biostratigraphic change and high degree of taxonomic similarity between the Bomfordi and Convergens subzones. These criteria, applied to the ammonoid genera which are known from the Bas-Auran area, also indicate relatively high values of palaeontological and stratigraphic completeness at the base of levels RB070–RB071 (= level 23 in Sturani 1967; i.e., the Bajocian/Bathonian boundary). The ammonoid biostratigraphic succession of Bas-Auran shows no evidence of biochronostratigraphic mixing, taphonomic condensation, signs of non-sequence or biostratigraphic discontinuities across the Bajocian/Bathonian boundary interval. Moreover, with forty-six successive ammonoid fossil-assemblages of the Convergens Subzone, along up to 5 metres of thickness belonging to three biohorizons, the Ravin du Bès Section displays maximum values of biostratigraphic and biochronostratigraphic completeness.





Fig. 9. Phylloceratina and Lytoceratina biochronostratigraphic data at the Bajocian/Bathonian boundary in the Ravin du Bès Section (from Pavia *et al.* 2008).



Microfossils

The Bajocian/Bathonian boundary may be characterized by secondary (auxiliary) biostratigraphic markers, such as nannofossils. According to the results of Erba (1990a, b; Cobianchi *et al.* 1992; Mattioli & Erba 1999), calcareous nannofossils are present in all beds and facilitate the characterization of the Bajocian-Bathonian transition. The Ravin du Bès Section appears to be suitable for the biostratigraphical study of microfossils, such as foraminifers or ostracods, but there are no published studies. According to preliminary results (Bodergat in Mangold 1999), ostracods are present in all marly samplings, but are badly preserved between bed 23 and bed 13. The marine taxa are different from those known in the Paris basin and England. The Subalpine taxa, specially the genera *Pontocyprilla*, *Isobythocypris* and *Cordobairdia*, indicate deeper environments (more than 200 m). Palynomorphs are poorly preserved and are not yet stratigraphically useful across the boundary (Poulsen 1997, Mangold 1999).

Calcareous nannofossils (E. Erba & D. Tiraboschi)

Nannofossil biostratigraphic investigation was performed on 59 samples (approximately every 20 cm) collected from the Ravin du Bès section in the Bas-Auran area; sample figures correspond to the bed numbers of the lithostratigraphic column of Fig. 7. This study is a revision of the previous work by Erba (1990a, b), extended to limestone layers and additional marlstone beds. Simple smear slides were prepared for both limestone and marlstone, using standard techniques and without centrifuging cleaning /concentration in order to retain the original sediment composition. Few milligrams of powdered sediments were mounted on a glass with the Norland Optical Adhesive, and then analyzed using a light polarizing microscope, at 1250x magnification.

All studied samples contain calcareous nannofossils. A total of 37 taxa were identified and their distribution is given in Fig. 11. The nannofossil total abundance fluctuates from extremely rare to common; the preservation is poor to moderate, with evidence of dissolution and overgrowth. Limestone levels generally contain depauperated and poorly preserved nannofloras, with stronger overgrowth and dissolution.

The nannofloras are characteristic of the Upper Bajocian–Lower Bathonian interval. Assemblages are dominated by *Watznaueria britannica* and *Watznaueria communis*, with common *Schizosphaerella punctulata*, *Watznaueria* aff. *W. manivitiae*, *Watznaueria manivitiae*, *Cyclagelosphaera margerelii*, *Cyclagelosphaera deflandrei*, *Lotharingius crucicentralis*, *Lotharingius velatus*, *Lotharingius sigillatus* and *Ethmorhabdus gallicus*.

Based on absence of *Carinolithus superbus* and of *Watznaueria barnesiae*, the lowermost portion of the investigated interval (samples 110 through 68b) corresponds to the Tethyan *W. communis* Subzone (NJT 10b) indicating a Late Bajocian age (Mattioli & Erba 1999). This subzone corresponds to the upper part of the Boreal NJ 10 Zone and the lower part of the NJ 11 Zone of Bown & Cooper (1998). The first occurrence (FO) of *Pseudoconus enigma* in sample 89 identifies the NJ10/NJ11 zonal boundary (Figs. 12–13). This taxon is rare and occurs only in limestones, with the only exception of a single specimen in marlstone sample 20, and that's why Erba (1990b) did not report this species.

The last occurrence (LO) of *Hexalithus magharensis* was observed in sample 82 indicating a latest Bajocian age (Mattioli & Erba 1999). Similarly, Erba (1990b) recorded this event in the Parkinsoni Zone (latest Bajocian) of the Digne area, whereas Kaenel *et al.* (1996) found an older age for the LO of *H. magharensis*, calibrated between the end of the Early Bajocian and the beginning of the Late Bajocian.

The FO of *Stephanolithion speciosum octum* was observed in sample 76; the taxon is extremely rare and scarce in the studied section. This event has been correlated to the base of the Parkinsoni Zone in NW Europe and Portugal (Kaenel *et al.* 1996), but within the Zigzag Zone in SE France (Erba 1990b). Bown *et al.* (1988) and Bown & Cooper (1998) report the FO of *S. speciosum octum* at the base of the Boreal NJ 11 Zone.

The FO of *W. barnesiae* (NJT11) was observed in sample 68a of earliest Bathonian age (Zigzag Zone). This event defines the base of the Tethyan NJT11 Zone (Mattioli & Erba 1999), comparable to most of the Boreal NJ11 Zone and NJ12a Subzone (Bown *et al.* 1988, Bown & Cooper 1998).

The uppermost portion of the studied interval corresponds to the Tethyan NJT 11 Zone (Mattioli & Erba 1999), since *Cyclagelosphaera wiedmannii* was not observed.

From sample 89 upwards, rosetta-shaped specimens likely of the genus *Rucinolithus* were consistently observed. They show highest abundance in the interval between sample 45 through 22 (Fig. 11), both in limestone and marlstone beds. Two morphotypes were distinguished, namely small (< 8 microns) and large (> 8 microns) *Rucinolithus* spp., based on their diameter. More detailed investigations are in progress to characterize the taxonomy of these morphotypes (Tiraboschi & Erba *in prep.*).

Our results are consistent with previous biostratigraphic data from the Upper Bajocian–Lower Bathonian interval in SE France (Erba 1990b), Portugal, NW Europe (Kaenel & Bergen 1993, Kaenel *et al.* 1996), Lombardian Basin (Chiari *et al.* 2007) and Boreal Realm (Bown & Cooper 1998). For the first time *P. enigma* has been documented from mid to low latitudes allowing a direct calibration between Tethyan and Boreal nannofossil events and biozones (Figs. 12–13).




Fig. 12. Nannofossil events detected in the Ravin du Bès Section. Tethyan biozones after Mattioli & Erba (1999) and Boreal biozones after Bown & Cooper (1998).




Fig. 13. Upper Bajocian and Lower Bathonian calcareous nannofossils from the Bas-Auran area. All specimens at 1250x magnification.

1. *Watznaueria communis*, crossed nicols, sample 110.
2. *Watznaueria britannica*, crossed nicols, sample 102.
3. *Watznaueria fossacincta*, crossed nicols, sample 110.
4. *Watznaueria* aff. *W. barnesiae*, crossed nicols, sample 60.
5. *Watznaueria barnesiae*, crossed nicols, sample 68a.
6. *Watznaueria barnesiae*, crossed nicols, sample 10.
7. *Cyclagelosphaera margerelii*, crossed nicols, sample 40.
8. *Watznaueria manivittiae*, crossed nicols, sample 67.
9. *Watznaueria* aff. *W. contracta*, crossed nicols, sample 48.
10. *Stephanolithion speciosum octum*, transmitted light, sample 76.
11. *Hexalithus magharensis*, crossed nicols, sample 110.
12. *Hexalithus magharensis*, transmitted light, sample 110.
13. *Stephanolithion speciosum speciosum*, transmitted light, sample 76.
14. *Rucinolithus* sp., crossed nicols, sample 6.
15. *Pseudoconus enigma*, crossed nicols, sample 53.

Correlation (S.R. Fernández-López)

Ammonites are the most relevant taxonomic group for global biochronostratigraphic correlation of the Bajocian/Bathonian boundary. Late Bajocian and Early Bathonian ammonites are found worldwide in the three major, oceanic or marine, palaeogeographical units: Tethyan, Pacific and Boreal domains (Cariou *et al.* 1985; Hillebrandt *et al.* 1992a, b; Taylor *et al.* 1992; Westermann 1993a, 2000; Page 1996a; Enay & Cariou 1999).

Figure 14 shows standard zonations for the four ammonite biogeographical provinces represented in western Europe. Ammonites of the Zigzag Zone have a wide distribution through the Northwest European, Sub-Mediterranean and Mediterranean provinces of the West Tethyan Subrealm. The Northwest European Province, in which parkinsoniids are common, comprises the following epeiric areas: England (Torrens 1980; Callomon 1995, 2003; Callomon & Cope 1995; Page 1996b, 2001; Dietze and Chandler 1998; Chandler *et al.* 1999), Normandy, Boulonnais, Lorraine, Alsace, northern Jura (Mangold & Rioult 1997, Rioult *et al.* 1997, Thierry 2003), northern Germany (Westermann 1958; Metz 1990, 1992), northern and central Poland (Kopik 2006, Zaton 2006).

The Sub-Mediterranean Province, in which Bathonian *Morphoceras* [M] - *Ebrayiceras* [m] occur associated with parkinsoniids and scarce phylloceratids and lytoceratids, includes the following epeiric areas: Lusitanian Basin (Fernández-López *et al.* 2006a, b), Iberian Basin (Fernández-López 2000, 2001), Aquitaine, Causses, Centre-west France, Nièvre (Delance *et al.* 1979, Courville *et al.* 1999, Enay *et al.* 2001), Mâconnais, Ardèche, southern Jura (Elmi 1967; Mangold 1971a, b, c, 1997a, b; Rulleau 2006), western Alps and Subalpine Basin (Sturani 1967, Pavia & Sturani 1968, Pavia 1973, 1984, Torrens 1987, Innocenti *et al.* 1990, Zany *et al.* 1990, Joly 2000), southern Germany (Dietl 1978, 1981, 1982, 1983, 1986, 1988; Dietl *et al.* 1978, 1983; Dietl & Hugger 1979; Dietl & Kapitzke 1983; Callomon *et al.* 1987; Schairer 1987, 1994; Dietze & Chandler 1996; Köstler & Schairer 1996; Dietze *et al.* 1997, 2002, 2004; Schweigert & Dietze 1998; Dietze 2000; Dietze & Schweigert 2000; Schweigert *et al.* 2002, 2003; Ohmert *et al.* 2004; Dietze & Dietl 2006), peri-Carpathian Poland (Wierzbowski *et al.* 1999; Luczynski *et al.* 2000; Matyja & Wierzbowski 2000, 2001; Zaton & Marynowski 2006), western Carpathians (Schlögl & Rakús 2004; Schlögl *et al.* 2005, 2006), South Transdanubian Mecksek (Galác 1995a, Geczy & Galác 1998), Romania (Galác 1994, Patruilus 1996), Balkans (Stephanov 1972), northwestern-central Iran (Seyed-Emami *et al.* 1985, 1989, 1991, 1998a, b) and north-eastern Iran (Majidifard 2003).

The Mediterranean Province, in which Late Bajocian and Early Bathonian morphoceratids occur associated with common phylloceratids and lytoceratids, comprises the following shelfal or oceanic areas: Betic Basin (Mangold 1981; Sandoval 1983, 1986, 1990; Sandoval *et al.* 2001), Majorca (Sandoval 1994), Sicily (Wendt 1963, 1971; Galác 1985, 1999a, b; Pavia & Cresta 2002; Pavia *et al.* 2002; Martire & Pavia 2004; Pavia 2007), Alps (Sturani 1971, Krystyn 1972, Joly 2000, Martire 1989, Mangold & Gygi 1997), North Transdanubian Bakony (Galác 1980, 1993, 1995b).

In the north-eastern Tethyan border (Donetz, Crimea, Caucasus, Great Balkhan, Turkmenistan, Tadzhikistan, Uzbekistan, Kazakhstan) latest Bajocian to Early Bathonian parkinsoniids and morphoceratids have been described, below Middle Bathonian specimens of *Bullatimorphites*, *Procerites* and *Siemiradzka* (Rostovtsev 1985; Tseretely 1989; Beznosov & Mitta 1998, 2000).

In Tibet and South-East Asia, Bathonian *Siemiradzka*, *Procerites* and *Wagnericeras* have been identified (Cariou & Enay 1999, Yin *et al.* 2000, Yin 2005). Upper Bajocian *Leptosphinctes* and *Cadomites* have been recognized in Japan, below Upper Bathonian *Pseudoneuquenicer* (Sato 1992).

Lower Bathonian morphoceratids, parkinsoniids and phylloceratids have been recognized in several basins of the southern Tethyan border: Morocco, Algeria and Tunisia (Elmi 1971, Elmi & Almería 1984, Enay *et al.* 1987b, Ouahhabi 1994, Soussi *et al.* 2000). *Oraniceras*, *Micromphalites* and *Oxycerites* occur in Lower Bathonian, whereas *Ermoceras*, *Leptosphinctes* and *Oppelia* characterize Upper Bajocian deposits.

In the Ethiopian Province, including Libya, Egypt, Israel, southern Turkey, southern Iran and Saudi Arabia (Parnes 1981, 1984, 1988; Enay *et al.* 1986, 1987a; Enay & Mangold 1994, 1996), the Lower Bathonian comprises the Tuwaiqensis (*Tulites*) and Clydocromphalus (*Micromphalites*) zones of the Arabian Province, including *Procerites* and *Zigzagiceras* in basal levels. Upper Bajocian deposits with *Ermoceras*, *Thambites*, *Leptosphinctes* and *Spiroceras* represent the Runcinatum (*Ermoceras*), Mogharensis (*Ermoceras*) and Planus (*Thambites*) zones.

		NW European Province	Sub-Mediterranean Province	Mediterranean Province	Boreal Province			
		England, Normandy, Boulonnais, Lorraine, Alsace, northern Germany, northern and central Poland.	Portugal, Iberian Basin, Aquitaine, Causses, Centre-west France, Nièvre, southern Jura, Mâconnais, Ardèche, southern Germany peri-Carp. Poland, Balkans, northern and central Iran.	Betic Basin, Sicily, Appennines, Switzerland, Austria, Hungary p.p (Villany and Mecsek), Hellenids, Serbia.	Eastern Greenland, Siberia, northern Alaska, northern Canada.			
Lower Bathonian	Zigzag	Tenuiplicatus	Aurigerus	Tenuiplicatus	Postpollubrum	Ishmae	14	
		Yeovilensis		Recinctus		Zigzag	Yeovilensis	Greenlandicus
		Macrescens	Zigzag	Macrescens	Macrescens		11	
		Convergens		Parvum	Dimorphitiformis	Arcticus	10	
Upper Bajocian	Parkinsoni	Bomfordi	Parkinsoni	Bomfordi	Parkinsoni	Dimorphus	9	
		Truellei		Densicosta		Daubenyi	Pompeckji	8
				Acris				

Fig. 14. Ammonite zones and subzones of the Uppermost Bajocian and Lower Bathonian in several palaeobiogeographical provinces: Northwest European (Westermann & Callomon 1988, Callomon & Cope 1995), Sub-Mediterranean (Mangold 1990, Rioult *et al.* 1997, Mangold & Rioult 1997), Mediterranean (Sandoval 1983, 1990; Galacz 1980, 1993) and Boreal (Callomon 2003) provinces.

In the Indo-Malgach Province, Late Bajocian and Middle Bathonian ammonites, but not Early Bathonian, have been described from Kenya, Madagascar and India (Singh *et al.* 1982, 1983; Jaitly & Singh 1983, 1984; Pandey & Agrawal 1984; Pandey & Westermann 1988; Galacz 1990; Pandey & Callomon 1995; Roy *et al.* 2007).

In south-western Pacific areas, Sula, Irian Jaya and New Guinea, latest Bajocian to Early Bathonian *Pretulites* and Early to Middle Bathonian *Satoceras*, as well as Bathonian specimens of *Asphinctites*, *Rugiferites* and *Bullatimorphites* have been described (Westermann & Getty 1970, Westermann & Callomon 1988, Sukanto & Westermann 1992, Westermann 1995, Callomon & Rose 2000).

Separate Late Bajocian and Early Bathonian ammonoid faunas have been distinguished, associated with characteristic Sphaeroceratinae and Eurycephalitinae, in the southern East-Pacific Subrealm of the Tethyan Realm: New Zealand (Westermann & Hudson 1991; Westermann 1993b; Westermann *et al.* 2000, 2002), Argentina, Chile and Peru (Westermann & Riccardi 1980; Westermann *et al.* 1980; Riccardi 1985, 1991; Riccardi *et al.* 1990a, b, 1991, 1994; Riccardi & Westermann 1991a, b, 1999; Hillebrandt *et al.* 1992a, b; Fernández-López *et al.* 1994; Gröschke & Hillebrandt 1994; Hillebrandt 1995, 2001; Gröschke 1996; Parent 1998). *Leptosphinctes*, *Lupherites*, *Strenoceras*, *Spiroceras* and *Megasphaeroceras* occur in the Upper Bajocian of the Andean Province. *Lobosphinctes intersertus* Buckman has been identified in Chacay Melehue (Argentina) below a Bathonian *Cadomites*-Tulitidae mixed assemblage. The first occurrence of several genera such as *Oxycerites*, *Zeissoceras*, *Prohectioceras* and *Rugiferites*, below the oldest representatives of Bathonian *Bullatimorphites*, have been used to recognize Lower Bathonian deposits. New species of Bathonian *?Zigzagiceras* and *Morphoceras* have been proposed (Gröschke & Hillebrandt 1995; Riccardi & Westermann 1999).

In Mexico (Sandoval & Westermann 1986, Sandoval *et al.* 1990) Upper Bajocian begins with the upper Floresi Zone of Oaxaca, containing the Mediterranean *Subcollina lucretia* (Orbigny). The overlying Zapotecum Zone includes *Parastrenoceras*, *Leptosphinctes* and *Oppelia*. The Upper Bathonian Retrocostatum Zone has been identified by *Prohectioceras blanazense*, associated with *Epistrenoceras*, *Lilloettia* and *Neuquenicerias*.

In the Western Interior of the United States of America (Schoshonean Province, Imlay 1981), western Canada and southern Alaska (Athabaskan Province; Imlay 1980, 1982, 1984; Hall & Westermann 1980; Hall & Stronach 1981; Callomon 1984; Hall 1984, 1988, 1989; Poulton *et al.* 1991, 1994) the Upper Bajocian Rotundum Zone includes *Leptosphinctes*, *Lupherites*, *Spiroceras* and *Megasphaeroceras*, below the *Epizigzagiceras-Parareineckeia* association. The *Parachondroceras-Sohlites* assemblages from Oregon may be Upper Bajocian or Lower Bathonian (Imlay 1984).

The Boreal Realm (Eastern Greenland, Siberia, Northern Alaska and Northern Canada) became clearly differentiated in the Late Bajocian and several zonations for the Early Bathonian have been proposed. The Cardioceratidae, in particular *Cranocephalites* and *Arctocephalites*, constituted characteristic elements of the Boreal Realm at the Bajocian/Bathonian boundary (Callomon 1985). The Zone of *Arctocephalites arcticus* (Newton & Teall), above the Zone of *Cranocephalites pompeckji* (Madsen), may represent the basal Bathonian zone in the Boreal Realm (Callomon 1993, 1994, 2003; Rawson 1982; Zakharov *et al.* 1998). The Zone of *Arctocephalites spathi* from northern Yukon probably is coeval with the Boreal Arcticus Zone of eastern Greenland (Poulton 1987). Boreal Arctocephalitinae are associated with parkinsoniids in the south-eastern part of the Russian platform, allowing the correlation between the regional Michalskii-Besnosovi zonal boundary and the Boreal Arcticus-Greenlandicus boundary or the Northwest European Parkinsoni-Zigzag boundary (Mitta 2001, 2004, 2005, 2006, 2007; Mitta & Seltzer 2002; Mitta *et al.* 2004; Saltykov 2007; Zakharov 2007).

Several authors have proposed diverse biozonations for the Upper Bajocian and Lower Bathonian based in different taxonomic groups of macroinvertebrates (Fig. 15): brachiopods (Manceñido & Dagys 1992, Vörös 2001, Alméras *et al.* 2007), belemnites (Challinor 1992, Challinor *et al.* 1992, Combémoré 1997), nautiloids (Branger 2004), bivalves (Damborenea *et al.* 1992, Hallam 1994, Damborenea 2002, Ruban 2006), echinoderms (Thierry *et al.* 1997, Moyne *et al.* 2005), corals (Beauvais 1992).

	Ammonite NW European standard zonation		Brachiopods (Alméras <i>et al.</i> 2007)		Belemnites Combémoré, 1997	
			North-western Tethyan border	Southern Tethyan border		
Lower Bathonian	Tenuiplicatus		Cymatorhynchia reynesi (ex. Formosarhynchia dumortieri)	Rugitela cadomensis	Duvalia disputabilis (partim)	
	Zigzag	Yeovilensis		Tubithyris whatleyensis		Rugitela cadomensis
		Macrescens		Caucasella voutensis		Sphaeroidothyris szajnochai
		Convergens				Burmishynchia athiensis
Upper Bajocian	Parkinsoni	Bomfordi	Megateuthis elliptica (partim)			
		Truellei				
			Callirhynchia oranensis Cymatorhynchia reynesi			

Fig. 15. Zonations for brachiopods and belemnites (from Alméras *et al.* 2007 and Combémoré 1997).

The following taxonomic groups of microfossils are of biochronostratigraphic relevance also (Fig. 16): foraminifera (Bassoulet 1997, Ruget & Nicollin 1997, Gräfe 2005, Cai *et al.* 2006, Saltykov 2007, Wernli & Görög 2007), ostracods (Braun & Brooke 1992, Bodergat 1997), dinoflagellate cysts (Riding & Thomas 1992, Fauconnier 1997, Poulsen & Riding 2003), radiolarians and calcareous nannofossils (Pessagno & Mizutani, 1992, Baumgartner *et al.* 1995, Cordey *et al.* 2005, Chiari *et al.* 2007). Palaeobotanical and palynological data have been recently published by: Kimura *et al.* 1992, Sarjeant *et al.* 1992, Cleal & Rees 2003, Wang *et al.* 2005, Vaez-Javadi & Mirzaei-Ataabadi 2006, Jana & Hilton 2007.

		Ammonite NW European standard zonation		Ostracoda Bodergat, 1997	Foraminifera Ruguet & Nicollin, 1997	Dinoflagellate cysts Riding & Thomas, 1992 Fauconnier, 1997
Lower Bathonian	Tenuiplicatus		Levis - Bathonica (<i>partim</i>)	Quasicitrella (<i>partim</i>)	L. quenstedti, L. galeata, L. polymorpha and L. argonauta (<i>partim</i>)	Ctenidodinium predae (<i>partim</i>)
	Zigzag	Yeovilensis		Bessinensis -Malzi- Bessinensis		
		Macrescens				
		Convergens				
Upper Bajocian	Parkinsoni	Bomfordi	Regularis - Richteri (<i>partim</i>)		Acanthaulax crispa (<i>partim</i>)	
		Truellei				

Fig. 16. Zonations for ostracods (from Bodergat, 1997), foraminifera (from Ruguet & Nicollin, 1997) and dinoflagellate cysts (from Riding & Thomas 1992, Fauconnier 1997).

Isotope stratigraphy

From a geochemical point of view, in the French Subalpine Basin during the Jurassic Period, several authors have emphasized that the manganese content of pelagic carbonates is related to second-order sea-level changes and episodes of hydrothermal activity that affected the chemistry of global sea water. The main transgressive phases are marked by a manganese content increase, whereas regressive phases are characterized by decreasing trends (Corbin 1994, Corbin *et al.* 2000). In the Chaudon-Norante section, 4 km north of the Bas-Auran area, the Early Bathonian maximum transgressive is marked by sedimentary condensations, associated with high manganese content (from 300 to 1370 mg kg⁻¹). In contrast, the Middle and Late Bathonian regressive phase coincides with low manganese content periods. These stratigraphical patterns in divalent manganese can be of either local or regional significance, being concentrated, most probably as a very early diagenetic phase, only in oxygen-depleted waters that typically underlie zones of elevated organic productivity (Jenkyns *et al.* 2002). For strontium isotope (⁸⁷Sr/⁸⁶Sr ratio), oxygen isotope ($\delta^{18}\text{O}$) or carbon isotope ($\delta^{13}\text{C}$) chemostratigraphy, no data are currently available.

Volcanogenic deposits suitable for direct radio-isotope dating are not known in the section. The age of the Bajocian/Bathonian boundary has been dated 167.7 ± 3.5 Ma by Gradstein & Ogg (2004) and Gradstein *et al.* (2005).

Magnetostratigraphy (R. Lanza)

In the autumn 1994, the Bas-Auran section was extensively sampled across the Bajocian/Bathonian boundary. Some 30 hand-samples were collected and eventually cored in the rock-magnetism laboratory of the Torino University. All specimens were measured with a JR-5 spinner magnetometer and thermally demagnetized using a Schonstedt furnace (Degiorgis 1996).

The specimens are characterized by three remanent magnetization components:

— a viscous (VRM) component close to the present field and removed at temperature values about 120 °C

— a secondary component stable up to values of 300 to 450 °C.

— a high-temperature component, stable between 350–450 and 480–500 °C.

The secondary component has been interpreted as a Tertiary magnetic overprint: declination points to the NW (310° to 330°), inclination is positive (30° to 50°).

The high-temperature component has been isolated only in 11 out of the 32 analyzed specimens. Its definition is difficult and often poor, because its intensity is very low, usually 10–25 % of the initial NRM. This component may be regarded as the more stable fraction of the primary Jurassic remanence acquired when the rocks formed. The fact that the primary remanence can be isolated only in few specimens and its poor definition prevent any reliable magnetostratigraphic interpretation.

The results found by Degiorgis (1996) in the Bas-Auran section have been fully substantiated in the whole southern Subalpine Chains by Aubourg & Chabert-Pelline (1999).

Gamma-ray spectrometry

Field gamma-ray spectrometry data have been obtained by G. Pavia, P. Lazarin and L. Leroy (April 2007) and are presented in Fig. 17. Spectral gamma-ray data from the Ravin du Bès Section show an increase in the total gamma-ray counts at the Aurigerus Zone. The values are relatively low and display insignificant variation at the Bajocian-Bathonian boundary, but they show a positive peak at the top of the Lower Bathonian. Total gamma-ray logs have been used in sequence stratigraphy on the basis that gamma-ray peaks commonly correspond to maximum flooding surfaces (cf. Parkinson 1996; Deconinck *et al.* 2003; Pawellek & Aigner 2003, 2004; Pellenard *et al.* 2003; Raddadi *et al.* 2005; Ruf *et al.* 2005; Schnyder *et al.* 2006). High gamma-ray counts, low sedimentation rates and high concentrations of ammonites may be associated with the development of condensed sections. These features, however, developed both in condensed deposits of deep carbonate environments during transgressions or episodes of relative sea-level rise and in expanded deposits of shallow carbonate epicontinental platforms during regressions or episodes of relative sea-level fall (Fernández-López *et al.* 2002). The stratigraphic trend in spectral gamma-ray data associated with sedimentary condensation on the Bas-Auran area, from the Bajocian Bomfordi Subzone towards Bathonian Tenuiplicatus Subzone, provides support for an Early Bathonian deepening half-cycle of second order, lacking evidence of stratigraphic gaps at the Bajocian-Bathonian transition.




Fig. 17. Gamma-ray log of the Ravin du Bès Section, showing a positive peak at the top of the "Marno-calcaires à *Cancellophycus*" in the Lower Bathonian Tenuiplicatus Subzone (Aurigerus Zone).

Protection of the site (M. Guiomar)

The sites of Bas-Auran and Le Bès are part of the protected territory of “La Réserve Naturelle Géologique de Haute Provence”. The Geological Reserve, which covers 55 communes in the “Alpes de Haute-Provence” and Var departments, has been entrusted with the main missions of protecting, enhancing and raising awareness to the environment and supporting economic development of this heritage. These different missions are based on scientific knowledge of geological sites and inventories. This is why the Reserve wishes to encourage the number of scientific studies of its territory. While some of this research is instigated and supported by the Reserve itself and conducted in collaboration with its partners, academic or other, there is also independent research, conducted with technical support from the Reserve, in particular for the authorization to collect fossils. Two types of regulations apply in protected territory: those relative to sites listed as natural reserves (Réserve Naturelle) by Ministerial decree and those relative to protected areas, by Prefectoral order (arrêtés préfectoraux). This explains why the sites of Bas-Auran and Le Bès do not have the same status. Bas-Auran is listed as a natural reserve (RN): no collecting or surface removal is possible, except by ministerial authorization. The section of Le Bès is part of the protected area where fossils may be collected and authorizations delivered for extraction or excavation (files managed by the Reserve), mainly in the context of research projects. Authorizations delivered by the Reserve in no way affect private property rights and all applications for excavation must be accompanied by a request to the owner of the land. According to the site’s scientific value, the Reserve may request that a geological site be added or removed from listing to facilitate its management and/or protection; exceptionally, the Reserve may approach local communities for the acquisition of certain lands. For the section(s) to obtain GSSP status, it is necessary to undertake all the requisite measures to maintain free access to them as well as consider minimum developments (ease of access, safety) as well as enhancement (to be defined with scientists and managers), whenever possible.

The Bathonian ASP in Cabo Mondego Section (Portugal) (S.R. Fernández-López, M.H. Henriques & C. Mangold)

An auxiliary section and point (ASP) for the base of the Bathonian Stage is located in Cabo Mondego, 40 km west of Coimbra, 7 km north of Figueira da Foz (40°11’19’’N, 8°54’30’’W, Section 02 in Fig. 18). These classical fossiliferous deposits have been studied by numerous specialists (Ruget-Perrot 1961; Elmi *et al.* 1971; Mangold 1971c, 1990; Rocha *et al.* 1981, 1987; Mangold & Rioult 1997; Fernández-López & Henriques 2002; Fernández-López *et al.* 2006a, b).

These Bathonian deposits correspond to the Cabo Mondego Formation and comprise limestone-marl alternations, with ammonoids, bivalves (*Bositra*), rhynchonellid brachiopods, crinoids and belemnites. Bioturbation structures are common (*Zoophycos*, *Thalassinoides*, *Chondrites*). These fossiliferous deposits were developed in an open sea, in hemipelagic environment of distal and outer carbonate ramp, below wave base (Watkinson 1989, Soares *et al.* 1993, Azerêdo *et al.* 2003).

The base of the Bathonian has been established at the base of marly interval 123 of Section-02 (Fig. 19) and at the base of the marly interval FC1 of Section-90. From a biostratigraphic point of view, 10 metres of thickness with 62 successive ammonoid fossil-assemblages from 77 successive fossiliferous stratigraphic intervals have been recognized and sampled in the Parvum Subzone. The base of the Bathonian has been established by the lowest occurrence of representatives of the *Morphoceras* [M] - *Ebrayiceras* [m] group in the marly interval 02CM123. The Lower Bathonian index ammonite *Morphoceras parvum* occurs in the marly interval 02CM139. The Lower Bathonian index ammonite *Gonolkites convergens* occurs in the marly interval 02CM181.

As in the Bas-Auran area, palaeontological data about the youngest members of Bigotitinae and oldest members of Zigzagiceratinae are of particular relevance for the biostratigraphic subdivision and correlation of the basal Bathonian Zigzag Zone. The Diniensis, Mondegoensis and Protozigzagiceras biohorizons of the Parvum Subzone have been recognized in Cabo Mondego (Fernández-López *et al.* 2007). The Diniensis Biohorizon is represented by the stratigraphic intervals 02CM123–02CM145 in the Section 02 and FC1–FC17 in the Section 90 (Fig. 19). The Mondegoensis Biohorizon is represented by the stratigraphic intervals 02CM146–02CM182 in the Section 02 and

FC18–FC43 in the Section 90, taking into account the occurrence of *B. mondegoensis* at the level 02CM146. The Protozigzagiceras Biohorizon is represented by the stratigraphic intervals 02CM183–02CM198 in the Section 02 and FC44–FD11 in the Section 90, taking into account the occurrence of *Protozigzagiceras* at the level 04CM183.

New provisions for the conservation and protection of this Portuguese outcrop have now been implemented under national laws, after the classification of the Cabo Mondego area as a Natural Monument of the Portuguese Republic in 2007 (Henriques & Ramalho 2005, Page *et al.* 2006).




Fig. 18. Location maps of three stratigraphic sections of Cabo Mondego (Portugal, from Fernández-López *et al.* 2006a).




Fig. 19. Ammonite biostratigraphic data at the Bajocian/Bathonian boundary in the Section-90 and Section-02 of Cabo Mondego (Portugal, from Fernández-López et al. 2006ab modified).

Summary

The leading candidate Global Boundary Stratotype Section and Point for the base of the Bathonian Stage at the Ravin du Bès Section (France) satisfies most of the requirements recommended by the ICS (e.g. in [Remane *et al.* 1996](#), [Gradstein *et al.* 2003](#), [Morton 2006](#), cf. [Fig. 20](#)):

- The exposure extends over 13 m in thickness, comprising five metres of fossiliferous levels below the boundary and eight metres above. The stratigraphic succession can be recognized laterally over several hundred metres distance.
- At the Bajocian-Bathonian transition, no vertical (bio-, ichno- or tapho-) facies changes, stratigraphic gaps or hiatuses have been recorded. There is no evidence of taphonomic condensation (i.e. mixture of fossils of different age or different chronostratigraphic units). In relation to the rate of sedimentation, the Bomfordi and Convergens subzones are over 10 m thick.
- Structural complexity, synsedimentary and tectonic disturbances, or important alterations by metamorphism are not relevant constraints in the Bas-Auran area.
- The hemipelagic, bed-scale limestone-marl alternations show a maximum value of biostratigraphic completeness for the Bajocian/Bathonian transition. Through five metres of thickness, forty-six successive ammonoid fossil-assemblages in Ravin du Bès Section belonging to three biohorizons of the Parvum Subzone have been recognized. The Bomfordi Subzone attains a minimum thickness of 5 m and includes 42 successive ammonoid fossil-assemblages.
- The boundary has been characterized by both primary and secondary (auxiliary) biostratigraphic markers. There is a well-preserved, abundant and diverse fossil record across the boundary interval, with key markers (ammonites and nannofossils) for worldwide correlation of the uppermost Bajocian and Lower Bathonian. The section appears to be suitable for biostratigraphic study of microfossils, such as foraminifera, but as yet there are no published studies.
- Regional analyses of sequence stratigraphy and manganese chemostratigraphy are available. A transgressive systems tract associated with a deepening phase and sedimentary starvation, within 3rd and 2nd order deepening/shallowing cycles, was developed in the Bas-Auran area of the French Subalpine Basin, during the Early Bathonian. No data are currently available for strontium isotope ($^{87}\text{Sr}/^{86}\text{Sr}$ ratio), oxygen isotope ($\delta^{18}\text{O}$) or carbon isotope ($\delta^{13}\text{C}$) chemostratigraphy.
- The stratigraphic trend in spectral gamma-ray data provides support for an Early Bathonian deepening half-cycle of second order, lacking evidence of stratigraphic gaps at the Bajocian-Bathonian transition.
- Bajocian and Bathonian deposits have been remagnetized with a steady normal polarity. The requirement of suitability for magnetostratigraphy and geochronometry, however, can be indirectly satisfied by reference to the Bathonian magnetostratigraphic scale of [Steiner *et al.* \(1987\)](#), [O'Dogherty *et al.* 2006](#)) as defined in the Subbetic Cordillera.
- Volcanogenic deposits suitable for direct radio-isotope dating are not known in the section. According to the data published by [Gradstein & Ogg \(2004\)](#) and [Ogg \(2004\)](#), the age of the Bajocian/Bathonian boundary is 167.7 ± 3.5 Ma in other basins.
- The criteria of accessibility, conservation and protection are assured by the “Réserve Naturelle Géologique de Haute Provence”, protected under national law and recognised by UNESCO. The park is managed by the “Centre de Géologie de Digne”.
- The Cabo Mondego Section (Portugal) is suggested as the Bathonian ASP within the same GSSP proposal. It provides complementary biochronostratigraphic data mainly related to the ammonite succession at the Sub-Mediterranean Parvum Subzone and its correlation with the Northwest European Convergens Subzone. Accessibility, conservation and protection are guaranteed, after the classification of the Cabo Mondego area as a Natural Monument of the Portuguese Republic in 2007.

The requirements for a GSSP (ICS)	Ravin du Bès Section (Bas-Auran)
GEOLOGICAL REQUIREMENTS	
Exposure over an adequate thickness	Yes
Continuous sedimentation. No gaps or condensation close to the boundary	Yes
Rate of sedimentation	About 4.5 m for the Parvum Subzone and at least 5 m for the Bomfordi Subzone
Absence of syndepositional and tectonic disturbances	Yes
Absence of metamorphism and strong diagenetic alteration	Yes (for macrofossils)
BIOSTRATIGRAPHIC REQUIREMENTS	
Abundance and diversity of well-preserved fossils	Abundant and well preserved ammonites
Absence of vertical facies changes at or near the boundary	Yes
Favorable facies for long-range biostratigraphic correlations	Yes
OTHER METHODS	
Radioisotopic dating	No information
Magnetostratigraphy	No significant result
Geochronometry	No information
Sequence stratigraphy	Graciansky <i>et al.</i> 1993, 1998
Gamma-ray spectrometry	Yes, supporting sequence-stratigraphy results
OTHER REQUIREMENTS	
GSSP indicated by a permanent fixed marker	Yes, if accepted
Physical and logistical accessibility	Yes
Free access for research	Yes
Permanent protection of the site	Part of the European Geopark: "Réserve Naturelle Géologique de Haute Provence"

Fig. 20. Summary of the requirements of the International Commission on Stratigraphy for Ravin du Bès Section (Bas-Auran).

Acknowledgements. This work has been supported by the CGL2004-0694/BTE (MEC-CSIC) Project and grants from the 2006 and 2007 Geoconservation Projects, respectively coordinated at the Madrid and Torino universities.

References

- Alm eras, Y., Faure, P., Elmi S., Enay, R. & Mangold, C. 2007: Zonation des brachiopodes du Jurassique moyen sur la marge sud de la T ethys occidentale (Maroc, Alg erie occidentale) Comparaison avec la marge nord-t ethysienne fran aise. *Geobios* 40, 1–19.
- Arkell, W.J. 1951–1959: *A monograph of English Bathonian ammonites*. 264 pp. Palaeontographical Society, London.
- Arkell, W.J. 1956: *Jurassic geology of the world*. 806 pp. Oliver & Boyd Ltd., London.
- Aubourg, C. & Chabert-Pelline, C. 1999: Neogene remagnetization of normal polarity in the Late Jurassic black shales from the southern Subalpine Chains (French Alps). Evidence for late anticlockwise rotations. *Tectonophysics* 308 (1999), 473–486.
- Azer do, A.C., Duarte, L.V., Henriques, M.H. & Manuppella, G. 2003: Da din mica continental no Tri sico aos mares do Jur ssico Inferior e M dio. *Cadernos de Geologia de Portugal* 2003, 1–43.
- Bassoullet, J.-P. 1997: Les grandes foraminif res. *Bulletin du Centre de Recherches Elf Exploration Production, M moires* 17, 293–304.
- Baumgartner, O.P., O’Doherty, L., Gorican, S., Urquhart, E., Pillevuit, A. & De Wever, P. 1995: Middle Jurassic to lower Cretaceous Radiolaria of Tethys: occurrences, systematics, Biochronology. *M moires de G ologie* 23, 1–1162.
- Beauvais, L. 1992: Corals of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 324–327. Cambridge University Press, Cambridge.
- Beznosov, N.V. & Mitta, V.V. 1998: *Catalogue of Ammonitida and key sections of the Upper Bajocian – Lower Bathonian of North Caucasus* [in Russian]. 72 pp. VNIGNI, Nedra, Moscow.
- Beznosov, N.V. & Mitta, V.V. 2000: *Jurassic geology and ammonites of Great Balkhan (Western Turkmenistan)* [in Russian]. 115 pp. VNIGNI, Nedra, Moscow.
- Bodergat, A.-M. 1997: Les ostracodes marines du Jurassique europ en. Utilisation stratigraphique. *Bulletin du Centre de Recherches Elf Exploration Production, M moires* 17, 197–223.
- Bown, P.R. & Cooper, M.K.E. 1998: Jurassic. In Bown, P.R. (ed.): *Calcareous Nannofossil Biostratigraphy*, 34–85. Kluwer Academic Publishers, London.
- Bown, P.R., Cooper, M.K.E. & Lord, A.R. 1988: A calcareous nannofossil biozonation scheme for the early to Mid Mesozoic. *Newsletter on Stratigraphy* 20, 91–114.
- Branger, P. 2004: Middle Jurassic Nautiloidea from western France. *Revista Italiana di Paleontologia e Stratigrafia* 110, 141–149.
- Braun, W.K. & Brooke, M.M. 1992: Ostracods of western Canada. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 296–299. Cambridge University Press, Cambridge.
- Buckman, S.S. 1909–1930: *Yorkshire Type Ammonites and Type Ammonites*. 790 pls. Wesley, London.
- Cai, H.-W., Li, J.-G. & Zhang, B.-G. 2006: Early and Middle Jurassic foraminifera from Tingri and Nyalam regions of southern Tibet, China. *Acta Palaeontologica Sinica* 45 (4), 437–452.
- Callomon, J.H. 1984: A review of the biostratigraphy of the post-Lower Bajocian Jurassic ammonites of western and northern North America. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 143–174. Cambridge University Press, Cambridge.
- Callomon, J.H. 1985: The evolution of the Jurassic ammonite family Cardioceratidae. *Special Papers in Palaeontology* 33, 49–90.
- Callomon, J.H. 1993: The ammonite succession in the Middle Jurassic of East Greenland. *Bulletin of the Geological Society of Denmark* 40, 83–113.
- Callomon, J.H. 1994: Jurassic ammonite biochronology of Greenland and the Arctic. *Bulletin of the Geological Society of Denmark* 41, 128–137.
- Callomon, J.H. 1995: Time from fossils: S.S. Buckman and Jurassic high-resolution geochronology. *Geological Society, London, Memoirs* 16, 127–150.
- Callomon, J.H. 2003: The Middle Jurassic of western and northern Europe: its subdivisions, geochronology and correlations. *Geological Survey of Denmark and Greenland Bulletin* 1, 61–73.

- Callomon, J.H. & Cope, J.C.W. 1995: The Jurassic Geology of Dorset. In Taylor, P.D. (ed.): *Field Geology of the British Jurassic*, 51–104. Geological Society, London.
- Callomon, J.H. & Rose G. 2000: Middle Jurassic ammonites from the island of Babar in the southern Moluccan Forearc, Indonesia. *Revue de Paléobiologie* 8, 53–64.
- Callomon, J.H., Dietl, G., Galácz, A., Grandl, H., Niederhöfer, H.J. & Zeiss, A. 1987: Zur Stratigraphie des Mittel- und unteren Oberjuras in Sengenthal bei Neumarkt/Opf. (Fränkische Alb). *Stuttgarter Beiträge zur Naturkunde* 132, 1–53.
- Cariou, E. & Enay, R. 1999: Les ammonites du Bathonien et du Callovien de Thakkhola (Népal Central): biochronologie et intérêt paléobiogéographique. *Geobios* 32, 701–726.
- Cariou, E., Contini, D., Dommergues, J.L., Enay, R., Geysant, J.R., Mangold C. & Thierry J. 1985: Biogéographie des Ammonites et évolution structurale de la Téthys au cours du Jurassique. *Bulletin de la Société géologique de France* 1, 679–697.
- Challinor, A.B. 1992: Belemnites of the southwest Pacific. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 308–310. Cambridge University Press, Cambridge.
- Challinor, A.B., Doyle, P., Howlett, P.J. & Nal'nyaeva, T.I. 1992: Belemnites of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 334–341. Cambridge University Press, Cambridge.
- Chandler, R.B., Glover, L. & Smith, D. 1999: A temporary section in the Inferior Oolite (Middle Jurassic) at Coldharbour Business Park, Dodge Cross, Sherborne. *Proceedings of the Dorset Natural History and Archaeological Society* 120 (1998), 69–72.
- Chiari, M., Cobianchi, M. & Picotti, V. 2007: Integrated stratigraphy (radiolarians and calcareous nannofossils) of the Middle to Upper Jurassic Alpine radiolarites (Lombardian Basin, Italy): Constraints to their genetic interpretation. *Palaeogeography, Palaeoclimatology, Palaeoecology* 249, 233–270.
- Cleal, C.J. & Rees, P.M. 2003: The Middle Jurassic flora from Stonesfield, Oxfordshire, UK. *Palaeontology* 46, 739–801.
- Cobianchi, M., Erba, E. & Pirini Radrizzani, C. 1992: Evolutionary trends of calcareous nannofossil genera *Lotharingius* and *Watznaueria* during the Early and Middle Jurassic. *Memorie Scienze Geologica* 43, 19–25.
- Combémourel, R. 1997: Bélemnites. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires* 17, 157–167.
- Corbin, J.C. 1994: *Evolution géochimique du Jurassique du Sud-Est de la France: influence des variations du niveau marin et de la tectonique*. PhD thesis, Paris VI Univ., unpublished.
- Corbin, J.C., Person, A., Iatzoura, A., Ferre, B. & Renard, M. 2000: Manganese in pelagic carbonates: indication of major tectonic events during the geodynamic evolution of a passive continental margin (the Jurassic European Margin of the Tethys–Ligurian Sea). *Palaeogeography, Palaeoclimatology, Palaeoecology* 156, 123–138.
- Cordey, F., Boughdiri, M. & Sallouhi, H. 2005: First direct age determination from the Jurassic radiolarian-bearing siliceous series (Jédidi Formation) of northwestern Tunisia. *Comptes Rendus Geoscience* 337, 777–785.
- Courville, P., Thierry, J. & Cariou, E. 1999: Modalités évolutives du genre *Bullatimorphites* (Ammonitina) au Bathonien-Callovien (Jurassique moyen) en Europe occidentale. *Comptes Rendus de l'Académie des Sciences de Paris* 323, 81–88.
- Damborenea, S.E. 2002: Jurassic evolution of Southern Hemisphere marine palaeobiogeographic units based on benthonic bivalves. *Geobios M. S.* 24, 51–71.
- Damborenea, S.E., Polubotko, I.V., Sey, I.I. & Paraketsov, K.V. 1992: Bivalves zones and assemblages of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 300–307. Cambridge University Press, Cambridge.
- Deconinck, J.F., Hesselbo, S.P., Debuissier, N., Averbuch, O., Baudin, F. & Bessa, J. 2003: Environmental controls on clay mineralogy of an Early Jurassic mudrock (Blue Lias Formation, southern England). *International Journal of Earth Sciences* 92, 255–266.
- Degiorgis, C. 1996: *Stratigrafia magnetica del Baiociano: la sezione di Puerto Escaño (Cordillera Subbética, Spagna) e la sezione del Bas Auran (Alpes de Haute Provence, Francia)*. 109 pp. Graduate Thesis, University of Torino.

- Delance, J.H., Laurin, B. & Marchand, D. 1979: Observations sur la stratigraphie du Bathonien et du Callovien inférieur dans la région de Saint-Benin-d'Azy. *Bulletin Scientifique de Bourgogne* 32, 71–95.
- Dietl, G. 1978: Die heteromorphen Ammoniten des Dogger (Stratigraphie, Taxonomie, Phylogenie, Ökologie). *Stuttgarter Beiträge zur Naturkunde* 33, 1–97.
- Dietl, G. 1981: Zur systematischen Stellung von *Ammonites subfurcatus* Zieten und deren Bedeutung für die subfurcatum-Zone (Bajocium, Mittl. Jura). *Stuttgarter Beiträge zur Naturkunde* 81, 1–11.
- Dietl, G. 1982: Das wirkliche Fundniveau von *Ammonites aspidoides* OPPEL (Ammonoidea, Mittl. Jura) am locus typicus. *Stuttgarter Beiträge zur Naturkunde* 87, 1–21.
- Dietl, G. 1983: Die Ammoniten-Gattung *Strenoceras* aus dem südwestdeutschen Subfurcaten-Oolith (Bajocium, Mittl. Jura). *Stuttgarter Beiträge zur Naturkunde* 90, 1–37.
- Dietl, G. 1986: Erstnachweis von *Oecoptychius subrefractus* (S. Buckm.) (Ammonoidea) aus dem Unter-Bathonium (Mittl. Jura) von SW-Deutschland. *Stuttgarter Beiträge zur Naturkunde* 87, 1–21.
- Dietl, G. 1988: Der Hamiten-Ton (Ober-Bajocium, Mittl. Jura) in Bauaufschlüssen der neuen Bundesautobahn A8, Streckenabschnitt Aichelberg-Gruibingen. *Jahreshefte der Gesellschaft für Naturkunde in Württemberg* 143, 59–77.
- Dietl, G. & Hugger, R. 1979: Zur Stratigraphie des Ober-Bajocium (Braunjura δ/ϵ -Grenzschichten) der Zollernalb (Schwäbische Alb, Baden-Württemberg). *Stuttgarter Beiträge zur Naturkunde* 43, 1–14.
- Dietl, G. & Kapitzke, M. 1983: Das Bathonium (Mittlerer Jura) zwischen Aalen und Bopfingen, östl. Schwäb. Alb. 1. Mittel-Bathonium. *Stuttgarter Beiträge zur Naturkunde* 93, 1–27.
- Dietl, G., Flaig, R. & Glück, E. 1978: Zur Stratigraphie der Ober-Bajocium (Braunjura δ/ϵ Grenzschichten) am Plettengerg bei Balingen, Württemberg. *Stuttgarter Beiträge zur Naturkunde* 40, 1–16.
- Dietl, G., Hugger R. & Schaaf, D. 1983: Die Lage der Bajocium/Bathonium-Grenze (Mittlerer Jura) in der südwestlichen Schwäbischen Alb, SW-Deutschland. *Jahreshefte der Gesellschaft für Naturkunde in Württemberg* 138, 75–84.
- Dietze, V. 2000: Feinstratigraphie und Ammonitenfauna der Acris-Subzone (Parkinsoni-Zone, Ober-Bajocium, Mittlerer Jura) am Ipfl (östliche Schwäbische Alb, Süddeutschland). *Stuttgarter Beiträge zur Naturkunde* 295, 1–43.
- Dietze, V. & Chandler, R.B. 1996: Die Zone des *Zigzagoceras zigzag*. *Fossilien* 1996/3, 159–166.
- Dietze, V. & Chandler, R.B. 1998: New Ammonites from the Zigzag Bed of Dorset. *Dorset Proceedings* 119 (1997), 109–116.
- Dietze V. & Dietl G. 2006: Feinstratigraphie und Ammoniten-Faunen-horizonte im Ober-Bajocium und Bathonium des Ipfl-Gebietes (Schwäbische Alb, Südwestdeutschland). *Stuttgarter Beiträge zur Naturkunde* 360, 1–51.
- Dietze, V. & Schweigert, G. 2000: Zur Stratigraphie und Ammonitenführung des Ober-Bajociums und Bathoniums, insbesondere der Zigzag-Zone, Convergens-Subzone, von Röttingen (östliche Schwäbische Alb, Südwestdeutschland). *Stuttgarter Beiträge zur Naturkunde* 284, 1–15.
- Dietze, V., Krieger, T. & Schweigert, G. 1997: Über *Oecoptychius subrefractus* (BUCKMAN), *Asphinctites tenuiplicatus* (BRAUNS) und *Polysphinctites secundus* (WETZEL) (Ammonoidea) aus dem Unter-Bathonium (Mittlerer Jura) der Oberpfalz (Nordost-Bayern, Süddeutschland). *Stuttgarter Beiträge zur Naturkunde* 245, 1–25.
- Dietze, V., Schweigert, G., Callomon, J.H. & Gauthier, H. 2002: Garantiana- und frühe Parkinsoni-Zone (Ober-Bajocium, Mittlerer Jura) am Ipfl (östliche Schwäbische Alb, SW-Deutschland) mit Bemerkungen zur Phylogenie der Ammonitengattung *Garantiana* Mascke, 1907. *Stuttgarter Beiträge zur Naturkunde* 315, 1–89.
- Dietze, V., Ermer, G., Görlich, M., Ivankic, Z., Krieger, Th. & Röper, M. 2004: Das Bajocium und Bathonium (Mittel-Jura) bei Greding (Fränkische Alb, Süddeutschland). *Archaeopteryx* 22, 61–74.
- Elmi, S. 1967: Le Lias supérieur et le Jurassique moyen de l'Ardèche. *Documents des Laboratoires de Géologie Lyon* 19, 509–845.

- Elmi, S. 1971: Les faunes à *Prohecticoceras* (Oppeliidae, Ammonitina) du Bathonien inférieur et moyen des confins algéro-marocains. *Geobios* 4, 243–264.
- Elmi, S. & Alméras, Y. 1984: Physiography, palaeotectonics and palaeoenvironments as controls of changes in ammonite and brachiopod communities (an example from the early and middle Jurassic of western Algeria). *Palaeogeography, Palaeoclimatology, Palaeoecology* 47, 347–360.
- Elmi, S., Mangold, C., Mouterde, R. & Ruget, C. 1971: Révision de l'étage Bathonien au Cap Mondego (Portugal). *Annales Instituti Geologici Publici Hungarici* 54, 439–450.
- Enay, R. & Cariou, E. 1999: Jurassic ammonite faunas from Nepal and their bearing on the palaeobiogeography of the Himalayan belt. *Journal of Asian Earth Sciences* 17, 829–848.
- Enay, R. & Mangold, C. 1994: Première zonation par ammonites du Jurassique d'Arabie Saoudite, une référence pour la province arabique. *Geobios M.S.* 17, 161–174.
- Enay, R. & Mangold, C. 1996: Dimorphisme dans le genre *Ermoceras* Douvillé, 1916 (Stephanoceratidae, Bajocien supérieur) et implication nomenclaturale. *Comptes Rendus de l'Académie des Sciences de Paris* 322, 791–798.
- Enay, R., Le Nindre, C., Mangold, C., Manivit, J. & Vaslet, D. 1986: The Jurassic of Central Saudi Arabia: new data on lithostratigraphic units, paleoenvironments, ammonite fauna, ages and correlations. Technical Record, Bureau des Recherches Géologiques et Minières, 06-3, 67 pp. Deputy Ministry for mineral resources, Jiddah, Kingdom of Saudi Arabia.
- Enay, R., Le Nindre, C., Mangold, C., Manivit, J. & Vaslet, D. 1987a: Le Jurassique d'Arabie saoudite centrale: nouvelles données sur la lithostratigraphie, les paléoenvironnements, les faunes d'ammonites, les âges et les corrélations. *Geobios M.S.* 9, 13–65.
- Enay, R., Mangold, C., Du Dresnay, R. & Rakus, M. 1987b: Arrivals of Arabian origin among the ammonite faunas of Morocco during the Bajocian-Bathonian. *Palaeogeography, Palaeoclimatology, Palaeoecology* 61, 107–120.
- Enay, R., Gauthier, H., Trevisan, M., Berton, J.B., Brivet, L., Brodbeck, J.L., Demaizieres, J.F., Donie, P., Fourel, A. & Trehour, M. 2001: Les *Micromphalites* (Ammonitina) du Bathonien inférieur de la Nièvre (France): installation sur la marge européenne de la Téthys de formes sud-téthysiennes d'origine arabique et description d'un néotype de *M. busqueti* (de Gross.). *Revue de Paléobiologie* 20, 503–524.
- Erba, E. 1990a: Calcareous nannofossils from the Bas Auran section. In Rocha, R.B. & Soares, A.F. (eds.): *2nd International Symposium on Jurassic Stratigraphy 1987 (1988)*, 343–345. Centro de Estratigrafia e Paleobiologia, Univ. Nova de Lisboa, Lisbon.
- Erba, E. 1990b: Calcareous nannofossils biostratigraphy of some Bajocian sections from the Digne area (SE France). *Memorie descrittive della Carta Geologica d'Italia* 60, 237–256.
- Fauconnier, D. 1997: Kystes de dinoflagellés des domaines nord-ouest européen et sud-téthysien. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires* 17, 225–241.
- Fernández-López, S. 2000: Lower Bathonian ammonites of Serra de la Creu (Tivissa, Catalan Basin, Spain). *Revue de Paléobiologie* 8, 45–52.
- Fernández-López, S. 2001: Upper Bathonian ammonites of the Catalan Basin (Tivissa and Cap Salou, Spain). *Hantkeniana* 3, 25–39.
- Fernández-López, S.R. 2007a: Ammonoid taphonomy, palaeoenvironments and sequence stratigraphy at the Bajocian/Bathonian boundary on the Bas Auran area (Subalpine Basin, SE France). *Lethaia* 40, 377–391.
- Fernández-López, S.R. 2007b: Candidate sections for GSSP of the base of Bathonian Stage (Middle Jurassic). *Volumina Jurassica*, in litt.
- Fernández-López, S.R. & Henriques, M.H. 2002: Upper Bajocian – Lower Bathonian ammonites of Cabo Mondego section (Portugal). In Martire, L. (ed.): *6th International Symposium on the Jurassic System, Abstracts and Programs*, 65–66. International Subcommission on Jurassic Stratigraphy, Mondello.
- Fernández-López, S., Chong Díaz, G., Quinzio Sinn, L.A. & Wilke, H.-G. 1994: The Upper Bajocian and Bathonian in the Cordillera de Domeyko, North-Chilean Precordillera: sedimentological and biostratigraphical results. *Geobios M.S.* 17, 187–201.

- Fernández-López, S., Henriques, M.H.P. & Duarte, L.V. 2002: Taphonomy of ammonite condensed associations. Jurassic examples from carbonate platforms of Iberia. *Abhandlungen der Geologischen Bundesanstalt* 57, 423–430.
- Fernández-López, S.R., Henriques, M.H. & Mangold, C. 2006a: Ammonite succession at the Bajocian/Bathonian boundary in the Cabo Mondego region (Portugal). *Lethaia* 39, 253–264.
- Fernández-López, S.R., Henriques, M.H., Mangold, C. 2006b: Ammonite horizons at the basal Bathonian zone (Parvum Subzone) in Cabo Mondego, Portugal. *Volumina Jurassica* 4, 161.
- Fernández-López, S.R., Henriques, M.H., Mangold, C. & Pavia, G. 2007: New Early Bathonian Bigotitinae and Zigzagiceratinae (Ammonoidea, Middle Jurassic). *Rivista Italiana di Paleontologia e Stratigrafia* 113, 383–399.
- Ferry, S. 1990: Post-field comments. Mesozoic eustacy record on western Tethyan margins. Post-meting field-trip in the Vocontian Trough. *Publication of the Association des Sédimentologues français* 12, 121–140.
- Galácz, A. 1980: Bajocian and Bathonian Ammonites of Gyenespuszta Bakony Mts., Hungary. *Geologica Hungarica* 39, 1–228.
- Galácz, A. 1985: A Bathonian (Middle Jurassic) ammonite faunula from Monte Kumeta (Western Sicily). *Fragmenta Mineralogica et Palaeontologica* 12, 19–26.
- Galácz, A. 1990: New collection of successive ammonite faunas from the Bajocian of Mombasa (Kenya, East Africa). *Memoire Descrittive della Carta Geologica d'Italia* 40, 199–204.
- Galácz, A. 1993: Comparison of Hungarian Bathonian (Middle Jurassic) ammonite assemblages; suggestions on habitats. *Discussiones Palaeontologicae* 39, 25–33.
- Galácz, A. 1994: The age of the ammonite fauna from the classic Middle Jurassic locality of Swinitza (Banat, Romania). *Palaeopelagos S.P.* 1, 167–179.
- Galácz, A. 1995a: Ammonite stratigraphy of the Bathonian red limestone of the Mecsek Mts, south Hungary. *Annales Universitatis Scientiarum Budapestinensis* 30 (1994), 111–150.
- Galácz, A. 1995b: Revision of the Middle Jurassic ammonite fauna from Csóka-hegy, Vértes Hills (Transdanubian Hungary). *Hantkeniana* 1, 119–129.
- Galácz, A. 1999a: A Lower Bathonian ammonite fauna from Erice (Western Sicily). *Annales Universitatis Scientiarum Budapestinensis* 32, 149–168.
- Galácz, A. 1999b: Middle Jurassic ammonites of Arabian origin in Western Sicily. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 10, 605–613.
- Garnier, A. 1872: Réunion extraordinaire à Digne. *Bulletin de la Société Géologique de France* 29, 626–656.
- Géczy, B. & Galácz, A. 1998: Bathonian ammonites from the classic Middle Jurassic locality of Villány, South Hungary. *Revue de Paléobiologie* 17, 479–511.
- Goguel, J. 1966: Carte géologique à l'échelle 1:80000, feuille Castellane. BRGM, Paris.
- Graciansky, P.-C. de, Durozoy, G. & Gigot, P. 1982: Carte géologique de la France à 1:50000, Digne. 75 pp. BRGM, Paris.
- Graciansky, P.-C. de, Dardeau, G., Dumont, T., Jacquin, T., Marchand, D., Mouterde, R. & Vail, P.R. 1993: Depositional sequence cycles, transgressive-regressive facies cycles, and extensional tectonics, example from the southern subalpine Jurassic basin, France. *Bulletin de la Société géologique de France* 164, 709–718.
- Graciansky, P.-C. de, Dardeau, G., Dommergues, J.L., Dyrlet, C., Marchand, D., Dumont, T., Hesselbo, S.P., Jacquin, T., Goggin, V., Meister, C., Mouterde, R., Rey, J. & Vail, P.R. 1998a: Ammonite biostratigraphic correlation and Early Jurassic sequence stratigraphy in France: Comparisons with some U.K. sections. *SEPM Special Publication No. 60*, 583–622.
- Graciansky, P.-C. de, Dardeau, G., Bodeur, Y., Elmi, S., Fortwengler, D., Jacquin, T., Marchand, D. & Thierry, J. 1998b: Les Terres Noires du Sud-Est de la France (Jurassique moyen et supérieur): interprétation en termes de stratigraphie séquentielle. *Bulletin du Centre de Recherches Elf Exploration-Production* 22, 35–66.
- Gradstein, F.M. & Ogg, J.G. 2004: Geologic Time Scale 2004 – why, how and where next! *Lethaia* 37, 175–181.
- Gradstein, F.M., Finney, S.C., Lane, R. & Ogg, J.G. 2003: ICS on stage. *Lethaia* 36, 371–378.
- Gradstein, F.M., Ogg, J.G. & Smith, A.G. (eds.) 2005: *A Geologic Time Scale 2004*. 589 pp. Cambridge University Press, Cambridge.

- Gräfe, K.-U. 2005: Benthic foraminifers and palaeoenvironment in the Lower and Middle Jurassic of the Western Basque-Cantabrian Basin (Northern Spain). *Journal of Iberian Geology* 31, 217–233.
- Gröschke, M. 1996: Zwei neue Ammoniten der Gattung *Iniskinites* aus dem Bathonium (Mitteljura) von Nordchile. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1996, 140–146.
- Gröschke, M. & Hillebrandt, A. von 1994: The Bathonian in Northern Chile. *Geobios M.S.* 17, 255–264.
- Guillaume, L. 1938: Observations sur la limite inférieure des marnes à *Posinomya alpina* entre Digne et Castellane. *Compte Rendu Sommaire des Séances de la Société Géologique de France*, 189–199.
- Hall, R.L. 1984: Lithostratigraphy and biostratigraphy of the Fernie Formation (Jurassic) in the southern Canadian Rocky Mountains. *Canadian Society of Petroleum Geologists Memoir* 9, 233–247.
- Hall, R.L. 1988: Late Bajocian and Bathonian (Middle Jurassic) ammonites from the Fernie Formation, Canadian Rocky Mountains. *Journal of Paleontology* 62, 575–586.
- Hall, R.L. 1989: New Bathonian (Middle Jurassic) ammonite faunas from the Fernie Formation, southern Alberta. *Canadian Journal of Earth Sciences* 26, 16–22.
- Hall, R.L. & Stronach, N.J. 1981: First record of late Bajocian (Jurassic) ammonites in the Fernie Formation, Alberta. *Canadian Journal of Earth Sciences* 18, 919–925.
- Hall, R. & Westermann, G.E.G. 1980: Lower Bajocian (Jurassic) cephalopod faunas from Western Canada and proposed assemblage zones for the Lower Bajocian of North America. *Palaeontographica Americana* 9, 1–93.
- Hallam, A. 1994: *An outline of Phanerozoic Biogeography*. 246 pp. Oxford University Press, Oxford.
- Halloy J.J. d'O. d' 1843: *Précis élémentaire de géologie*. 790 pp. A. Bertrand Ed., Paris.
- Hardenbol, J., Thierry, J., Farley, M.B., Jacquin, T., Graciansky, P.-C. de & Vail, P.R. 1998: Mesozoic and Cenozoic sequence chronostratigraphic framework of European Basins. In Graciansky, P.C. de, Hardenbol, J., Jacquin, T. & Vail, P.R. (eds.): *Mesozoic and Cenozoic Sequence Stratigraphy of European Basins*, 3–14. SEPM (Society for Sedimentary Geology) Special Publication, No. 60, Tulsa, Oklahoma.
- Harland, W.B., Cox, A.V., Llewellyn, P.G., Pickton, C.A.G., Smith, A.G. & Walters, R. 1982: A Geologic Time Scale. 131 pp. Cambridge University Press, Cambridge.
- Haug, E. 1891: Les Chaînes Subalpines entre Gap et Digne. *Bulletin du Service de la Carte Géologique de France* 3, 21, 1–192.
- Henriques, M. & Ramalho, M.M. 2005: Jurassic Heritage of Cabo Mondego (Central Portugal). In Henriques, M.H. (Coord.): *Jurassic Heritage and Geoconservation in Portugal: Selected Sites*. Field Trip Guide Book. IV International Symposium ProGEO on the Conservation of the Geological Heritage, 37–43. Geosciences Centre, Coimbra.
- Hillebrandt, A. von 1995: A Late Bathonian morphoceratid (Jurassic, Ammonitina) from Peru. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 35, 27–37.
- Hillebrandt, A. von 2001. Ammonite stratigraphy of the Bajocian in Northern Chile. *Hantkeniana* 3, 49–87.
- Hillebrandt, A. von, Smith, P., Westermann, G.E.G. & Callomon, J.H. 1992a: Ammonites zones of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 247–272. Cambridge University Press, Cambridge.
- Hillebrandt, A. von, Westermann, G.E.G., Callomon, J.H., Detterman, R.L., 1992b: Ammonites of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 342–359. Cambridge University Press, Cambridge.
- Imlay, R.W. 1980: Middle Jurassic (Bathonian) ammonites from Southern Alaska. *U. S. Geological Survey Professional Paper* 1091, 1–42.
- Imlay, R.W. 1981: Jurassic (Bathonian and Callovian) ammonites in Eastern Oregon and Western Idaho. *U. S. Geological Survey Professional Paper* 1142, 1–24.
- Imlay, R.W. 1982: Late Bajocian ammonites from Southern Alaska. *U. S. Geological Survey Professional Paper* 1189, 1–21.

- Imlay, R.W. 1984: Jurassic ammonite successions in North America and biogeographic implications. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 1–12. Cambridge University Press, Cambridge.
- Innocenti, M., Mangold, C., Pavia, G. & Torrens, H.S. 1990: A proposal for the formal ratification of the basal boundary stratotype of the Bathonian Stage based on a Bas Auran section (S.E. France). In Rocha, R.B. & Soares, A.F. (eds.): *2nd International Symposium on Jurassic Stratigraphy 1987 (1988)*, 333–346. Centro de Estratigrafia e Paleobiologia, Universidade Nova de Lisboa, Lisbon.
- Jacquin, T., Dardeau, G., Durllet, C., Graciansky, P.C. de & Hantzpergue, P. 1998: The North Sea Cycle: an overview of 2nd-order transgressive/regressive facies cycles in Western Europe. In Graciansky, P.C. de, Hardenbol, J., Jacquin, T. & Vail, P.R. (eds.): *Mesozoic and Cenozoic Sequence Stratigraphy of European Basins*, 445–466. SEPM (Society for Sedimentary Geology) Special Publication, No. 60, Tulsa, Oklahoma.
- Jana, B.N. & Hilton, J. 2007: Resolving the age of the Mesozoic Kuar Bet Beds (Kachchh, Gujarat, India): A reinvestigation of palaeobotanical and palynological assemblages. *Journal of Asian Earth Sciences* 30, 457–463.
- Jaitly, A.K. & Singh, C.S.P. 1983: Discovery of the Late Bajocian-*Leptosphinctes* BUCKMAN (Jurassic Ammonitina) from Kachchh, Western India. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1983, 91–96.
- Jaitly, A.K. & Singh, C.S.P. 1984: On the Bathonian (Middle Jurassic) ammonites *Micromphalites* Buckman and *Gracilisphinctes* Buckman from Kachchh, Western India. *Geological Magazine* 121, 319–321.
- Jenkyns, H., Jones, C., Gröcke, D., Hesselbo, S. & Parkinson, N. 2002: Chemostratigraphy of the Jurassic System: applications, limitations and implications for palaeoceanography. *Journal of the Geological Society, London* 159, 351–378.
- Joly, B. 2000: Les Juraphyllitidae, Phylloceratidae, Neophylloceratidae (Phyllocerataceae, Phylloceratina, Ammonoidea) de France au Jurassique et au Cretace. *Geobios M. S.* 23, 1–204.
- Kaenel, E. de & Bergen, J.A., 1993: New Early and Middle Jurassic coccolith taxa and biostratigraphy from the eastern proto-Atlantic (Morocco, Portugal and DSDP Site 547B). *Eclogae Geologicae Helvetiae* 86, 861–907.
- Kaenel, E. de, Bergen, J.A. & Salis Perch-Nielsen, K. von 1996: Jurassic calcareous nannofossil biostratigraphy of Western Europe. Compilation of recent studies and calibration of bioevents. *Bulletin de la Société Géologique de France* 167, 15–28.
- Kimura, T., Lebedev, E.L., Markovich, E.M. & Samylina, V.A. 1992: Macroflora of eastern Asia and other circum-Pacific areas. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 311–316. Cambridge University Press, Cambridge.
- Köstler, L. & Schairer, G. 1996: *Morphoceras* aus dem “Parkinsonien-Oolith” (Mittlerer Jura) von Segenthal/Opf. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 36, 81–85.
- Kopik, J. 2006: Bathonian ammonites of the families Sphaeroceratidae BUCKMAN and Tullitidae BUCKMAN from the Polish Jura Chain (Southern Poland). *Polish Geological Institute, Special Papers* 21, 1–68.
- Krystyn, L. 1972: Die Oberbajocium- und Bathonium-Ammoniten der Klaus-Schichten des Steinbruches Neumühle bei Wien (Österreich). *Annalen des Naturhistorischen Museum in Wien* 76, 195–310.
- Lemoine, M. 1984: Mesozoic evolution of the western Alps. *Annales Geophysicae* 2, 171–172.
- Lemoine, M. 1985: Structuration jurassique des Alpes occidentales et palinspastiques de la Téthys ligure. *Bulletin de la Société Géologique de France* 8, 126–137.
- Luczynski, P., Matyja, B. A. & Wierzbowski, A. 2000: Ontogeny and ecological interpretation of the ammonites of the *Asphinctites-Polysphinctites* group (Tenuiplicatus Zone, lower Bathonian). *Bajocian-Bathonian Working Groups Meeting*, 23–24. Budapest.
- Majidifard, M.R. 2003: Biostratigraphy, Lithostratigraphy, ammonite taxonomy and microfacies analysis of the Middle and Upper Jurassic of northeastern Iran. Dissertation zur Erlangung des Naturwissenschaftlichen Doktorgrades der Bayerischen Julius-Maximilians-Universität Würzburg, 201 pp.

- Manceñido, M.O. & Dagys, A.S. 1992: Brachiopods of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 328–333. Cambridge University Press, Cambridge.
- Mangold, C. 1971a: Stratigraphie des étages Bathonien et Callovien du Jura Méridional. *Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon* 41, 1, (1970), 1–376.
- Mangold, C. 1971b: Les Perisphinctidae (Ammonitina) du Jura Méridional au Bathonien et au Callovien. *Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon* 41 (1970), 2, 1–246.
- Mangold, C. 1971c: Morphoceratidae (Ammonoidea, Perisphinctidae) bathoniens du Jura méridional, de la Nièvre et du Portugal. *Geobios* 3 (1970), 43–130.
- Mangold, C. 1981: Le Bathonien de l'Est du Subbétique (Espagne du Sud). *Cuadernos de Geología* 10 (1979), 271–281.
- Mangold, C. 1984: Report of the Bathonian Working Group. In Michelsen O. & Zeiss A. (eds.): *1st International Symposium on Jurassic Stratigraphy (IUGS)*, 67–75. Geological Survey of Denmark, Copenhagen.
- Mangold, C. 1990: Le Bathonien du Cap Mondego (N de Figueira da Foz, Portugal). Biochronologie et corrélations. *Cahiers de l'Université Catholique de Lyon* 4, 89–105.
- Mangold, C. 1997a: Tendances évolutives chez les Morphocératidés (Périsphinctacés, Ammonitina). *Cahiers de l'Université Catholique de Lyon* 10, 93–101.
- Mangold, C. 1997b: Le Jurassique moyen. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires* 17, 355–361.
- Mangold C. 1999. Report of the Bajocian-Bathonian boundary Working Group. *International Subcommission on Jurassic Stratigraphy, Newsletter* 26, 50–52.
- Mangold, C. & Gygi, R. 1997. Bathonien ammonites from Canton Aargau, Northern Switzerland: stratigraphy, taxonomy and biogeography. *Geobios* 30, 497–518.
- Mangold, C. & Rioult, M. 1997: Bathonien. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires* 17, 55–62.
- Martire, L. 1989: *Analisi biostratigrafica e sedimentologica del rosso ammonitico veronese dell'Altopiano di Asiago*. 166 pp. Thesis Univ. Torino, unpublished.
- Martire, L. & Pavia, G. 2004: Jurassic sedimentary and tectonic processes at Montagna Grande (Trapanese Domain, Western Sicily, Italy). *Rivista Italiana di Paleontologia e Stratigrafia* 110, 23–33.
- Mattioli, E. & Erba, E. 1999: Synthesis of calcareous nannofossil events in the Tethyan Lower and Middle Jurassic. *Rivista Italiana di Paleontologia e Stratigrafia* 105, 343–376.
- Matyja, B.A. & Wierzbowski, A. 2000: Ammonites and stratigraphy of the uppermost Bajocian and Lower Bathonian between Czestochowa and Wielun, Central Poland. *Acta Geologica Polonica* 50, 191–209.
- Matyja, B.A. & Wierzbowski, A. 2001: Palaeogeographical distribution of early Bathonian ammonites of the *Asphinctites-Polysphinctites* group. *Hantkeniana* 3, 89–103.
- Maubeuge, P.L. 1950: *Sur le Bathonien et en particulier sur le Bathonien lorrain (note préliminaire)*. 16 pp. Impr. Thomas, Nancy.
- Metz, M. 1990: Ein neues Bajocium-Profil (Mittlerer Jura) im Osnabrücker Bergland (Niedersachsen). *Osnabrücker naturwissenschaftliche Mitteilungen* 16, 7–30.
- Metz, M. 1992: Die Faunenhorizonte der “Subfurcaten-Schichten” (Bajocium, Niortense-Zone) in Nordwestdeutschland. *Osnabrücker naturwissenschaftliche Mitteilungen* 18, 25–65.
- Mitta, V.V. 2001: Distribution of the Bajocian-Bathonian ammonites in the South-West chains of Hissar Range. *Hantkeniana* 3, 105–129.
- Mitta, V.V. 2004: *Sokurella galaczi* gen. et sp. nov. and other Middle Jurassic Parkinsoniidae (Ammonoidea) from the lower reaches of the Volga River. *Paleontological Journal* 38, 257–264.
- Mitta, V.V. 2005: Late Bathonian Cardioceratidae (Ammonoidea) from the Middle Reaches of the Volga River. *Paleontological Journal* 39, 629–644.
- Mitta, V.V. 2006: The First discovery of *Arctocephalites* (Cardioceratidae, Ammonoidea) from the Middle Jurassic of the Pechora Basin (Northern Russia) [in Russian]. In Barskov, I. &

- Leonova, T. (eds.): *Contributions to current cephalopod research*, 82–84. Paleontological Institute, Moscow.
- Mitta, V.V. 2007: The Boundary of the Bajocian and Bathonian stages in the European Russia [in Russian]. In Zakharov, V.A. (ed.): *Jurassic System of Russia: problems of Stratigraphy and Paleogeography. Second all-Russian meeting*, 161–163. Yaroslavl State Pedagogical University, Yaroslavl.
- Mitta, V.V. & Seltzer, V.B. 2002: First finds of Arctocephalitinae (Ammonoidea) in the Jurassic of the south-eastern Russian Platform, and the correlation of the Boreal Bathonian Stage with the standard scale [in Russian]. *Transactions of the Scientific Research Geological Institute of the N.G. Chernyshevskii Saratov State University* 10, 12–39.
- Mitta, V.V., Barskov, I.S., Gründel, J., Zakharov, V.A., Seltzer, V.B., Ivanov, A.V., Rostovtseva, J.A. & Tarasova, L.O. 2004: The Upper Bajocian and Lower Bathonian in the section near Saratov [in Russian]. *Vernadsky Museum Novitates* 2, 1–39.
- Morton, N. 1974: The definition of standard Jurassic Stages. *Mémoire du Bureau de Recherches géologiques et minières* 75 (1971), 83–93.
- Morton, N. 2006: Chronostratigraphic units in the Jurassic and their boundaries: Definition, recognition and correlation, causal mechanisms. *Progress in Natural Science* 16, 13, 1–11.
- Moyne, S., Thierry, J., Marchand, D., Nicolleau, P., Pineau, J.P., Courville, P. & Saucède, T. 2005: Le genre *Nucleolites* (Echinoidea, Cassiduloidea) du Bajocien à l'Oxfordien dans le Bassin de Paris: apport des données architecturales à la systématique et à la phylogénie. *Geobios* 38, 519–532.
- O'Dogherty, L., Sandoval, J., Bartolini, A., Bruchez, S., Bill, M. & Guex, J. 2006: Carbon–isotope stratigraphy and ammonite faunal turnover for the Middle Jurassic in the Southern Iberian palaeomargin. *Palaeogeography, Palaeoclimatology, Palaeoecology* 239, 311–333.
- Ogg, J.G. 2004: Status of Divisions of the International Geologic Time Scale. *Lethaia* 37, 183–199.
- Ohmert, W., Gassmann, G., Kutz, M., Mattes, R. & Müller, R. 2004: Neue Ammonitenfunde aus der Hauptrogenstein-Formation (Mitteljura) des Breisgau (Oberrhein). *Jahresberichte und Mitteilungen des Oberrheinischen Geologischen Vereines* 86, 337–350.
- Olivero, D. 1994: La trace fossile *Zoophycos* du Jurassique du Sud-Est de la France. Signification paléoenvironnementale. *Documents des Laboratoires de Géologie Lyon* 129, 1–329.
- Olivero, D. 2003: Early Jurassic to Late Cretaceous evolution of *Zoophycos* in the French Subalpine Basin (southeastern France). *Palaeogeography, Palaeoclimatology, Palaeoecology* 192, 59–78.
- Olivero, D. & Atrops, F. 1996: Les séries à *Zoophycos* du Bathonien-Callovien de l'Arc de Castellane (SE de la France) dans la zone de transition plate-forme/bassin: stratigraphie et paléotectonique. *Comptes Rendus de l'Académie des Sciences de Paris* 323, 81–88.
- Olivero, D., Mangold, C. & Pavia, G. 1997: La formation des Calcaires à *Zoophycos* du Verdon (Bathonien inférieur à Callovien moyen) des environs de Castellane (Alpes-de-Haute-Provence, France): biochronologie et lacunes. *Comptes Rendus de l'Académie des Sciences de Paris* 324, 33–40.
- Oppel, A. 1856–1858: *Die Juraformation Englands, Frankreichs und des südwestlichen Deutschlands, nach ihren einzelnen Gliedern eingeteilt und verglichen*. 857 pp. Ebner & Seubert, Stuttgart.
- Oppel, A. 1862: Über jurassische Cephalopoden. *Palaeontologische Mittheilungen aus dem Museum des Königl. Bayer. Staates* 3, 127–266.
- Oppel, A. 1865: Geognostische Studien in den Ardèche Departement. *Palaeontologische Mittheilungen aus dem Museum des Königl. Bayer. Staates* 5, 305–322.
- Orbigny, A. d' 1842–1851: *Paléontologie française. Terrain Jurassique. 1. Céphalopodes*. 641 pp. Paris.
- Orbigny, A. d' 1849–1852: *Cours élémentaire de paléontologie et de géologie stratigraphiques*. 1145 pp. Masson Ed., Paris.
- Ouahhabi, B. 1994: *Le Lias et le Dogger inférieur des Beni Snassen orientaux (Maroc). Stratigraphie, paléontologie et dynamique du bassin. Comparaison avec les Monts d'Oujda et avec les régions limitrophes*. 496 pp. Thesis Univ. Oujda.
- Page, K.N. 1996a: Mesozoic Ammonoids in space and time. In Landmann, N.H., Tanabe, K. and R.A. Davis (eds.): *Ammonoid Paleobiology. Topics in Geobiology* 13, 755–794.

- Page, K.N. 1996b: Observations on the succession of stratigraphically useful ammonite faunas in the Bathonian (Middle Jurassic) of south-west England, and their correlation with a Sub-Mediterranean "Standard Zonation". *Proceedings of the Ussher Society* 9, 45–53.
- Page, K.N. 2001: Up a Bathonian backwater – a review of the ammonite evidence for correlating sequences with interdigitating non-marine facies in central and northern England. *Hantkeniana* 3, 131–148.
- Page, K.N., Meléndez, G. & Henriques, M.H. 2006: Jurassic Global Stratotype Section and Points (GSSPs) – a potential serial World Heritage Site? *Volumina Jurassica* 7, 253.
- Pandey, D.K. & Agrawal, S.K. 1984: On two new species of the Middle Jurassic ammonite genus *Clydoniceras* Blake from Kachchh, western India. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1984, 321–326.
- Pandey, D.K. & Callomon, J.H. 1995: Contributions to the Jurassic of Kachchh, Western India. III. The Middle Bathonian ammonite families *Clydoniceratidae* and *Perisphinctidae* from Pachchham Island. *Beringeria* 16, 125–145.
- Pandey, D.K. & Westermann, G.E.G. 1988: First record of Bathonian *Bullatimorphites* (Jurassic, Ammonitina) from Kachchh, India. *Journal of Paleontology* 62, 148–150.
- Parkinson, D.N. 1996: Gamma-ray spectrometry as a tool for stratigraphical interpretation: examples from the western European Lower Jurassic. *Geological Society, London, Special Publications* 103, 231–255.
- Parent, H. 1998: Upper Bathonian and Lower Callovian ammonites from Chacay Melehué (Argentina). *Acta Paleontologica Polonica* 42, 69–130.
- Parnes, A. 1981: Biostratigraphy of the Mahmal Formation (Middle and Upper Bajocian) in Makhtesh Ramon (Negev, Southern Israel). *Geological Survey of Israel, Bulletin*. 74, 1–55.
- Parnes, A. 1984: Irregularly coiled Stephanoceratidae from the Upper Bajocian of Gebel Maghara, Sinai, Egypt. *Geological Survey of Israel, Current Research 1983-84*, 29–32.
- Parnes, A. 1988: Middle Jurassic (Middle Bajocian - Middle Bathonian) ammonites from Gebel Maghara, Sinai, Egypt. In the collections of the Geological Survey of Israel. *Journal of African Earth Sciences* 7, 707–733.
- Patrului, D. 1996: Ammonites hétéromorphes et autres Parkinsoniidés du Bathonien-Callovien inférieur de Vadu Crisului (Monts Apuseni - Roumanie). *Mémoires Institut de Géologie et de Géophysique* 36, 13–19.
- Pavia, G. 1973: Ammoniti del Baiociano superiore di Digne (Francia SE, Dip. Basses-Alpes). *Bollettino della Società Paleontologica Italiana* 10, 2 (1971), 75–142.
- Pavia, G. 1983a: Il genere *Ptychophylloceras* Spath, 1927 (Ammonoidea, Phyllocerataceae) nel Baiociano sudeuropeo. *Atti della Accademia Nazionale dei Lincei, Memorie* 1982, 17, 1–31.
- Pavia, G. 1983b: New data on *Orthogarantiana* (Torrensia) Sturani, 1971 (Ammonitina, Stephanocerataceae) in the European Upper Bajocian. *Bollettino del Museo Regionale di Scienze Naturali di Torino* 1, 201–214.
- Pavia, G. 1984: Bajocien et Bathonien de l'Arc de Digne. In Debrand-Passard, S. (ed.). Synthèse géologique du Sud-Est de la France. *Mémoire B.R.G.M.* 125, 199–200.
- Pavia, G. 1994: Taphonomic remarks on d'Orbigny's type-Bajocian (Bayeux, west France). *Servizio Geologico Nazionale, Miscellanea* 5, 93–111.
- Pavia, G. 2000: New *Subcollina* (Ammonitida) from the topmost Lower Bajocian: their phylogenetic and palaeogeographic significance. *GeoResearch Forum* 6, 397–406.
- Pavia, G. 2007: *Lissoceras monachum* (Gemmellaro), a ghost Ammonitida of the Tethyan Bathonian. *Bollettino della Società Paleontologica Italiana* 45 (2006), 217–226.
- Pavia, G. & Cresta, S. (coords.) 2002: Revision of Jurassic ammonites of the Gemmellaro collections. *Quaderni del Museo Geologico "G.G. Gemmellaro"* 6, 1–408.
- Pavia, G. & Sturani, C. 1968: Étude biostratigraphique du Bajocien des Chaînes Subalpines aus environs de Digne (Basses-Alpes). *Bollettino della Società Geologica Italiana* 87, 305–316.
- Pavia, G., Martire, L., Canzoneri, V. & D'Arpa, C. 2002: Rocca chi Parra quarry, a condensed Rosso Ammonitico succession: depositional and erosional geometries, neptunian dykes and ammonite assemblages. In Santantonio, M. (ed.): *General Field Trip Guidebook, VI International Symposium on the Jurassic System*, 42–48. Palermo.

- Pavia, G., Fernández-López, S.R. & Mangold, C. 2008: Ammonoid succession at the Bajocian-Bathonian transition in the Bas Auran area, Digne district, South-East France. *Rivista Italiana di Paleontologia e Stratigrafia*, in litt.
- Pawellek, T. & Aigner, T. 2003: Stratigraphic architecture and gamma ray logs of deeper ramp carbonates (Upper Jurassic, SW Germany). *Sedimentary Geology* 159, 203–240.
- Pawellek, T. & Aigner, T. 2004: Dynamic stratigraphy as a tool in economic mineral exploration: ultra-pure limestones (Upper Jurassic, SW Germany). *Marine and Petroleum Geology* 21, 499–516.
- Pellenard, P., Deconinck, J.-F., Huff, W.D., Thierry, J., Marchand, D., Fortwengler, D. & Trouiller, A. 2003: Characterization and correlation of Upper Jurassic (Oxfordian) bentonite deposits in the Paris Basin and the Subalpine Basin, France. *Sedimentology* 50, 1035–1060.
- Pessagno, E. Jr. & Mizutani, S. 1992: Radiolarian biozones of North America and Japan. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 293–295. Cambridge University Press, Cambridge.
- Poulsen, N.E. 1997: Report on samples from Bas Auran, SE France. *Jurassic Microfossil Group Newsletter* 6, 9–10.
- Poulsen, N.E. & Riding, J.B. 2003: The Jurassic dinoflagellate cyst zonation of Subboreal Northwest Europe. *Geological Survey of Denmark and Greenland Bulletin* 1, 115–144.
- Poulton, T.P. 1987: Zonation and correlation of Middle Boreal Bathonian to lower Callovian (Jurassic) ammonites, Salmon Cache Canyon, Porcupine River, northern Yukon. *Geological Survey of Canada, Bulletin* 358, 1–155.
- Poulton, T.P., Hall, R.L., Tipper, H.W., Cameron, B.E.B. & Carter, E.S. 1991: Current status of Middle Jurassic biostratigraphy of the Queen Charlotte Islands, British Columbia. *Geological Survey of Canada, Paper* 90-10, 237–252.
- Poulton, T.P., Hall, R.L. & Callomon, J.H. 1994: Ammonite and bivalve assemblages in Bathonian through Oxfordian strata of northern Bowser Basin, northwestern British Columbia, Canada. *Geobios M.S.* 17, 415–421.
- Raddadi, M.C., Vanneau, A.A., Poupeau, G., Carrio-Schaffhauser, E., Arnaud, H. & Rivera, A. 2005: Interpretation of gamma-ray logs: The distribution of uranium in carbonate platform. *Comptes Rendus Geosciences* 337, 1457–1461.
- Rawson, P.F. 1982: New Arctocephalitinae (Ammonoidea) from the Middle Jurassic of Kong Karls Land, Svalbard. *Geological Magazine* 119, 95–100.
- Remane, J., Bassett, M.G., Cowie, J.W., Gorbbrandt, K.H., Lane, H.R., Milchelson, O. & Wang, N. 1996: Revised guidelines for the establishment of global chronostratigraphic standards by the International Commission of Stratigraphy (ICS). *Episodes* 19, 77–81.
- Riccardi, A.C. 1985: Los Eurycephalitinae andinos (Ammonitina, Jurásico Medio): modelos evolutivos y resolución paleontológica. *Boletín Gent. Inst. Fitolec. Castelar* 13, 1–27.
- Riccardi, A.C. 1991: Jurassic and Cretaceous marine connections between the Southeast Pacific and Tethys. *Palaeogeography, Palaeoclimatology, Palaeoecology* 87, 155–189.
- Riccardi, A.C. & Westermann, G.E.G. 1991a: Middle Jurassic ammonite fauna and biochronology of the Argentine-Chilean Andes. Part III: Bathonian-Callovian Eurycephalitinae, Stephanocerataceae. *Palaeontographica* 216, 1–110.
- Riccardi, A.C. & Westermann, G.E.G. 1991b: Middle Jurassic ammonite fauna and biochronology of the Argentine-Chilean Andes. Part IV: Bathonian-Callovian Reineckeidae. *Palaeontographica* 216, 111–145.
- Riccardi, A.C. & Westermann, G.E.G. 1999: An early Bathonian Tethyan ammonite fauna from Argentina. *Palaeontology* 42, 193–209.
- Riccardi, A.C., Westermann, G.E.G. & Elmi, S. 1990a: The Middle Jurassic Bathonian-Callovian Ammonite Zones of the Argentine-Chilean Andes. *Geobios* 22 (1989), 553–597.
- Riccardi, A.C., Westermann, G.E.G. & Damborenea, S.E. 1990b: Middle Jurassic of South America and Antarctic Peninsula. *Newsletter on Stratigraphy* 21, 105–128.
- Riccardi, A.C., Westermann, G.E.G. & Elmi, S. 1991: Biostratigraphy of the upper Bajocian-middle Callovian (Middle Jurassic), South America. *Journal of South American Earth Sciences* 4, 149–157.

- Riccardi, A.C., Damborenea, S.E., Manceñido, M.O. & Ballent, S.C. 1994: Middle Jurassic biostratigraphy of Argentina. *Geobios M.S.* 17, 423–430.
- Riding, J.B. & Thomas, J.E. 1992: Dinoflagellate cysts of the Jurassic System. In Powell A.J. (ed.): *A stratigraphic index of dinoflagellate cysts*, 7–97. Chapman & Hall, London.
- Rioult, M. 1964: Le stratotype du Bajocien. *Colloque du Jurassique Luxembourg 1962, Comptes Rendus et Mémoires*, 239–258. Institut Grand-Ducal, Luxembourg.
- Rioult, M., Contini, D., Elmi, S. & Gabilly, J. 1997: Bajocien. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires* 17, 41–53.
- Rocha, R., Manuppella, G., Mouterde, R., Ruget, C. & Zbyszewski, G. 1981: Carta geológica de Portugal, 1/50000, folha 19-C, Figueira da Foz. Serviços Geológicos de Portugal, Lisbon.
- Rocha, R.B., Mouterde, R., Soares, A.F. & Elmi, S. 1987: *Excursion A - Biostratigraphie et évolution séquentielle du Bassin au Nord du Tage au cours du Lias et du Dogger*. 84 pp. International Subcommission on Jurassic Stratigraphy, Centro de Estratigrafia e Paleobiologia da UNL (INIC), Lisbon.
- Rostovtsev, K.O. 1985. *The Jurassic deposits of the south part of the Transcaucasus* [in Russian]. 188 pp. Nauka, Leningrad.
- Roy, P., Bardhan, S., Mitra, A. & Jana, S.K. 2007: New Bathonian (Middle Jurassic) ammonite assemblages from Kutch, India. *Journal of Asian Earth Sciences* 30, 629–651.
- Ruban, D.A. 2006: Taxonomic diversity dynamics of the Jurassic bivalves in the Caucasus: Regional trends and recognition of global patterns. *Palaeogeography, Palaeoclimatology, Palaeoecology* 239, 63–74.
- Ruf, M., Link, E., Pross, J. & Aigner, T. 2005: Integrated sequence stratigraphy: Facies, stable isotope and palynofacies analysis in a deeper epicontinental carbonate ramp (Late Jurassic, SW Germany). *Sedimentary Geology* 175, 391–414.
- Ruget-Perrot, C. 1961: Études stratigraphiques sur le Dogger et le Malm inférieur du Portugal au Nord du Tage. *Serviços Geológicos de Portugal, Memória* 7, 1–197.
- Ruget, C. & Nicollin, J.P. 1997: Les petits foraminifères benthoniques dégagés. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires* 17, 281–291.
- Rulleau, L. 2006: *Biostratigraphie et paléontologie du Lias supérieur et du Dogger de la région lyonnaise*. 382 pp. Lafarge Ciments, Lozanne.
- Saltykov, V.F. 2007: The sequence of the ammonite and foraminifer zones in Bajocian – Bathonian section of the Lower Volga area [in Russian]. In Zakharov, V.A. (ed.): *Jurassic System of Russia: problems of Stratigraphy and Paleogeography. Second all-Russian meeting*, 201–203. Yaroslavl State Pedagogical University, Yaroslavl.
- Sandoval, J. 1983: *Bioestratigrafía y paleontología (Stephanocerataceae y Perisphinctaceae) del Bajocense y Bathonense en las Cordilleras Béticas*. 613 pp. Tesis Doctoral, Universidad de Granada, Granada.
- Sandoval, J. 1986: Middle Jurassic Haploceratidae (Ammonitina) from the Subbetic Zone (South Spain). *Geobios* 19, 435–463.
- Sandoval, J. 1990: A revision of the Bajocian divisions in the Subbetic Domain (southern Spain). *Memoire descrittive della Carta Geologica d'Italia* 40, 141–162.
- Sandoval, J. 1994: The Bajocian Stage in the Island of Majorca: biostratigraphy and ammonite assemblages. *Servizio Geologico Nazionale, Miscellanea* 5, 293–215.
- Sandoval, J. & Westermann, G.E.G. 1986: The Bajocian ammonite fauna of Oaxaca. *Journal of Paleontology* 60, 1220–1271.
- Sandoval, J., Westermann, G.E.G. & Marshall, M.C. 1990: Ammonite fauna, stratigraphy and ecology of the Bathonian-Callovian (Jurassic) Tecocoyunca Group, South Mexico. *Palaeontographica* 210, 93–149.
- Sandoval, J., O'Dogherty, L. & Guex, J. 2001: Evolutionary rates of Jurassic ammonites in relation to sea-level fluctuations. *Palaos* 16, 311–335.
- Sarjeant, W.A.S., Volkheimer, W. & Zhang, W.P. 1992: Jurassic Palynomorphs of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 273–292. Cambridge University Press, Cambridge.
- Sato, T. 1992: Southeast Asia and Japan. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 194–213. Cambridge University Press, Cambridge.

- Schairer, G. 1987: Ammoniten aus Bajoc und Bathon (mittlerer Jura) von Segenthal. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 27, 31–50.
- Schairer, G. 1994: *Polysphinctites polysphinctus* Buckman aus dem “Parkinsonien-Oolith” von Segenthal. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 34, 159–162.
- Schlögl, J. & Rakús, M. 2004: Ammonites of Arabian origin from the Early Bathonian of the Czorsztyn Unit, Pieniny Klippen Belt (Western Carpathians, Slovakia). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 2004, 449–460.
- Schlögl, J., Rakús, M., Mangold, C. & Elmi, S. 2005: Bajocian – Bathonian ammonite fauna of the Czorsztyn Unit, Pieniny Klippen Belt (Western Carpathians, Slovakia); its biostratigraphical and palaeobiogeographical significance. *Acta Geologica Polonica* 55, 339–359.
- Schlögl, J., Elmi, S., Rakús, M., Mangold, C. & Ouahhabi, M. 2006: Specialization and iterative evolution of some Western Tethyan Bathonian ammonites [*Benatinites* (B.) nov., *B. (Lugariceras)* nov. and *Hemigarantia*]. *Geobios* 39, 113–124.
- Schnyder, J., Ruffell, A., Deconinck, J.F. & Baudin, F. 2006: Conjunctive use of spectral gamma-ray logs and clay mineralogy in defining late Jurassic–early Cretaceous palaeoclimate change (Dorset, U.K.). *Palaeogeography, Palaeoclimatology, Palaeoecology* 229, 303–320.
- Schweigert, G. & Dietze, V. 1998: Revision der dimorphen Ammonitengattungen *Phlycticeras* HYATT – *Oecoptychius* NEUMAYR (Strigoceratidae, Mitteljura). *Stuttgarter Beiträge zur Naturkunde* 269, 1–59.
- Schweigert, G., Dietze, V. & Dietl, G. 2002: Erstnachweis der Ammoniten-Gattung *Parkinsonia* BAYLE, 1878 in der Garantiana-Zone (Tetragona-Subzone). *Stuttgarter Beiträge zur Naturkunde* 320, 1–15.
- Schweigert, G., Dietl, G. & Dietze, V. 2003: Neue Nachweise von *Phlycticeras* und *Oecoptychius* (Ammonitina: Strigoceratidae: Phlycticeratinae). *Stuttgarter Beiträge zur Naturkunde* 335, 1–21.
- Seyed-Emami, K., Schairer, G. & Bolourchi, M.H. 1985: Ammoniten aus der unteren Dalichy-Formation (oberes Bajocium bis unteres Bathonium) der Umgebung von Abe-Garm (Avaj, NW-Zentraliran). *Zitteliana* 12, 57–85.
- Seyed-Emami, K., Schairer, G. & Alavi-Naini, M. 1989: Ammoniten aus der unteren Dalichai-Formation (Unterbathon) östlich von Semnan (SE-Alborz, Iran). *Müncher geowissenschaftliche Abhandlungen* 15, 79–91.
- Seyed-Emami, K., Schairer, G., Aghanabati, S.A. & Fazl, M. 1991: Ammoniten aus dem Bathon der Gegend von Tabas - Nayband (Zentraliran). *Müncher Geowissenschaftliche Abhandlungen A* 19, 65–100.
- Seyed-Emami, K., Schairer, G., Aghanabati, S.A., Fürsich, F.T., Senowbari-Daryan, B. & Majidifard, M.R. 1998a: *Cadomites* aus der unteren Baghamshah-Formation (Oberbathon, Mittlerer Jura) SW Tabas (Zentraliran). *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 38, 111–119.
- Seyed-Emami, K., Schairer, G., Aghanabati, S.A., Fürsich, F.T., Senowbari-Daryan, B. & Majidifard, M.R. 1998b: *Bullatimorphites* aus dem Oberbathon (Mittlerer Jura) SW Tabas (Zentraliran). *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 38, 121–134.
- Singh, C.S.P., Jaitly, A.K. & Pandey, D.K. 1982: First report of some Bajocian-Bathonian (Middle Jurassic) ammonoids and the age of the oldest sediments from Kachchh, W. India. *Newsletters on Stratigraphy* 11, 37–40.
- Singh, C.S.P., Pandey, D.K. & Jaitly, A.K. 1983: Discovery of *Clydoniceras* Blake and *Gracilisphinctes* Buckman (Bathonian-Middle Jurassic Ammonites) in Kachchh, Western India. *Journal of Paleontology* 57, 821–824.
- Soares, A.F., Rocha, R.B., Elmi, S., Henriques, M.H.P., Mouterde, R., Alméras, Y., Ruget, C., Marques, J.F., Duarte, L.V., Carapito, M.C. & Kullberg, J.C. 1993: Le sous-bassin nord lusitanien: histoire d'un rift avorté (Trias-Jurassique moyen, Portugal). *Comptes Rendus de l'Académie des Sciences de Paris* 317, 1659–1666.

- Soussi, M., Mangold, C., Enay, R., Boughdiri, M. & Ben Ismail, M.H. 2000: Le Jurassique inférieur et moyen de la Tunisie septentrionale; corrélations avec l'axe Nord-Sud et paléogéographie. *Geobios* 33, 437–446.
- Steiner, M., Ogg, J., Sandoval, J. 1987: Jurassic magnetostratigraphy. 3. Bathonian-Bajocian of Carcabuey, Sierra Harana and Campillo de Arenas (Subbetic Cordillera, Southern Spain). *Earth and Planetary Science Letters* 82, 357–372.
- Stephanov, J. 1972: Monograph on the Bathonian ammonite genus *Siemiradzka* Hyatt, 1900 (nomenclature, taxonomy and phylogeny). *Travaux sur la Géologie de Bulgarie* 21, 5–82.
- Sturani, C. 1967: Ammonites and stratigraphy of the Bathonian in the Digne-Barrême area (South Eastern France). *Bolletino della Società Paleontologica Italiana* 5, 3–57.
- Sturani, C. 1971: Ammonites and stratigraphy of the “*Posidonia alpine*” beds of the Venetian Alps (Middle Jurassic, mainly Bajocian). *Memorie degli Istituti di Geologia e Mineralogia dell' Università di Padova* 28, 1–190.
- Sukanto, R. & Westermann, G.E.G. 1992: Indonesia and Papua New Guinea. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 181–193. Cambridge University Press, Cambridge.
- Taylor, D.G., Callomon, J.H., Hall, R., Smith, P.L., Tipper, H.W. & Westermann, G.E.G. 1992: Ammonites of the circum-Pacific region. In Westermann, E.G.E. (ed.): *The Jurassic of the Circum-Pacific*, 342–359. Cambridge University Press, Cambridge.
- Thierry, J. 2003: Les ammonites du Bathonien-Callovien du Boulonnais: biodiversité, biostratigraphie et biogéographie. *Geobios* 36, 93–126.
- Thierry, J., Clavel, B., Hantzpergue, P., Neraudeau, D., Rigollet, L. & Vadet, A. 1997: Distribution chronologique et géographique des échinides jurassiques en France: essai d'utilisation biostratigraphique. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires* 17, 253–271.
- Tiraboschi, D. & Erba, E. (in prep.): Calcareous nannofossil biostratigraphy of the Ravin du Bès section (Upper Bajocian-Lower Bathonian, SE France): evolutionary trends of *Watznaueria banesia* and new morphotypes of the genus *Rucinolithus*.
- Torrens, H. 1965: Revised zonal scheme for the Bathonian Stage of Europe. *Carpatho-Balkan Geological Association, VII Congress, Reports II, 1*, 47–55.
- Torrens, H. 1974a: Bathonian Stage. In N. Morton (editor): *The definition of standard Jurassic stages. Memoires du Bureau des Recherches Géologiques-Minières (Paris)* 75 (1971), 83–93.
- Torrens, H. 1974b: Standard zones of the Bathonian. *Memoires du Bureau des Recherches Géologiques-Minières (Paris)* 75 (1971), 581–604.
- Torrens, H. 1980: Bathonian correlation chart. In Cope, J.C.W., Duff, K.L., Parsons, C.F., Torrens, H.S., Wimbledon, W.A. & Wright, J.K. (eds.): *A correlation of Jurassic rocks in the British Isles. Part 2: Middle and Upper Jurassic. Geological Society, Spec. rep. 15*, 21–45.
- Torrens, H. 1987: Ammonites and stratigraphy of the Bathonian rocks in the Digne-Barrême area (South-Eastern France, Dept. Alpes de Haute Provence). *Bolletino della Società Paleontologica Italiana* 26, 93–108.
- Torrens, H. 2002: From d'Orbigny to the Devonian: some thoughts on the history of the stratotype concept. *Comptes rendues Palevol* 1, 335–345.
- Tseretely, L. 1989: *The representatives of family Oppeliidae BONARELLI in Bathonian deposits of Transcaucasus*. 112 pp. Metsniereba, Tbilisi.
- Vaez-Javadi, F. & Mirzaei-Ataabadi, M. 2006: Jurassic plant macrofossils from the Hojedk Formation, Kerman area, east-central Iran. *Alcheringa* 30, 63–96.
- Vörös A. 2001: Bajocian and Bathonian brachiopods in Hungary: a review. *Hantkeniana* 3, 177–182.
- Wang, Y., Mosbrugger, V. & Zhang, H. 2005: Early to Middle Jurassic vegetation and climatic events in the Qaidam Basin, Northwest China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 224, 200–216.
- Watkinson, M. P. 1989: *Triassic to Middle Jurassic sequences from the Lusitanian Basin Portugal, and their equivalents in other North Atlantic margin basins*. 390 pp. Unpublished Ph. D. Thesis, The Open University, Milton Keynes.
- Wendt, J. 1963: Stratigraphisch-paläontologische Untersuchungen im Dogger Westsiziliens. *Bolletino della Società Paleontologica Italiana* 2, 57–145.

- Wendt, J. 1971: Geologia del Monte Erice (provincial di Trapani, Sicilia occidentale). *Geologica Romana* 10, 53–76.
- Wernli, R. & Görög, A. 2007: Protoglobigérines et Oberhauserellidae (Foraminifères) du Bajocien-Bathonien du Jura méridional, France. *Revue de micropaléontologie* 50, 185–205.
- Westermann, G. 1958: Ammoniten-Fauna und Stratigraphie des Bathonien NW-Deutschlands. *Beihefte zum Geologischen Jahrbuch* 32, 1–103.
- Westermann, G.E.G. 1993a: Global bio-events in mid-Jurassic ammonites controlled by seaways. *Sytematics Association Special Volume* 47, 187–226.
- Westermann, G.E.G. 1993b: New Mid-Jurassic Ammonitina from New Zealand: implications for biogeography and oceanography. *GeoResearch Forum* 1, 179–185.
- Westermann, G.E.G. 1995: Mid-Jurassic Ammonitina from the Central Ranges of Irian Jaya and the origin of stephanoceratids. *Hantkeniana* 1, 105–118.
- Westermann, G.E.G. 2000: Marine faunal realms of the Mesozoic: review and revision under the new guidelines for biogeographic classification and nomenclature. *Palaeogeography, Palaeoclimatology, Palaeoecology* 163, 49–68.
- Westermann, G.E.G. & Callomon, J.H. 1988: The Macrocephalitinae and associated Bathonian and Early Callovian (Jurassic) ammonoids of the Sula Islands and New Guinea. *Palaeontographica* 203, 1–90.
- Westermann, G.E.G. & Getty, T.A. 1970: New Middle Jurassic Ammonitina from New Guinea. *Bulletins of American Paleontology* 57, 231–321.
- Westermann, G.E.G. & Hudson, N. 1991: The first find of Eurycephalitinae (Jurassic Ammonitina) from New Zealand and biogeographic implications. *Journal of Paleontology* 65, 689–693.
- Westermann, G.E.G. & Riccardi, A.C. 1980: The Upper Bajocian ammonite *Strenoceras* in Chile: first circum-Pacific record of the Subfurcatum Zone. *Newsletters on Stratigraphy* 9, 19–29.
- Westermann, G.E.G., Riccardi, A.C., Palacios, O. & Rangel, C. 1980: Jurásico medio en el Perú. *Instituto Geológico Minero y Metalúrgico, Boletín* 9, 1–47.
- Westermann, G.E.G., Hudson, N. & Grant-Mackie, J.A. 2000: Bajocian (Middle Jurassic) Ammonitina of New Zealand. *New Zealand Journal of Geology & Geophysics* 43, 33–57.
- Westermann, G.E.G., Hudson, N. & Grant-Mackie, J.A. 2002: New Jurassic Ammonitina from New Zealand: Bathonian–Callovian Eurycephalitinae. *New Zealand Journal of Geology & Geophysics* 45, 499–525.
- Wierzbowski, A., Jaworska, M. & Krobicki, M. 1999: Jurassic (Upper Bajocian-lowest Oxfordian) ammonitico rosso facies in the Pieniny Klippen Belt, Carpathians, Poland: its fauna, age, microfacies and sedimentary environment. *Studia Geologica Polonica* 115, 7–74.
- Yin, J. 2005: Middle Jurassic (Bathonian–Callovian) ammonites from the Amdo Area, Northern Tibet. *Acta Palaeontologica Sinica* 44, 1–16.
- Yin, J., Callomon, J.H. & Enay, R. 2000: A hiatus of 8 My duration in the Middle Jurassic with ammonite succession in Tethyan Himalaya (South Tibet). *Geobios* 33, 201–210.
- Zakharov, V.A. 2007: State of stage and zonal scales of the Jurassic System: global and regional aspects. In Zakharov, V.A. (ed.): *Jurassic System of Russia: problems of Stratigraphy and Paleogeography. Second all-Russian meeting*, 68–74. Yaroslavl State Pedagogical University, Yaroslavl.
- Zakharov, V.A., Bogomolov, Yu.I., Il'ina, V.I., Konstantinov, A.G., Kurushin, N.I., Lebedeva, N.K., Meledina, S.V., Nikitenko, B.L., Sobolev, E.S. & Shurygin B.N. 1998: Boreal zonal standard and biostratigraphy of the Siberian Mesozoic. *Russian Geology and Geophysics* 38 (1997), 965–993.
- Zany, D., Atrops, F., Marchand, D. & Thierry, J. 1990: Nouvelles données biostratigraphiques sur les séries du Bathonien et du Callovien des environs de Digne (Alpes-de-Haute-Provence). *Géologie Méditerranéenne* 17, 39–53.
- Zaton, M. 2006: *Tulites cadus* BUCKMAN, 1921 (Ammonoidea) from the Middle Bathonian of the Polish Jura and its biostratigraphic significance. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 243, 191–199.
- Zaton, M. & Marynowski, L. 2006: Ammonite fauna from uppermost Bajocian (Middle Jurassic) calcitic concretions from the Polish Jura - biogeographical and taphonomical implications. *Geobios* 39, 426–442.

- Ziegler, P.A. 1999: Evolution of the Arctic-North Atlantic and the Western Tethys. *American Association Petroleum Geologists Memoir 43*, 1–198.
- Zurcher, P. 1895: Compte rendu de la course du 23 septembre de Digne à Barrême. *Bulletin de la Société Géologique de France 23*, 866–873.

