

THE ORIGIN OF DWARF EARLY-TYPE GALAXIES IN THE VIRGO CLUSTER

E. Toloba¹, A. Boselli², R.F. Peletier³ and J. Gorgas¹

Abstract. We present new medium resolution kinematic data for a sample of 21 dwarf early-type galaxies (dEs) mainly in the Virgo cluster. These data are used to study the origin of the dE population inhabiting rich clusters. Within them we detect two different subpopulations: half of the sample (52%) are rotationally supported and the other half are pressure supported. We also find that the rotationally supported dEs are located in the outer parts of the cluster, present disk morphological shapes and are younger than those pressure supported that are concentrated in the core of the cluster without any underlying structures. Our analysis reveals that the rotationally supported objects have rotation curves similarly shaped to those of star forming galaxies of similar luminosities and follow the Tully-Fisher relation. This is expected if dE galaxies are the descendant of low luminosity star forming systems which recently entered the cluster and lost their gas due to a ram pressure stripping event, quenching their star formation activity and transforming them into quiescent systems, but conserving their angular momentum.

1 Introduction

The physical processes involved in galaxy formation and evolution still remain uncertain. Two main scenarios are nowadays in debate: in *the monolithic collapse* spheroids formed at a very early epoch from a rapid gas collapse followed by a strong starburst that passively evolved to the present while in *the hierarchical scenario* the most massive galaxies formed from a subsequent merging of smaller structures. A way of quantifying the relative role of these different mechanisms

¹ Departamento de Astrofísica y CC. de la Atmósfera, Universidad Complutense de Madrid, 28040 Madrid, Spain

² Laboratoire d'Astrophysique de Marseille, UMR 6110 CNRS, 38 rue F. Joliot-Curie, 13388 Marseille, France

³ Kapteyn Astronomical Institute, Rijksuniversiteit Groningen, Postbus 800, 9700 AV Groningen, The Netherlands

is to study dwarf galaxies, the most numerous objects in the universe (Ferguson & Binggeli 1994). Among those, dEs are the dominant galaxy population in high density regions such as clusters. Their low mass and large number make them ideal probes of those mechanisms through which a cluster environment can alter the appearance of galaxies, *i.e.* ram pressure and tidal interactions. If Λ CDM models predict dEs as the descendants of the cosmological building blocks, observations of nearby clusters favour the scenario where dEs formed after gas removal of late-type galaxies recently entering high density environments that quenched their star formation activity.

Several works have shown that the dE population is more complex and heterogeneous than originally thought (Lisker *et al.* 2006a; Lisker *et al.* 2006b; Lisker *et al.* 2007; Lisker *et al.* 2008) and an unexpected range in their stellar populations has been found (Michielsen *et al.* 2008). These evidences indicate a complex formation process shaping the evolution of dEs in clusters. The study of their kinematic properties is a powerful tool to unravel their origin.

2 Observations and data reduction

The observed sample consists of 18 dEs in the Virgo cluster and 3 in the field in the magnitude range $-17.5 > M_B > -15.5$. It was selected from the Virgo Cluster Catalog (VCC, Binggeli *et al.* 1985) requiring to have SDSS photometry and to be within GALEX pointings (see details in Toloba *et al.* submitted).

The observations were carried out at WHT(4.2 m) and INT(2.5 m) at Roque de los Muchachos Observatory (La Palma). We obtained long-slit spectroscopy (3445–8950 Å) along the major axis of the dEs reaching, with $2''$ slit-width in 1 hour integration per object, a spectral resolution of $\sim 45 \text{ km s}^{-1}$ (FWHM).

The reduction was done using standard procedures for long-slit spectra, using REDUCE (Cardiel 1999), a package specially focused on the parallel treatment of errors. Details on the observations and analysis are presented in Toloba *et al.* submitted.

3 Results

Given the radial decrease of the galaxy surface brightness, the S/N in the outer parts of some of the galaxies was not enough to reach a plateau on the rotation curves. To be cautious, for those galaxies where the v_{max} (maximum rotation) is measured at a radius $< 6''$, the value is considered as a lower limit and indicated with an arrow in all the Figures.

As indicated in Toloba *et al.* (2009), Figure 1 shows the anisotropy diagram where v_{max}/σ is the ratio between the maximum rotational velocity and the central velocity dispersion of the galaxies. The C_4 parameter quantifies how different the isophotes are from a perfect ellipse ($C_4 > 0$ means disk isophotes, $C_4 < 0$ boxy isophotes, Kormendy & Bender 1996). The agreement between Lisker *et al.* (2006a) and C_4 classifications is remarkable. All galaxies with disk isophotes (except VCC917) have been classified by Lisker *et al.* (2006a) as disks.




Fig. 1. Anisotropy diagram. The solid line is the model for an isotropic oblate system flattened by rotation (Binney 1978). Triangles are giant ellipticals (from Emsellem *et al.* 2007), with slow rotators ($v_{max}/\sigma < 0.1$) in dark grey, and fast rotators ($v_{max}/\sigma > 0.1$) in light grey. Blue, red and green symbols show our sample of dEs as classified by Lisker *et al.* (2006a) into disk, no disk and not in their sample respectively. The filled dots and the open squares indicate galaxies with and without disks on the basis of C_4 . Lower limits on v_{max} are indicated with arrows. The black symbols represent the median for dEs with (black dot)/without disk (black square) based on C_4 classification.

Figure 1, where rotationally supported galaxies (above the solid line) are separated from pressure dominated ones (below the solid line), shows that: 1) as ellipticals, dEs can be separated into slow ($v_{max}/\sigma < 0.1$) and fast ($v_{max}/\sigma > 0.1$) rotators Emsellem *et al.* (2007); 2) a large fraction (52%) of the dEs are rotationally supported and 3) all rotationally supported galaxies have morphological and/or photometric signs of a spiral origin.

The apparent discordance with Geha *et al.* (2003), who found that the majority of the dEs are not rotating, is due to the fact that their data are limited to the core of the galaxies, never reaching radii larger than $6''$, where the increase of the rotational velocity is generally observed (Toloba 2009; Toloba *et al.* 2010, submitted).

The left panel of Figure 2 shows $(v_{max}/\sigma)^* = \frac{v_{max}/\sigma}{\sqrt{\epsilon/(1-\epsilon)}}$, the anisotropic parameter corrected for inclination, as a function of the Virgocentric distance. It shows that rotationally supported systems are generally located in the cluster outskirts or in the field, while the pressure supported dwarfs are found only in the central regions of the cluster. This idea has been suggested before (Geha *et al.* 2003; van Zee *et al.* 2004), but no clear confirmation was found. Our observations reach distances further away from M 87 and the evidence for a trend is now clear. We emphasize, however, that any relation with the clustercentric distance is smeared out by projection effects.

The right panel of Figure 2 presents the anisotropic parameter *vs.* the ages from Michielsen *et al.* (2008). It shows that dEs with disks are ~ 3 Gyr younger




Fig. 2. Anisotropy parameter $vs.$ the angular distance to M 87 (centre of Virgo) in the left and $vs.$ the age in the right. The symbols are as in Figure 1. The light and dark grey rectangles limit the regions for fast and slow rotating Es (Emsellem *et al.* 2007). The solid line ($(v_{max}/\sigma)^* = 0.8$) divides the diagrams into rotationally and pressure supported galaxies. In the left panel the open triangles are from van Zee *et al.* (2004). In the right panel the ages are from Michielsen *et al.* (2008).




Fig. 3. *Left:* observed rotation curves of rotationally supported dEs (grey symbols) in comparison to the fitted rotation curves of late-type spiral galaxies (black solid and blue dashed lines) from Catinella *et al.* (2006). Blue dots represent the median of our rotationally supported dEs and the grey area indicates the 1σ deviation. *Right:* Tully-Fisher relation for our rotationally supported dEs (in dark blue) compared to the data on dEs by van Zee *et al.* (2004) (in light blue) and on normal spirals by Giovanelli *et al.* (1997b) (in grey) and De Rijcke *et al.* (2007) (DR07, in red).

than the others and while dEs with no signs of a spiral origin might be of all ages but on average old (5/8 of the squares are older than 7 Gyr), disk galaxies are preferentially young (9/12 of the dots are younger than 7 Gyr), suggesting that the 2 populations might have a different origin.

Catinella *et al.* (2006) made a systematic study of the shape of the rotation curves of late-type spiral galaxies as a function of luminosity. They fitted the rotation curves following the Polyex model which is an analytical function that

depends on 3 parameters: V_0 , r_{PE} and α , the amplitude, the exponential scale of the inner region and the slope of the outer part of the rotation curve, respectively. The mean fitted rotation curves from Catinella *et al.* (2006) are normalised to the optical radius (R_{opt} , radius containing 83% of the total I -band luminosity), and the velocities are corrected from inclination. This correction has been made using $\cos^2 i = ((1 - \epsilon)^2 - q_0^2) / (1 - q_0^2)$ (Giovanelli *et al.* 1997b), where i is the inclination, ϵ is the ellipticity and $q_0 = 0.3$, a conservative value for dwarf galaxies shaped as thick disks (see Toloba *et al.* submitted). To reach the dwarf regime in the fitted rotation curves by Catinella *et al.* (2006) we have extrapolated linearly the three Polyex parameters to $M_I = -18.49$, the mean I -band magnitude of the dEs here analysed.

Figure 3 shows that the rotation curves of star forming and quiescent systems are surprisingly similar despite the fact that the former are gas rich systems while the latter do not show any gaseous contribution (left panel).

If the rotationally supported dEs have the same shape of the rotation curves as the low luminosity late-type spirals, they should also follow the Tully-Fisher relation shown in the right panel of Figure 3, in particular considering that the rotation velocity is probably underestimated.

4 Discussion and conclusions

The strong morphological segregation observed in high density environments (Sandage *et al.* 1985; Ferguson & Binggeli 1994; Blanton *et al.* 2005) indicates that the cluster environment plays a major role in the formation of dEs. Figures 1 and 2 indicate that dEs can be divided into rotationally and pressure supported systems. They show that the rotationally supported dEs, characterised by a disk structure, are preferentially located in the field and in the periphery of the cluster, while the pressure supported systems, without disk underlying structures, are closer to the center. They also indicate that rotationally supported dEs have, on average, younger stellar populations than pressure supported dEs. Figure 3 indicates that the rotationally supported population have the same shape of the rotation curve observed in low luminosity late-type spiral systems and follow the Tully-Fisher relation, the most representative scaling relation for spiral galaxies. Since the angular momentum of these objects is conserved, the most plausible scenario for gas stripping is the ram pressure exerted by the dense and hot intergalactic medium (IGM) on the fragile inter-stellar medium of the low luminosity star forming galaxies freshly entering the cluster environment (see Boselli *et al.* 2008a; Boselli *et al.* 2008b).

For these rotationally supported objects gravitational interactions with the cluster potential or with other cluster members (galaxy harassment) can be excluded since they would, on relatively short time scales, reduce the angular momentum of the perturbed galaxies, leading to the formation of pressure supported systems. This process, however, could still be invoked to explain the kinematic and structural properties of the remaining pressure supported dEs populating the core of the cluster (half of our sample). Indeed the properties observed in these

pressure dominated objects can also be explained if they were star forming systems that entered into the cluster in early epochs, maybe through the accretion of groups where preprocessing was active, and later modified by galaxy harassment. Another possibility is to be star forming systems that entered into the cluster several Gyr ago, where ram pressure, although less efficient than today because of the lower density of the IGM and of the smaller velocity dispersion of the cluster still in formation, had the time through multiple cluster crossing to remove the gas and stop the star formation activity. This pressure supported systems are also consistent with an *in situ* formation, through the isotropic collapse of the gas at early epochs. In this scenario pressure supported systems would be the low luminosity extension of Es, with the exception that they probably did not undergo major merging events (De Lucia *et al.* 2006).

In summary, we present medium resolution ($R \sim 3800$) kinematic data for 21 dEs mainly located in the Virgo cluster. Their analysis has shown that there exists two populations of dEs in the Virgo cluster. Those located in the outskirts are rotationally supported and also show disk-like underlying structures as well as younger ages than those located in the core of Virgo, that are pressure supported objects without any underlying structure. The rotationally supported population also shows rotation curves similarly shaped to late-type galaxies of similar luminosity and they also follow the Tully-Fisher relation.

These observations are consistent with a picture where the dEs result from the transformation of star forming systems that entered the cluster and lost their gas through their interaction with the environment. The observed conservation of the angular momentum in the rotationally supported dEs suggests a ram pressure stripping event as the responsible of the gas removal whereas for the pressure supported systems other mechanisms such as harassment and the collapse of primordial gas inside the cluster also explain their observed properties.

References

- Binggeli, B., Sandage, A., & Tammann, G.A., 1985, AJ, 90, 1681
 Binney, J., 1978, MNRAS, 183, 501
 Blanton, M.R., Lupton, R.H., Schlegel, D.J., *et al.*, 2005, ApJ, 631, 208
 Boselli, A., Boissier, S., Cortese, L., & Gavazzi, G., 2008a, ApJ, 674, 742
 Boselli, A., Boissier, S., Cortese, L., & Gavazzi, G., 2008b, A&A, 489, 1015
 Cardiel, N., 1999, Ph.D. Thesis, Universidad Complutense de Madrid, Spain
 Catinella, B., Giovanelli, R., & Haynes, M.P., 2006, ApJ, 640, 751
 De Lucia, G., Springel, V., *et al.*, 2006, MNRAS, 366, 499
 De Rijcke, S., Zeilinger, W.W., *et al.*, 2007, ApJ, 659, 1172
 Emsellem, E., Cappellari, M., Krajnović, D., *et al.*, 2007, MNRAS, 379, 401
 Ferguson, H.C., & Binggeli, B., 1994, A&AS, 106, 67
 Geha, M., Guhathakurta, P., & van der Marel, R.P., 2003, AJ, 126, 1794
 Giovanelli, R., Haynes, M.P., Herter, T., *et al.*, 1997b, AJ, 113, 22
 Kormendy, J., & Bender, R., 1996, ApJ, 464, L119

- Lisker, T., Glatt, K., Westera, P., & Grebel, E.K., 2006a, *AJ*, 132, 2432
Lisker, T., Grebel, E.K., & Binggeli, B., 2006b, *AJ*, 132, 497
Lisker, T., Grebel, E.K., Binggeli, B., & Glatt, K., 2007, *ApJ*, 660, 1186
Lisker, T., Grebel, E.K., & Binggeli, B., 2008, *AJ*, 135, 380
Michielsen, D., Boselli, A., Conselice, C.J., *et al.*, 2008, *MNRAS*, 385, 1374
Sandage, A., Binggeli, B., & Tammann, G.A., 1985, *AJ*, 90, 1759
Toloba, E., Boselli, A., Gorgas, J., *et al.*, 2009, *ApJ*, 707, L17
Toloba, E., Boselli, A., Gorgas, J., *et al.*, *A&A*, submitted
van Zee, L., Skillman, E.D., & Haynes, M.P., 2004, *AJ*, 128, 121