

Accepted Manuscript



Pharmacology  
&  
Therapeutics

Lipid-lowering therapy with statins, a new approach to antiarrhythmic therapy

Juan Tamargo, Ricardo Caballero, Ricardo Gómez, Lucía Núñez, Miguel Vaquero, Eva Delpón

PII: S0163-7258(06)00205-1  
DOI: doi: [10.1016/j.pharmthera.2006.12.002](https://doi.org/10.1016/j.pharmthera.2006.12.002)  
Reference: JPT 5892

To appear in: *Pharmacology and Therapeutics*

Received date: 8 December 2006  
Accepted date: 8 December 2006

Please cite this article as: Tamargo, J., Caballero, R., Gómez, R., Núñez, L., Vaquero, M. & Delpón, E., Lipid-lowering therapy with statins, a new approach to antiarrhythmic therapy, *Pharmacology and Therapeutics* (2007), doi: [10.1016/j.pharmthera.2006.12.002](https://doi.org/10.1016/j.pharmthera.2006.12.002)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**P&T 21603**

**Lipid-lowering therapy with statins, a new approach to antiarrhythmic  
therapy**

**Short title: Antiarrhythmic effects of statins**

**Juan Tamargo, Ricardo Caballero, Ricardo Gómez, Lucía Núñez, Miguel Vaquero &  
Eva Delpón**

Department of Pharmacology, School of Medicine. Universidad Complutense. Madrid. Spain

**Author for correspondence:**

Juan Tamargo

Department of Pharmacology, School of Medicine

Universidad Complutense, 28040 Madrid, Spain

Phone: 91-3941472

Fax: 91-3941470

E-mail: [jtamargo@med.ucm.es](mailto:jtamargo@med.ucm.es)

**Abstract**

Inhibitors of 3-hydroxy-3-methylglutaryl coenzyme A reductase (statins) are the most effective and best-tolerated drugs to treat elevated levels of low-density lipoprotein cholesterol (LDL-C). In addition, they exhibit other effects unrelated to their lipid lowering effects (pleiotropic actions). In recent years, experimental and clinical evidence demonstrates that statins exert antiarrhythmic properties, reducing the recurrences of supraventricular and life-threatening ventricular arrhythmias both in patients with and without coronary artery disease. Thus, statins may constitute a novel therapeutic approach to cardiac arrhythmias. This article reviews the antiarrhythmic properties of statins as well as the possible mechanisms involved, including the lowering of LDL-C levels, the improvement of endothelial dysfunction and autonomic function, the stabilization of the atherosclerotic plaques, the antioxidant, anti-inflammatory, antithrombotic and cardioprotective properties and the modulation of transmembrane ion fluxes.

**Keywords:** statins, cardiac arrhythmias, atrial fibrillation, ventricular tachyarrhythmias, antiarrhythmic actions, pleiotropic effects.

**Abbreviations:**

AF: atrial fibrillation. APD: action potential duration. ATP: adenosine triphosphate.  $[Ca^{2+}]_i$ : intracellular  $Ca^{2+}$  concentration. CAD: coronary artery disease. CRP: C-reactive protein. GGPP: geranyl-geranyl pyrophosphate. HF: heart failure. HMG-CoA: 3-hydroxy-3-methylglutaryl coenzyme-A. HRV: heart rate variability.  $I_{Ca,L}$ : inward L-type  $Ca^{+}$  current. ICD: implantable cardioverter defibrillator.  $I_{Kur}$ : ultrarapid component of the delayed rectifier  $K^{+}$  current. IL: interleukin.  $I_{Na}$ : fast inward  $Na^{+}$  current.  $I_{to}$ : transient outward  $K^{+}$  current.  $K_{ATP}$ : ATP-sensitive  $K^{+}$  channels.  $K_v$ : voltage-gated  $K^{+}$  channels. LDL-C: low-density lipoprotein cholesterol. L-NAME: L-nitro-arginine methyl ester. LPC: L- $\alpha$ -lysophosphatidylcholine. LV: left ventricular. LVEF: left ventricular ejection fraction. MI: myocardial infarction. MMP: matrix metalloproteinase. NADPH: nicotinamide adenine dinucleotide phosphate. NO: nitric oxide. NOS3: endothelial nitric oxide synthase. NSVT: non-sustained ventricular tachycardia. NYHA: New York Heart Association. PAI-1: plasminogen activator inhibitor-1. SCD: sudden cardiac death. SMC: smooth muscle cell. TF: tissue factor. Th: T-lymphocytes helper. VF: ventricular fibrillation. VSMC: vascular smooth muscle cell. VT: ventricular tachycardia.

## Table of Contents

1. Introduction
2. Ventricular arrhythmias in patients with coronary artery disease
3. Mechanism involved in atrial fibrillation
4. Inhibitors of the HMG-CoA reductase
  - 4.1. Pleiotropic effects
5. Effects of statins on cardiac arrhythmias
  - 5.1. Ventricular arrhythmias
    - 5.1.1. Effects in experimental models
    - 5.1.2. Effects in clinical studies
  - 5.2. Atrial fibrillation
    - 5.2.1. Effects in experimental models
    - 5.2.2. Effects in clinical studies
6. Mechanisms of action involved in the antiarrhythmic effects of statins
  - 6.1. Antiischemic effects
  - 6.2. Role of pleiotropic effects on the antiarrhythmic effects of statins
    - 6.2.1. Anti-inflammatory effects
    - 6.2.2. Improvement of endothelial dysfunction
    - 6.2.3. Stabilization of vulnerable plaques
    - 6.2.4. Antioxidant effects
    - 6.2.5. Cardioprotective effects
    - 6.2.6. Improvement of autonomic function
    - 6.2.7. Effects on cardiac ion channels and transporters
      - 6.2.7.1. Role of membrane cholesterol on transmembrane ion fluxes
      - 6.2.7.2. Role of statins on transmembrane ion fluxes
7. Conclusions

## 1. Introduction

Cardiac arrhythmias are a leading cause of morbidity and mortality (Rubart & Zipes 2005). Sudden cardiac death (SCD) caused by ventricular tachyarrhythmias, such as ventricular tachycardia (VT) and fibrillation (VF), and less frequently by bradyarrhythmias and pulseless electrical activity, accounts for more than 300,000 deaths annually in the United States alone and it remains a major and unresolved public health problem (Rubart & Zipes, 2005). Coronary artery disease (CAD) is the underlying structural heart disease in at least 80% of victims of these life-threatening ventricular tachyarrhythmias, even when SCD is often associated with other heart diseases, such as dilated nonischemic and hypertrophic cardiomyopathies and heart failure (HF), drugs and assorted noncardiac diseases, and may also occur in healthy young individuals (Zipes & Wellens, 1998).

Although it was hoped that current antiarrhythmic drugs would protect against VT/VF, the results of clinical trials have been quite disappointing. In fact, standard antiarrhythmic drug therapy fails to reduce, or even increases, the incidence of SCD and exhibits important limitations arising from the risk of severe side effects and potentially lethal proarrhythmia (Pratt, et al., 1998). The reason for these results could be that for many decades, antiarrhythmic drug development has been concentrated on manipulating membrane cardiac ionic currents, while the triggering events and the arrhythmogenic substrates were not taken into consideration. Moreover, diseases most frequently associated with lethal arrhythmias, including CAD and most common cardiomyopathies, do not result from primary abnormalities in ionic channels.

Because most adult patients at risk of SCD in industrialised societies suffer from atherogenic dyslipemias and CAD (Zipes & Wellens, 1998), when considering possible therapeutic approaches, it is appropriate to pay attention to the cellular and molecular mechanisms underlying sudden arrhythmic death in atherosclerotic heart disease. If CAD and occlusive coronary thrombi associated with rupture of the atherosclerotic plaque are the trigger factors of the life-threatening ventricular tachyarrhythmias, the treatment should target the pathogenic mechanisms leading to CAD instead of using standard antiarrhythmic drugs. In fact, the greatest reduction in cardiovascular mortality, including SCD, in patients with a previous myocardial infarction (MI) and HF has been observed with beta-blockers (Gottlieb et al., 1998; López-Sendón et al., 2004) and non-antiarrhythmic drugs, i.e. those without major direct cardiac electrophysiological effects, such as angiotensin-converting enzyme inhibitors, angiotensin-receptor blockers, aldosterone antagonists, thrombolytic, antithrombotic and lipid-lowering agents (Alberte & Zipes, 2003).

Therefore, it can be speculated that the treatment of hypercholesterolemia in patients with CAD would result not only in a reduction of major coronary events, but ultimately also in a reduction of arrhythmic episodes and SCD. Recent evidence has indicated that inhibitors of the 3-hydroxy-3-methylglutaryl coenzyme-A (HMG-CoA) reductase (“statins”) can reduce the incidence of both supraventricular and ventricular arrhythmias in patients with or without CAD. However, this effect is not observed with other lipid-lowering drugs. Therefore, this article focuses on the antiarrhythmic effects of statins. We firstly analyzed the mechanisms involved in ventricular and atrial arrhythmias in patients with CAD, then the experimental and clinical evidence supporting the antiarrhythmic properties of statins and, finally, the possible mechanisms involved in these antiarrhythmic actions.

## **2. Ventricular arrhythmias in patients with coronary artery disease**

Atherosclerosis is primarily a chronic vascular inflammatory disorder characterized by infiltration of the arterial wall by macrophages and T-lymphocytes and elevated levels of inflammation markers such cytokines (tumor-necrosis factor [TNF]- $\alpha$ , IL-1, IL-6, IL-18, CD40 ligand), chemokines (IL-8, monocyte chemoattractant protein [MCP]-1), acute phase reactants (high-sensitivity CRP-hsCRP, serum amyloid A) and soluble cell adhesion molecules (E- and P-selectins, vascular, intercellular and their counter ligand CD11a/CD18 in the monocyte) (Bellosta et al., 1998; Bonetti et al., 2002; Davignon, 2004; Schonbeck & Libby, 2004; Jain & Ridker, 2005; Liao & Laufs, 2005; Ray & Cannon, 2005). Plaque rupture followed by superimposed thrombus formation is a major cause of acute coronary syndromes (Conti, 2001; Libby, 2001). Plaque composition and stability, rather than the volume or the severity of stenosis, are the main determinants of atherosclerotic complications (Dupuis et al., 1999). Unstable plaques, prone to fissuring or disruption are characterized by a thin fibrous cap, high-lipid content, few SMCs, and excess of activated macrophages and T-lymphocytes in the cap (Conti, 2001; Libby, 2001). Activated macrophages play a key role in destabilizing the plaque by their ability to secrete several cytokines, which in turn decreases collagen production by vascular SMCs, and MMPs and elastases, which degrades the extracellular matrix, leading to weakening of the fibrous cap and plaque instability (Galis et al., 1994).

We already mention that CAD is the underlying heart disease in 80% of victims of life-threatening ventricular tachyarrhythmias and recent occlusive thrombus is present in 15% to 64% of patients who died suddenly (Mehta et al., 1997; Zipes & Wellens, 1998; Rubart & Zipes, 2005). Post-mortem examinations suggest that SCD victims can be divided into: a)

patients without a prior history of heart disease in whom acute coronary thrombosis that leads to fatal VF (sometimes preceded by polymorphic VT) may be the first manifestation of coronary atherosclerosis, and b) substrate-related or nonischemic SCD, which occurs more frequently in the presence of a myocardial scar from a previous MI and impaired left ventricular (LV) function (Mehta et al., 1997). In these patients, scarred myocardium may provide the substrate for reentrant ventricular arrhythmias, most commonly monomorphic VT that may degenerate into VF. In fact, in experimental models acute ischemia superimposed on a previous MI is an arrhythmogenic mechanism, whereas the same degree of acute ischemia in the absence of myocardial scar is not arrhythmogenic (Furukawa et al., 1991). Also, ischemia superimposed at the border zone of a 1-month MI, where reentrant circuits are formed, is more arrhythmogenic than ischemia at a distance from the infarct zone in swine (Cinca et al., 1997). In many patients, however, the combination of both mechanisms, myocardial ischemia and scarring, is probably responsible for the genesis of lethal ventricular arrhythmias.

Abrupt cessation of coronary blood flow decreases cellular adenosine triphosphate (ATP) content and produces marked changes in the distribution of  $\text{Na}^+$ ,  $\text{Ca}^{2+}$  and  $\text{K}^+$  ions across the cardiac membrane (Tomaselli & Zipes, 2004; Rubart & Zipes, 2005). Net cellular  $\text{K}^+$  efflux through ATP-dependent, inward rectifier and other voltage-gated  $\text{K}^+$  channels and subsequent accumulation of  $\text{K}^+$  in the interstitial fluid leads to membrane depolarization, slows intracardiac conduction velocity and increases dispersion of ventricular repolarization and refractoriness, effects that facilitate re-entrant arrhythmias. Ischemia also inhibits  $\text{Na}^+$ - $\text{K}^+$  ATPase activity and increases that of the  $\text{Na}^+$ / $\text{H}^+$  exchanger, effects leading to an increase in intracellular  $\text{Na}^+$  concentration. These changes in  $\text{Na}^+$  and  $\text{K}^+$  across the cardiomyocyte membrane in turn increase the intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) via the  $\text{Na}^+$ - $\text{Ca}^{2+}$  exchanger operating in the reverse mode and activate L-type  $\text{Ca}^{2+}$  channels. This progressive  $\text{Ca}^{2+}$  overload increases the probability for triggered-induced ventricular arrhythmias associated to delayed afterdepolarizations, while augmented L-type  $\text{Ca}^{2+}$  current ( $I_{\text{Ca,L}}$ ) can also lead to ventricular arrhythmias triggered by early afterdepolarizations. Increased  $\text{Ca}^{2+}$  overload together with an increased production of oxygen free radicals, acidosis and increased catecholamine levels would lead to increased automaticity that may facilitate the development of VT/VF.

Recurrent VT in patients with CAD is believed to result also, in part, from cardiac structural remodeling due to the formation of anatomic barriers (interstitial fibrosis), regions of slow conduction and prolonged and heterogeneous repolarization across the ventricular wall

(Pfeffer & Braunwald, 1990). A MI often initiates the remodeling process, but ischemia, inflammation, neurohumoral activation, and other factors (including electrolyte and acid-base abnormalities, hypoxia, drug-induced proarrhythmic effects and superimposition of ischemia on a previous MI) continue for years. The importance of remodeling to arrhythmogenesis in CAD is suggested by the delay between MI and the first episode of VT (Anderson, 2003).

The observation that beta-blockers improve survival and reduce the incidence of SCD in patients after MI suggests that in both acute ischemia and substrate-related SCD the increased sympathetic tone plays an important role in the genesis of SCD from VT/VF. (Mehta et al., 1997; Shusterman et al., 1998). In addition to electrical and structural remodeling, neural remodeling in the form of sympathetic nerve sprouting may also result in ventricular arrhythmias in dogs with MI and complete atrioventricular block (Cao et al., 2000a) and in patients with severe HF (Cao et al., 2000b). Myocardial infarction results in sympathetic denervation of regions of necrosis/fibrosis and of myocardium in areas distal to the infarct and hyperinnervation in the periphery of injured myocardium, this latter effect being significantly higher in patients with ventricular tachyarrhythmias (Cao et al., 2000a,b). These changes can create gradients of refractoriness and excitability contributing to the arrhythmogenic substrate and are correlated directly with the occurrence of spontaneous ventricular arrhythmias and SCD. In addition, local elevated noradrenaline levels may induce coronary artery vasoconstriction, increasing myocardial regional ischemia. In a rabbit model, hypercholesterolemia results in cardiac hypertrophy and sympathetic hyperinnervation, increases  $I_{Ca,L}$  density and prolongs in an inhomogeneous manner the duration of the ventricular action potentials (QT interval) leading to an increased dispersion of repolarization even in the absence of coronary artery stenosis or MI (Liu et al., 2003). Interestingly, this neural and electrophysiological remodeling is significantly higher in patients with a history of tachyarrhythmias than in the patients without tachyarrhythmias (Cao et al., 2000a,b). The prolongation of the ventricular action potentials and the increase in  $I_{Ca,L}$  during sympathetic stimulation could lead to an increase in the  $[Ca^{2+}]_i$ , which appears to be the common denominator in the generation of triggered activity. Thus, increased sympathetic activity and sympathetic hyperinnervation may be, in part, responsible for the occurrence of ventricular tachyarrhythmias and SCD in post-MI patients (Cao et al., 2000a,b; Rubart & Zipes, 2005).

### 3. Mechanisms involved in atrial fibrillation

Atrial fibrillation (AF) is the most common arrhythmia in clinical practice and is associated with many forms of heart disease, including CAD, HF, arterial hypertension, valvular disease and cardiac surgery (Fuster et al., 2006). It is associated with significant morbidity and mortality, accounting for over one-third of all strokes over the age of 60 years (Wolf et al., 1987). AF is often a self-perpetuating disease. In many patients initial paroxysms of AF, that terminate spontaneously within a few days, become more frequent and of longer duration to the point where they become persistent (i.e. AF requires cardioversion to restore sinus rhythm) and then permanent over time (AF cannot be converted to sinus rhythm). In the past decade we learned that AF results in progressive changes in the electrophysiological, structural and contractile properties of the atria in a way that promotes its maintenance and recurrence (i.e. “AF begets AF”) and may alter the response to antiarrhythmic (Nattel, 2002; Tamargo et al., 2004a). Electrical changes include heterogeneous shortening of atrial action potential duration (APD) and refractoriness, loss of atrial refractoriness adaptation to rate and slow intra-atrial conduction, while structural changes include atrial dilatation, fibrosis, hypertrophy, fatty infiltration, increase in intracellular glycogen, mitochondrial swelling, sarcoplasmic reticulum disruption, myolysis and patchy apoptosis (Nattel, 2002; Tamargo et al., 2004a; Fuster et al., 2006).

Other factors potentially involved in the induction or maintenance of AF include focal inflammation, endothelial/endocardial dysfunction, oxidative stress, ischemia and autonomic nervous system activity (Tamargo et al., 2004a; Fuster et al., 2006). Evidence for an inflammatory contribution to some forms of AF was initially suggested by the high incidence (25-40%) of AF after cardiac surgery and the association of AF with myocarditis and pericarditis (Boos et al., 2006). Fromer et al (1990) analyze two cases of drug refractory ectopic atrial tachycardia and histopathologic examination of the resected right atria shows right atrial focal myocarditis without concomitant ventricular myocarditis. Thus, they conclude that focal atrial myocarditis may represent one cause of drug-resistant ectopic atrial tachycardia. The hypothesis that lone paroxysmal AF may be due to isolated atrial myocarditis is supported by the presence of atrial lymphocytic myocarditis in 66% of the endomyocardial biopsies of the right atrial septum of patients with drug-resistant paroxysmal lone AF (Frustaci et al., 1997) and the prevention of AF recurrences in patients treated with oral methylprednisolone (Dernellis & Panaretou, 2004) or nonsteroidal anti-inflammatory medications after coronary artery bypass graft surgery (Cheruku et al., 2004). In patients undergoing cardiopulmonary bypass surgery, the activation of the complement

system and the release of proinflammatory cytokines follow a biphasic behaviour. Interleukin-6 (IL-6) levels reached a maximum at 6 hours post-surgery, while C-reactive protein (CRP) levels peaked on the second and third postoperative days, coinciding with the peak incidence of atrial arrhythmias (Bruins et al., 1997). Moreover, a close correlation between the -174C/G IL-6 gene variant, the inflammatory response to surgery and the development of postoperative atrial arrhythmias has been recently reported (Gaudino et al., 2003). Furthermore, the presence of circulating autoantibodies against cardiac myosin heavy chain in a significant percentage of patients with idiopathic paroxysmal atrial fibrillation raises the possibility of an inflammatory autoimmune process in some patients with paroxysmal atrial fibrillation (Maixent et al., 1998).

The presence of systemic inflammation determined by elevations in the levels of high-sensitivity CRP (hs-CRP), a marker of systemic inflammation, has also been associated with persistence of AF. In addition, there is a significant association between circulating hs-CRP levels and risk of future cardiovascular events, including stroke, SCD, plaque rupture, peripheral vascular disease and myocardial infarction (Ridker, 2001a; Szmitko et al., 2003). Levels of CRP are higher in patients with atrial arrhythmias compared with controls in sinus rhythm, those with persistent AF have higher CRP levels than those with paroxysmal AF and both have higher levels than controls (Chung et al., 2001). Moreover, levels of CRP are higher in patients with paroxysmal AF compared with controls and a low CRP level predicts successful cardioversion (Dernellis & Panaretou, 2005). In the *Cardiovascular Health Study* high CRP levels are not only associated with the presence of AF but also predict the patients at increased risk for future development of AF (Avilés et al., 2003). Moreover, CRP levels remain a significant independent predictor of AF even after adjustment for multiple risk factors of AF, including hypertension and CAD (Avilés et al., 2003; Anderson et al., 2004). Thus, and even when the cause of elevated CRP levels in AF patients remains unknown, elevated levels of hs-CRP may reflect an inflammatory state, which subsequently may promote the persistence of the arrhythmia (Chung et al., 2001).

Recent evidence indicates that endothelial dysfunction and increased oxidative stress may also play an important role in maintaining AF (Mihm et al., 2001; Cai et al., 2002; Nattel, 2002; Tamargo et al., 2004a; Fuster et al. 2006). In experimental models, AF induced by rapid atrial pacing is associated with a marked decrease (70%) in endocardial nitric oxide (NO) bioavailability and endothelial NO synthase (NOS3) expression in the left atria, while both values remained unaltered in the aorta and right atria (Cai et al., 2002). Rapid atrial pacing also decreases atrial tissue levels of ascorbate, an antioxidant and peroxynitrite

decomposition catalyst, and increases atrial oxidative stress and peroxynitrite formation, leading to loss of NO bioactivity. Moreover, NAD(P)H-dependent superoxide production is increased in the fibrillating human atria, which suggests that reactive oxygen species may play an important role in the atrial oxidative injury and both in the electrical and structural remodeling of the atria in patients with AF (Kim et al., 2005). Supplementation with ascorbate attenuates the pacing-induced atrial effective refractory period shortening and the accumulation of peroxynitrite in chronically instrumented dogs and when given before and for 5 days following cardiac bypass surgery and reduces the incidence of postoperative AF (16.3% vs 34.9% in control subjects (Carnes et al., 2001).

Experimental data suggests that atrial ischemia may create a substrate for AF maintenance (Sinno et al., 2003; Fuster et al., 2006). In a canine model, atrial ischemia induced by occlusion of an atrial branch of the right coronary artery that did not provide blood flow to the ventricles promotes the persistence of AF induced by burst pacing, apparently by causing localized atrial conduction slowing within the ischemic zone, which may stabilize atrial reentry that maintains AF (Sinno et al., 2003). In addition, stenosis of the right coronary or the sinoatrial nodal arteries predisposes patients to AF after coronary artery bypass grafting (Mendes et al., 1995; Kolvekar et al., 1997).

Increased sympathetic or parasympathetic tone has been implicated in the initiation of AF. Autonomic ganglia containing parasympathetic and sympathetic fibers are present on the epicardial surface of both atria, clustered on the posterior wall near the ostia of the pulmonary veins, superior vena cava and coronary sinus. In animal models, parasympathetic stimulation shortens atrial and pulmonary vein refractory periods, potentiating initiation and maintenance of AF (Schauerte et al., 2000), whereas vagal denervation of the atria prevents induction of AF (Elvan et al., 1995). Pure autonomic initiation of clinical AF is uncommon and seen only in situations of high sympathetic or high vagal tone, but recordings of heart rate variability disclose autonomic perturbations that precede the onset of AF in some patients (Bettoni & Zimmermann, 2002; Tomita et al., 2003).

#### **4. Inhibitors of the HMG-CoA reductase**

Statins are the most effective and best-tolerated drugs to treat elevated levels of low-density lipoprotein cholesterol (LDL-C). They competitively inhibit HMG-CoA reductase, which is responsible for converting HMG-CoA to mevalonate, the rate-limiting step for cholesterol synthesis (Figure 1). The  $K_i$  (inhibition constant) values for the statin-enzyme complexes range between 0.1 and 44 nM, whereas Michaelis constant ( $K_m$ ) for HMG-CoA is 4  $\mu$ M

(Holdgate et al., 2003). Statins possess an HMG-like moiety that inhibits HMG-CoA reductase by binding to the active site of the enzyme, thus sterically inhibiting the substrate from binding. The statins differ from each other in the rigid, hydrophobic structures covalently linked to the HMG-like moiety. Lovastatin, pravastatin and simvastatin are fungal derivatives that share a common hexahydronaphthalene ring present in the structure of compactin, the first statin studied in humans. Fluvastatin, atorvastatin and rosuvastatin are fully synthetic inhibitors containing a fluorophenyl group that forms a structural analog of the HMG-CoA intermediate. Lipophilic statins (atorvastatin, lovastatin, simvastatin) can penetrate easily cell membrane in any organ, while hepatic uptake of hydrophilic statins (pravastatin and rosuvastatin) is dependent on the presence of a specific carrier-mediated mechanism (Yamazaki et al., 1993).

Statins inhibit cholesterologenesis in the liver and other tissues, which results in an increased expression of LDL receptors that clear cholesterol-rich LDL and LDL precursors from the circulation (Brown & Goldstein, 1998). Thus, statins reduce levels of total cholesterol (20-35%), LDL-C (25-55%) and triglycerides (10-20%) and increase high-density lipoprotein (HDL) cholesterol (5-10%) levels (Cheung et al., 2004, Cholesterol Treatment Trialists' Collaborators, 2005). Results from both secondary and primary large-scale, multicenter, prevention trials have consistently demonstrated that statin therapy significantly reduce the risk of major coronary events (unstable angina pectoris, MI, coronary death and need for coronary revascularization), stroke and all-cause and cardiovascular mortality (Cheung et al., 2004; Cholesterol Treatment Trialists' Collaborators, 2005). These beneficial effects are observed in patients with and without CAD, hypercholesterolemia, hypertension, diabetes mellitus and, particularly, in smokers.

#### **4.1. Pleiotropic effects**

Although statins exert their major effects on CAD by lowering LDL-C, a multitude of potentially cardioprotective effects have been ascribed to these drugs, largely based on *in vitro* and *ex vivo* data (Bonneti et al., 2002; Davignon, 2004). The pleiotropic effects of statins that are summarized in Table 3 have been related to the inhibition of the synthesis of isoprenoid intermediates of the mevalonate pathway, such as isopentenyl adenosine, farnesylpyrophosphate and geranylgeranyl pyrophosphate (GGPP) (Figure 1). These intermediates serve as lipid anchors for post-translational modification of a variety of proteins involved in intracellular signal transduction pathways, including heterotrimeric G proteins and small guanosine-triphosphate (GTP)-binding proteins, such as Ras, Rho and Rac1. Indeed, isoprenoids are essential for maintaining membrane fluidity and regulation of

cell growth and differentiation, gene expression, cytoskeletal assembly and cell motility, protein and lipid trafficking, nuclear transport, and host defence (Bellosta et al., 1998; Bonetti et al., 2002; Davignon, 2004; Liao & Laufs, 2005).

The pleiotropic actions of statins include improved endothelial function, modulation of autonomic function, stabilization of vulnerable plaques and antioxidant, anti-inflammatory, antithrombotic and cardioprotective effects (Table 3). In addition, statins may exert direct antiarrhythmic effects by modulating the physicochemical properties of the cardiac sarcolemma resulting in alterations of transmembrane cardiac ion fluxes that may directly affect the electrophysiological properties of cardiac muscle.

## **5. Effects of statins on cardiac arrhythmias**

### **5.1. Ventricular arrhythmias**

**5.1.1. Effects in experimental models.** In anesthetized rats subjected to 5-min ischemia and 10-min reperfusion, chronic oral administration of pravastatin (0.02, 0.2, or 2 mg/kg for 22 days, once daily) does not modify serum cholesterol but significantly reduces the incidence of ischemia-induced VT (70% vs 9% at 2 mg/kg) and the incidence of reperfusion-induced lethal VF (90% vs 20% at 0.2 mg/kg) (Chen et al., 2003). However, acute pravastatin (0.2 or 2 mg/kg, once orally) and chronically administered fluvastatin (0.2, 2, or 4 mg/kg) have no significant effect on these arrhythmias. In these experiments, there are no significant changes in blood pressure, heart rate, QT interval, and serum cholesterol among pravastatin-, fluvastatin-, and vehicle-treated groups. Thus, the antiarrhythmic effects of pravastatin are independent of its cholesterol-lowering effect.

**5.1.2. Effects in clinical studies.** Two large secondary prevention trials have provided evidence that statins reduced SCD. The *Scandinavian Simvastatin Survival Study* (1994) included patients with CAD and high cholesterol levels. During a follow-up of 5.4 years, the instantaneous death and death within 1 h (most likely due to VT and/or VF) in the absence of confirmed MI accounted for 63 of 189 in the placebo group as compared to 37 of 111 coronary deaths in the simvastatin group (crude reduction in SCD of 26%) and this reduction in presumable arrhythmic death was less than the effect on all CAD deaths (42%). The *Long-term Intervention with Pravastatin in Ischemic Disease* (The LIPID Study Group, 1998) study enrolled patients with CAD and a broad range of cholesterol levels. In the placebo group, 211 deaths were classified as sudden as compared to 182 in pravastatin-treated patients, which represents a reduction of SCD of 14%. Although in these trials no separate statistical analysis of the SCD was performed, these observations suggest that

statins might exert a beneficial effect on the incidence of lethal ventricular arrhythmias and reduce the relative risk of SCD [RR 0.20, 95% CI 0.05-0.33] in patients with CAD.

Several studies found that statins can suppress ventricular arrhythmias in patients with or without CAD (Table 2). De Sutter et al (2000) explore the antiarrhythmic effects of lipid-lowering drugs in an observational non-randomized study performed in 78 patients with CAD and life-threatening ventricular arrhythmias treated with an implantable cardioverter defibrillator (ICD). Patients with ventricular tachyarrhythmias in the setting of acute MI and those in whom the arrhythmia could not be induced at electrophysiological study after revascularization were excluded. After a mean follow-up of 16 months, patients receiving lipid-lowering drugs (statins-59%, fibrates-41%) have a lower incidence of recurrences of VT/VF (22% vs. 57%,  $P = 0.004$ ) and of the combined end-point of cardiac death and hospitalization (15% vs 45%,  $P = 0.015$ ). Moreover, in this study, ICD therapy for fast VT is also significantly lower in patients receiving lipid lowering drugs (11 vs 35%,  $P = 0.032$ ). These results suggest that the use of lipid lowering drugs is associated with a reduction of recurrences of ventricular tachyarrhythmias in patients with CAD and ICD implants.

In the *Antiarrhythmics Versus Implantable Defibrillators* (AVID) study including 237 patients with CAD and ventricular arrhythmias treated with an ICD, lipid-lowering therapy (statins-79%, fibrates-19%, bile acid resins-3%) is also associated with a reduction in the relative hazard for VT/VF recurrence (40%,  $P = 0.003$ ), all-cause mortality (36%,  $P = 0.03$ ) and cardiac mortality (39%,  $P = 0.04$ ) (Mitchell et al., 2003). The benefit appears early in follow-up in this study that included patients with several comorbid conditions (congestive HF-45%, chronic lung-16% and renal disease-8%) excluded from the Scandinavian Survival Study Group (1994) and LIPID trials (The LIPID Study Group, 1998). In the *Multicenter Automatic Defibrillator Implantation* (MADIT)-II study, 654 patients receiving an ICD were categorized by the percentage of days each patient received statins during follow-up (Vyas et al., 2006). In this study, the cumulative rate of ICD therapy for VT/VF or SCD, whichever occurred first, was significantly reduced in those with  $\geq 90\%$  statin usage compared to those with lower statin usage ( $\leq 10\%$ ,  $P = 0.01$ ). The time-dependent statin:no statin therapy hazard ratio was 0.65 for the endpoint of VT/VF or cardiac death ( $P < 0.01$ ) and 0.72 ( $P = 0.046$ ) for VT/VF after adjusting for relevant covariates. In another study performed on 281 patients who had CAD and underwent ICD implantation statin therapy was associated with a significant decrease in the risk of ventricular arrhythmias that required ICD therapy (30 vs 50%,  $P = 0.0007$ ), an effect that was attributed to a decrease in ischemia (Chiu et al., 2005). After adjusting for age, LV ejection fraction (LVEF), gender and beta-

blocker therapy, hazard ratio for the first ICD therapy among patients who used statins was 0.60 ( $P = 0.01$ ). These findings indicate that statins reduce the incidence of life-threatening ventricular arrhythmias in high-risk patients with CAD. Patients with ischemic heart disease are predisposed to worse short- and long-term outcomes after VF out-of-hospital cardiac arrest. In 87 patients who survive to hospital discharge with neurologic recovery, use of an ICD and statin therapy is associated with decreased mortality (Bunch et al., 2006)

Other studies have analyzed the effects of statins in patients undergoing coronary surgery. In a non-randomized retrospective study performed on 323 patients, preoperative statin therapy significantly reduced the incidences of death, MI, unstable angina and arrhythmias (sustained AF, atrial flutter, sustained or non-sustained VT and VF) ( $P = 0.006$ ) 1-year after coronary artery bypass graft surgery (Dotani et al., 2000). In addition, there was an increase in the cardiovascular adverse outcomes among patients taking other lipid-lowering drugs compared to those taking statins (27% vs 12%,  $P = 0.01$ ).

Ventricular late potentials correspond to fragmented activation of ventricular tissue and are thought to originate from areas of slow and inhomogeneous conduction within diseased myocardium. Pravastatin administered early after acute ST elevated MI ( $< 6$  h) decreases the rate of ventricular late potentials and the incidence of in-hospital ventricular arrhythmias (26 vs. 63%,  $P=0.021$ ) and cardiovascular events following thrombolytic therapy (Kayikcioglu et al., 2003). These effects were attributed to the prevention of new myocardial ischemic episodes due to early stabilization of vulnerable plaques, improvement in endothelial function, a decrease in platelet thrombus deposition and a reduction in inflammation. Data from the *National Registry of Myocardial Infarction 4*, a prospective, observational study, also found that new treatment with statins in the first 24 h of admission for acute MI was associated with a decreased risk of in-hospital mortality compared with no statin use (4% as compared with 15.4%,  $P < 0.001$ ) (Fonarow et al., 2005). Continuation with statin therapy in the first 24 hours of hospitalization for acute MI decreased risk of mortality compared with no statin use (5.3% vs 15.4%,  $P < 0.001$ ). In contrast, patients who have been treated with statins before hospitalization but whose therapy was discontinued had a mortality risk that was slightly higher than the risk in patients who did not use statins (16.5% vs 15.4%,  $P > 0.05$ ). Early statin use, whether newly initiated or continued, was also associated with a lower incidence of cardiac arrest, cardiogenic shock, cardiac rupture and VT/VF (4.5% and 3.7% vs 5.6%). Thus, the use of statins within the first 24 hours of hospitalization for acute MI is associated with a significant lower rate of early complications and in-hospital mortality.

Non-sustained VT (NSVT) occurring after acute ST-elevation MI is associated with an increased long-term mortality. In 346 patients with NSVT occurring after acute ST-elevation MI, those without statin treatment and no NSVT, 1 year mortality was 9.2%, but increased to 25% (OR 3.02; 95% CI 1.47-6.20) if NSVT were present (Lorenz et al., 2005). In patients on statin treatment and no NSVT, 1-year mortality was only 3.2%, and in the presence of NSVT 1-year mortality was not significantly increased anymore (5.3%). Thus, under statin therapy NSVT occurring after acute ST-elevation MI is not associated with an adverse long-term prognosis, suggesting that statin therapy may be associated with a stabilization of myocardium against pro-arrhythmic events (Lorenz et al., 2005).

QT interval dispersion in the surface electrocardiogram, a marker of repolarization inhomogeneity, is a noninvasive method for the evaluation of patients with increased risk of SCD or life-threatening ventricular arrhythmias. Fluvastatin treatment for 12 months decrease the QT dispersion in hyperlipidemic patients, suggesting that statins may also decrease inhomogeneity of ventricular repolarization (Mark & Katona, 2000). In postinfarction patients with severe LV dysfunction, increased QT variability is associated with an increased risk for VT/VF (Haigney et al., 2004). In a prospective case-control study on 80 patients with depressed LV function (LVEF < 0.30%) and in New York Heart Association (NYHA) class III for at least two months, atorvastatin shortened the QT interval and decreased QT variability and these effects were independent from changes in LDL-C levels (Vrtovcet al., 2005).

Nevertheless, the evidence demonstrating that statins reduce the recurrence of ventricular arrhythmias in CAD patients with ICDs is based on small observational studies or retrospective analyses and data from randomized, placebo-controlled studies were lacking. Very recently, however, two randomized trials support that the addition of statins to ICD device therapy in patients with ischemic and nonischemic diseases reduces life-threatening arrhythmias and improves survival. The *Cholesterol Lowering and Arrhythmia Recurrences after Internal Defibrillator Implantation* (CLARIDI), presented at the Heart Rhythm Society 2006 Annual Scientific Sessions, is a prospective, randomized, placebo-controlled study performed in 106 patients with CAD and life-threatening ventricular arrhythmias requiring an ICD (De Sutter et al., 2006). Forty-five percent of the patients have congestive HF (mean LVEF 39%) and 85% a previous history of acute MI. After 1-year follow-up, atorvastatin therapy (80 mg/daily) results in a significant reduction in the first recurrence (21% vs 38%) of ventricular arrhythmias requiring an appropriate ICD treatment. Moreover, a substudy of the DEFINITE (*Defibrillators in Nonischemic Cardiomyopathy Treatment Evaluation*) trial

demonstrates in ICD patients with nonischemic dilated cardiomyopathy (LVEF = 35%) at risk for ventricular arrhythmias that statin users have lower rates of total mortality and arrhythmic SCD than non statin users, regardless of whether they were randomized to ICD or medical therapy (Goldberger et al., 2006). Compared with no statin use, statin therapy reduces the rate of total mortality/resuscitated cardiac arrest (4.5% vs 18.4%,  $P < .001$ ) and SCD (0.9% vs 5.2%,  $P = 0.043$ ).

## 5.2. Atrial fibrillation

**5.2.1. Effects in experimental models.** Several experimental studies have demonstrated that statins reduce the incidence of AF. In a canine model of AF induced by rapid atrial pacing simvastatin suppresses tachypacing-induced shortening of atrial refractoriness and virtually abolishes AF (Shiroshita-Takeshita et al., 2004). However, these effects are not shared by the antioxidant vitamins C and E. In a canine model of sterile pericarditis, elevated CRP levels are associated with sustained AF, suggesting that electrophysiological changes resulting from inflammation may perpetuate the arrhythmia (Kumagai et al., 2004). In this model, atrial myocytes show active perimyocarditis which consisted of patchy inflammatory infiltration and lipid degeneration and extensive interstitial fibrosis. These inflammatory changes may contribute to atrial structural remodeling and increase the propensity of the arrhythmias to persist. Atorvastatin-treated animals have lower CRP levels, longer atrial refractoriness and shorter intra-atrial conduction times and shorter AF duration than the control group (Kumagai et al., 2004). These results suggest that atorvastatin can prevent atrial electrical and structural remodeling and the maintenance of AF by inhibiting inflammation.

**5.2.2. Effects in clinical studies.** Clinical studies also support the efficacy of statins in maintaining sinus rhythm in patients with persistent AF (Table 1). In a retrospective study performed in 62 patients with lone persistent AF, statin therapy decreases the risk of AF recurrences (42% vs 84%,  $P = 0.032$ ) after successful direct-current cardioversion without affecting the defibrillation threshold (Siu et al., 2003). The beneficial effect of statins is observed during the first few months and remains during the long-term follow-up of 44 months. In another 449 patients with chronic CAD followed by an average of five years, statins significantly reduce the risk of developing AF (8% vs 15%,  $P = 0.01$ ) and this effect is independent of their cholesterol-lowering ability (Young-Xu et al., 2003). This association remains significant after adjustment for potential confounders, including age, hypertension, LV systolic function, HF, acute ischemic events, baseline cholesterol and changes in

cholesterol levels. Interestingly, in this study the incidence of AF is not reduced in patients treated with nonstatin lipid-lowering drugs such as gemfibrozil, cholestyramine, colestipol and probucol. Moreover, there is a relation between length of statin use and reduction of AF and the incidence of AF did not differ between users of nonstatin cholesterol-lowering drugs and nonusers. In another study, paroxysmal AF recorded in ambulatory electrocardiographic monitoring during everyday life was completely resolved in 65% of the patients treated with atorvastatin, but only in 10% of the patients in the placebo group ( $P > 0.019$ ) (Dernellis & Panaretou, 2005). The treatment group has also lower median CRP. In another prospective randomized study performed in 99 patients with persistent AF lasting  $> 48$  hours who were scheduled for electrical cardioversion, those receiving atorvastatin have a higher probability for maintaining sinus rhythm compared with those who receive no drug ( $P = 0.024$ ) (Ozaydin et al., 2006). Finally, in a multicenter registry of 25,268 patients with reduced LVEF ( $\leq 40\%$ ), prevalence of AF was significantly lower in patients taking lipid-lowering drugs (25.1%) compared with patients with untreated hyperlipidemia or with those not having hyperlipidemia (32.6% and 32.8%, respectively,  $P < .001$  for both comparisons; OR 0.69, 95% CI 0.64-0.74) (Hanna et al., 2006). This beneficial impact is independent of the lipid profile and other known arrhythmia risk factors.

Another two studies concluded that statins might help in both primary and secondary prevention AF. The first study found a lower rate of new-onset AF in 667 patients at risk taking statins (7.3% vs 11.6%), an effect restricted to those older than 65 years (Merckx et al., 2004). The other study showed that after successful pharmacologic or direct current cardioversion statin therapy was associated with a lower risk of recurrence of persistent AF among hypertensive patients after 1 year follow-up (50.6% vs 58.5%,  $P = 0.034$ ) (Colivicchi et al., 2004).

However, not all the available evidence supports the benefits of statins in preventing AF recurrence. In an open study performed in 114 patients with AF  $> 48$  h who were scheduled for electrical cardioversion, pravastatin (40 mg/day) for 3 weeks before and 6 weeks after cardioversion does not affect immediate electrical cardioversion success or the recurrence rate of AF after cardioversion (Tveit et al., 2004). The small sample size, the short length of treatment and the duration of AF prior to treatment could play a role in not finding differences between the groups.

AF is a frequent complication after cardiothoracic surgery, occurring in up to 40% of patients. Postoperative AF increases in-hospital morbidity and mortality, length of hospital stay and costs (Fuster et al., 2006). Preoperative use of statins reduces incidence of

postoperative AF following coronary artery bypass grafting (Auer et al., 2004; Marín et al., 2006) and after major noncardiac thoracic (lung and esophageal resection) surgery (Amar et al., 2005). In patients undergoing coronary artery bypass grafting, statin therapy is associated with increased levels of tissue inhibitor matrix metalloproteinase-1 and lower concentrations of matrix metalloproteinase-1 (MMP-1), which suggests that statins could also be protective against AF possibly by inhibiting extracellular matrix remodeling in the atria (Marín et al., 2006). In 253 patients from the SPPAF (*Study for prevention of postoperative atrial fibrillation*) trial, those undergoing cardiac surgery without statin therapy have a significantly higher risk of postoperative AF compared to statin users (45.9% vs 32.8%,  $P < 0.05$ ) (Auer et al., 2004). In patients who have undergone major noncardiac thoracic surgery, the preoperative use of statins is associated with a threefold decrease in the odds of developing AF and this benefit occurs independently of increases in the levels of two markers of inflammation, such as CRP or IL-6 (Amar et al., 2005).

Very recently, the *Atorvastatin for Reduction of Myocardial Dysrhythmias After cardiac surgery* (ARMYDA-3), a randomized, prospective, placebo-controlled trial, studied the effects of atorvastatin in 200 patients undergoing elective cardiac surgery with cardiopulmonary bypass (Patti et al., 2006). Treatment with atorvastatin, starting 7 days before cardiac surgery and continued in the postoperative period, significantly decreases postoperative AF (35% vs 57%) and shortens the postoperative hospital stay (6.3 vs 6.9 days,  $P = 0.001$ ). Subgroups analysis show that atorvastatin treatment results in a lower risk of AF in patients irrespective of age, sex, presence of diabetes mellitus, hypertension, and chronic obstructive pulmonary disease. Treatment benefit was more evident in patients undergoing coronary bypass operations and in those with a normal-sized left atrium, whereas it was absent after noncoronary surgery and in patients with left atrial enlargement.

## **6. Mechanisms of action involved in the antiarrhythmic effects of statins**

### **6.1. Antiischemic effects**

The mechanisms by which statins exert their antiarrhythmic effects have not yet been characterized at the cellular level. It can be hypothesized that in patients with CAD the antiarrhythmic effects and the reduction in SCD observed in secondary prevention trials may simply reflect their ability to lower the LDL-C levels, thereby attenuating the progression of CAD and, in some patients, promoting the regression of coronary atherosclerotic plaques. As a result, statins would reduce the number of new ischemic episodes that could have

triggered the arrhythmogenic substrate to initiate and/or maintain both atrial and ventricular tachyarrhythmias.

Statins exert cardioprotective effects in an isolated perfused rat heart model of global myocardial ischemia and reperfusion in normocholesterolemic, hypercholesterolemic and diabetic animal models (Lefer et al., 1999; Wagner et al., 2000; Lefer et al., 2001; Chen et al., 2003) and in apolipoprotein E-deficient (apoE<sup>-/-</sup>) mice fed with a high-cholesterol diet (Scalia et al., 2001). In these models, statins improve coronary blood flow, preserve LV developed pressure and cardiac contractile function, reduce the extent of myocardial necrosis, the leukocyte–endothelial cell interaction and the polymorphonuclear leukocytes accumulation in the ischemic myocardium and increase the NO production in the vasculature. Moreover, statins reduce transient ischemia even in patients with optimal anti-anginal therapy (Van Boven et al., 1996) and in patients with acute coronary syndrome, early and aggressive, lipid-lowering therapy with atorvastatin (80 mg/d) reduces recurrent ischemic events, mostly those requiring rehospitalization (Schwartz et al., 2001; Colivicchi et al., 2002).

Infarct size is markedly limited when brief periods of ischemia precede the sustained ischemia, a phenomenon known as ischemic preconditioning. Hypercholesterolemia blunted the infarct size-limiting effect of ischemic preconditioning and pravastatin restores the infarct size-limiting effect of ischemic preconditioning possibly via restoring the activation of ecto-5'-nucleotidase activity which, in turn, increases interstitial adenosine release (Ueda et al., 1999). Moreover, pretreatment with pravastatin enhances the adaptation of ischemia during coronary angioplasty, an effect abolished by aminophylline, which suggests that its cardioprotective effect may result from activation of adenosine receptors (Lee et al., 2004).

Furthermore, statins reduce transient ischemia even in patients with optimal anti-anginal therapy (Van Boven et al., 1996) and in patients with acute coronary syndrome, early and aggressive therapy with atorvastatin (80 mg/d) reduces recurrent ischemic events, mostly those requiring rehospitalization (Schwartz et al., 2001; Colivicchi et al., 2002).

Several lines of evidence, however, indicate that the antiarrhythmic effects of statins are independent of their LDL-C lowering effects. 1) These antiarrhythmic effects are observed both in patients with normocholesterolemia and/or hypercholesterolemia and with or without CAD. 2) Ischemia is rarely present in patients with recurrent ventricular tachyarrhythmias before the onset of recorded spontaneous sustained VT. Moreover, ventricular tachyarrhythmias can often be initiated by programmed electrical stimulation in the absence of ischemia and coronary revascularization does not prevent recurrent ventricular

arrhythmias in the chronic phase of MI (Anderson, 2003). 3) The magnitude of the survival benefit associated with statin therapy in patients with propensity to life-threatening VT/VF is higher than that reported from mortality trials in patients without such a demonstrated propensity (Scandinavian Survival Study Group, 1994; The LIPID Study Group, 1998). 4) The antiarrhythmic effect occurs relatively early in the follow-up. In fact, the Kaplan-Meier curves of arrhythmia-free survival for patients with and without statin therapy started to diverge very early (within 1 to 2 months of follow-up), long before any angiographically measurable regression of atherosclerosis (Mitchell et al., 2003; Vyas et al., 2006). Thus, reversal of atherosclerosis and/or retardation of plaque development associated could not be the sole reason for the antiarrhythmic actions of statins observed in patients with CAD and the benefits of statins extent to mechanisms beyond their LDL-C lowering effects, i.e. the so-called pleiotropic effects (Table 3).

## **6.2. Role of pleiotropic effects on the antiarrhythmic effects of statins**

The molecular mechanisms of the pleiotropic effects of statins have been analyzed in detail previously (Bellosta et al., 1998; Bonetti et al., 2002; Davignon, 2004; Liao & Laufs, 2005), in this article we shall focus our interest in the cardiac pleiotropic effects of statins that may be involved in their antiarrhythmic actions.

**6.2.1. Anti-inflammatory effects.** Plasma CRP are elevated in patients at risk for future cardiovascular events (Ridker, 2001a). Statins exhibit potent antiinflammatory effects reducing the levels of the hs-CRP in experimental models of AF (Kumagai et al., 2004) and in patients with combined hyperlipidemia and with CAD (Jialal et al., 2001; Ridker et al., 2001b; Albert et al., 2001). In the Cholesterol and Recurrent Events (CARE) trial, risk reduction in patients with CAD receiving statins was greater in those with higher CRP levels for a similar degree of LDL-C lowering (Ridker et al., 1999, 2001a). These results suggest that statin therapy may improve the inflammatory arrhythmogenic substrate in patients with both supraventricular and ventricular arrhythmias. Moreover, this antiinflammatory effect could be part of the mechanism of the stabilization of atherosclerotic plaques.

**6.2.2. Improvement of endothelial dysfunction.** Clinical studies provide evidence for a close association between the degree of endothelial dysfunction and clinical cardiovascular events in patients with cardiovascular risk factors, CAD or HF (Landmesser & Drexler, 2005). Statins improve coronary endothelium dysfunction and myocardial perfusion and preserve coronary microvascular integrity in hypercholesterolemic patients with CAD (Wassman et al., 2002a,b; Liao & Laufs, 2005; Landmesser & Drexler, 2005). High-doses

of atorvastatin improve peripheral endothelial function within 24 hours but this effect disappears 1 day after discontinuation of the drug (Laufs et al., 2001). This improvement occurred before levels of LDL-C and hs-CRP decrease after 2 days of treatment, suggesting that the beneficial effects of statins are independent of LDL-C lowering or the modulation of the inflammatory state. Improvement of coronary endothelial dysfunction and myocardial perfusion would be expected to reduce the ischemic arrhythmogenic substrate.

**6.2.3. Stabilization of vulnerable plaques.** Atherosclerotic plaque rupture underlies the onset of around 70% of acute coronary syndromes, so that modulation of plaque instability would reduce the incidence of acute coronary events and the ischemic arrhythmogenic substrate. Statins reduce the lipid core and lipid-laden macrophages and increase the relative volume of collagen within the plaque (Bellosta et al., 1998; Williams et al., 1998; Sukhova et al., 2002; Aikawa et al., 2001). They also inhibit monocyte infiltration (Bustos et al., 1998) and the activation of the monocyte/macrophage system reducing the release of proinflammatory cytokines and MMPs within the plaque (Bellosta et al., 1998; Luan et al., 2003). In patients with symptomatic carotid artery stenosis, plaques removed by endarterectomy from the statin-treated patients present significantly less lipid and oxidized LDL immunoreactivity, fewer macrophages and T cells and less MMP-2 immunoreactivity, and cell apoptosis than control plaques (Crisby et al., 2001). In addition, plaques had higher collagen content and immunoreactivity to tissue inhibitor of metalloproteinase-1 (a potent inhibitor of MMP-1 and MMP-9) in the pravastatin group. These effects, which occurred independently of LDL-C levels, indicate that statin therapy is associated with changes in atherosclerotic plaque composition that favor lesion stability and reduce the risk of rupture and of acute coronary events (Bellosta et al., 1998; Bonetti et al., 2002; Luan et al., 2003; Liao & Laufs, 2005).

**6.2.4. Antioxidant effects.** Formation of the oxygen-derived free radical superoxide by NAD(P)H oxidases plays a critical role in the development of a wide range of cardiovascular diseases, including AF (Kim et al., 2005). Reactive oxygen species may alter channel kinetics, including a decrease in Na<sup>+</sup> channel availability (Fukuda et al., 2005) and increasing the amplitude of the ultrarapid component of the delayed rectifier K<sup>+</sup> current (I<sub>Kur</sub>) (Caouette et al., 2003), leading to slowing of conduction velocity and shortening of the cardiac action potential duration, respectively, effects that facilitate the initiation and/or maintenance of AF. Moreover, abnormal activation NAD(P)H oxidases in response to neurohormones (angiotensin II, norepinephrine) has been shown to contribute to cardiac myocyte hypertrophy and fibrosis (Sorescu & Griendling, 2002).

Statins reduce the NADPH oxidase-dependent production of reactive oxygen species (superoxide and hydroxy radicals) by activated macrophages (Suzumura et al., 1999; Rikitake et al., 2001), endothelial cells (Wagner et al., 2000; Feron et al., 2001; Wassmann et al., 2002a) and isolated human atrial preparations obtained during cardiac bypass surgery (Maack et al., 2003). Moreover, statins enhance the activity of antioxidant enzymes such as catalase, glutathione peroxidase and paraoxonase (Wagner et al., 2000; Feron et al., 2001; Dechend et al., 2001; Wassmann et al., 2002a). However, the antioxidant effects of most statins appear only at suprapharmacological (micromolar) concentrations, so that its physiological relevance is uncertain. Only the active O-hydroxy metabolite of atorvastatin, but not the parent compound, exerts potent antioxidant effects inhibiting LDL oxidation at pharmacological (nanomolar) concentrations, being more potent than vitamin E or probucol (Mason et al., 2004).

**6.2.5. Cardioprotective effects.** Epidemiological studies found that hypercholesterolemia is associated with LV hypertrophy (Sundstrom et al., 2001), a major independent risk factor for total and cardiovascular mortality and SCD (Levy et al., 1990). Myocardial infarction frequently produces alterations in the geometry and structure (dilatation, hypertrophy, interstitial fibrosis) of noninfarcted ventricular myocardium, referred to as remodeling, that contributes to the development of depressed cardiac performance and is an important component of the arrhythmogenic substrate in post-MI patients (Pffefer y Braunwald, 1990). Statins inhibit cardiac hypertrophy induced in normotensive rats by aortic binding (Takemoto et al., 2001; Indolfi et al., 2002) or angiotensin II (AII) infusion (Oi et al., 1999), spontaneously hypertensive rats (Lee et al., 2005b), hypercholesterolemic rabbits (Lee et al., 2005a) and transgenic rats for human renin and angiotensinogen (dTGR) (Dechend et al., 2001). In hyperlipidemic patients, pravastatin reduces LV mass and the LV mass regression correlates with the magnitude of inhibition of free radical formation assessed by 8-iso-prostaglandin F<sub>2</sub> $\alpha$  formation (Lee et al., 2002). These antiproliferative effects are independent of blood pressure or LDL-C reduction but are reversed by mevalonate, suggesting that they are related to the blockade of the synthesis of isoprenoids.

Since CAD represents the most common cause of chronic HF, it may be not a surprise that statins may reduce the risk of developing HF after MI. In a normocholesterolemic murine model of congestive HF after MI, statin therapy normalizes the sympathetic outflow, reduces LV dilatation and LV end-diastolic pressure, cardiac hypertrophy and interstitial fibrosis of the noninfarcted LV without affecting the infarct size, improves LV ejection fraction and reduces death due to HF and arrhythmias (Dechend et al., 2001; Bauerssachs et

al., 2001; Hayashidani et al., 2002). In normolipidemic rats, pravastatin administration after MI attenuates cardiomyocyte hypertrophy and reduces the inducibility of ventricular tachyarrhythmias (Lee et al., 2003). It is important to note that although in these models HF was induced by an initial episode of ischemia and infarction, ongoing pathological ventricular remodeling is unrelated to further ischemic events on the failing ventricle. Thus, the beneficial effect of statins is unrelated to their impact on subsequent ischemia (Krum & McMurray, 2002). Neurohumoral activation plays a central role in the pathogenesis of LV hypertrophy and HF and statins inhibit the expression of endothelin-1, a potent vasoconstrictor and mitogen, and beta-adrenergic receptor-stimulated apoptosis in rat ventricular myocytes and decrease AT1 receptors (Laufs et al., 2006) and exert a favorable modulation of the autonomic nervous system (see 6.2.6.).

Even when HF patients were excluded in large statin trials and LVEF was either not measured or studied in only a small number of patients, post hoc analyses of statin trials have shown their beneficial effects in patients with ischemic HF (Sacks et al., 1996; Kjekshus et al., 1997; Krum & McMurray, 2002; Raina et al., 2006). Moreover, retrospective analyses of statin use in patients enrolled in HF clinical trials show that statin use is associated with improved survival and less development of HF after MI (Raina et al., 2006). Several studies have recently reported that in patients with normal (Fukuta et al., 2005, Folkeringa et al., 2006) or reduced LVEF ( $< 0.40$ ) (Huikuri et al., 1999; Christensen et al., 1999; Horwich et al., 2004; Ray et al., 2005; Sola et al., 2005, 2006), statin therapy improves exercise tolerance and LVEF, lowers NYHA class, reduces markers of inflammation (hs-CRP; IL-6 and TNF $\alpha$ ) and decreases hospitalizations for HF and all-cause mortality. There are no difference in survival among patients treated with statins who had ischemic or nonischemic HF. A similar benefit was observed in patients with nonischemic idiopathic dilated cardiomyopathy (Node et al., 2003).

Since LV hypertrophy and remodeling play an important role in the arrhythmogenic substrate that initiates and/or maintains cardiac arrhythmias in patients with hypertension, CAD and HF, it would be expected that statin therapy would reduce the arrhythmogenic risk in these populations. The possible putative mechanisms to explain the beneficial effects of statins in HF patients include: prevention of ischemic coronary events through LDL-C lowering, improvement of microvascular circulation and endothelial function, plaque stabilization, reduction of cardiac hypertrophy and fibrosis, neurohumoral activation and ventricular wall stress, and to their antioxidant, antiinflammatory and direct antiarrhythmic effects (Krum & McMurray, 2002; Laufs et al., 2006).

**6.2.6. Improvement of autonomic function.** We already mention that sympathetic activity and sympathetic hyperinnervation participate in the occurrence of ventricular tachyarrhythmias and SCD in post-MI patients (Cao et al., 2000a,b; Rubart & Zipes, 2005). Low heart rate variability (HRV), which probably reflects a decrease in vagal tone and/or an increase in sympathetic tone, is associated with an increased risk of coronary events, all-cause mortality and susceptibility to ventricular arrhythmias in patients with CAD and is negatively affected by hypercholesterolaemia (Quintana et al., 1997; Huikuri, 1999). In apoE<sup>-/-</sup> mice (Pelat et al., 2003) and in a rabbit model of pacing-induced congestive HF (Pliquett et al., 2003), statins normalize autonomic sympathetic outflow and baroreceptor responses, decrease plasma noradrenaline levels and renal sympathetic nerve activity and increase HRV. Hypercholesterolemia is also associated with a decreased HRV in men with and without CAD, suggesting an increased risk of SCD (Christensen et al., 1999).

In patients with hypercholesterolemia pravastatin increases parasympathetic modulation of heart rate, an effect associated with an increased expression of the  $\alpha$ -subunit of the heterotrimeric G-protein, G $_{\alpha i2}$  (Welzig et al., 2003). Atorvastatin improves autonomic function, as reflected by an increase in HRV level in hypercholesterolemic patients, with or without CAD (Pehlivanidis et al., 2001), and long-term administration of pravastatin in patients with stable angina pectoris (Van Boven et al., 1995). Statins also increase HRV in patients with previous MI with or without HF and this effect is independent from changes in plasma LDL-C levels (Riahi et al., 2002). Moreover, short-term treatment with atorvastatin increases HRV, decreases QT interval variability and shortens the QTc interval duration in patients with advanced chronic HF (mean LV ejection fraction 24% and NYHA class III) (Vrtovec et al., 2005). Thus, statin therapy exerts a favourable modulation on the sympathetic tone and autonomic dysfunction and may, therefore, reduce the risk of life-threatening arrhythmias in patients with CAD and HF.

### **6.2.7. Effects on cardiac ion channels and transporters**

**6.2.7.1. Role of membrane cholesterol on transmembrane ion fluxes.** Free (unsterified) cholesterol is a major lipid class of mammalian cell membranes where it plays an essential role in determining their physico-chemical properties and directly affects transmembrane ion fluxes by modulating the normal function of constituent membrane-integral proteins, like ion channels and ion pumps (Quinn, 1981; Tulenko et al., 1990; Yeagle, 1991).

Lipids are not homogeneously distributed within the lipid bilayer, but spontaneously aggregate to form specialized membrane microdomains enriched in cholesterol and sphingolipids (the so called “lipid rafts”), where many receptors and channels involved in

cell signalling preferentially accumulate. In SMCs, channels associated with lipid raft microdomains included voltage-gated  $\text{Na}^+$ , L-type  $\text{Ca}^{2+}$  and several  $\text{K}^+$  channels, including several voltage-gated (Kv1.3, Kv1.5 and Kv2.1) and the large conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels and the inward rectifier  $\text{K}^+$  channel subunit Kir3.1 (Martens et al., 2004).

Excess of membrane cholesterol modifies the physical properties of the membrane and significantly influences transmembrane ion fluxes. In fact, enrichment of human erythrocyte and cardiac membranes and rabbit kidney medulla basolateral membrane vesicles with cholesterol increases membrane core microviscosity and stimulates  $\text{Na}^+$ - $\text{Ca}^{2+}$  exchange, while inhibits the activity of  $\text{Na}^+$ - $\text{K}^+$ -ATPase,  $\text{Ca}^{2+}$ -ATPase,  $\text{Mg}^{2+}$ -ATPase and  $\text{Na}^+$ ,  $\text{Li}^+$ -countertransport (Ortega & Mas-Oliva, 1984; Kutryk & Pierce, 1988; Yeagle et al., 1988; Lijnen & Petrov, 1995).

Cholesterol enrichment of vascular SMC membranes with human LDL increases unstimulated and noradrenaline-stimulated  $\text{Ca}^{2+}$  influx, unstimulated  $\text{Ca}^{2+}$  efflux and cytosolic  $\text{Ca}^{2+}$  levels (Bialecki & Tulenko, 1989; Gleason et al., 1991). These effects that are associated with a decrease in membrane fluidity, are inhibited by L-type  $\text{Ca}^{2+}$  channel blockers (nifedipine, verapamil and diltiazem), suggesting that excess membrane cholesterol may unmask otherwise silent L-type  $\text{Ca}^{2+}$  channels that may be involved in altered arterial wall properties in dyslipemic patients. A significant increase in  $^{45}\text{Ca}^{2+}$  influx is also found in erythrocytes enriched with cholesterol (Locher et al., 1984). Moreover, vasosensitivity to noradrenaline and angiotensin II increases in hypercholesterolemic patients (Howes et al., 1997) and in isolated cholesterol-enriched arteries (Bialecki & Tulenko, 1989).

Hypercholesterolemia increases free cholesterol (and cholesterol ester) in cardiac sarcolemma and decreases  $\text{Na}^+$  current ( $I_{\text{Na}}$ ) density in isolated rabbit ventricular myocytes, an effect associated with a leftward shift in the inactivation potential and a slowing of the time course of recovery (Wu et al., 1995). However,  $I_{\text{Ca,L}}$  density is increased in hypercholesterolemic rabbits, an effect that may contribute to the prolongation of the ventricular APD (QT intervals) observed in this model (Wu et al., 1995) and in canine and ovine Purkinje fibers (Karagueuzian et al., 1982). In Langendorff-perfused rabbit hearts, hyperlipidemia also prolonged the QT interval and this effect was reversed by simvastatin (Lee et al., 2005a). In contrast, cholesterol-depleted rat heart myocytes in culture manifest a faster rate of depolarization and their spontaneous activity was more resistant to  $[\text{Na}^+]$  or  $[\text{Ca}^{2+}]$  deprivation, low pH, verapamil and tetrodotoxin (Hasin et al., 1980). These findings are consistent with an increase in  $\text{Na}^+$  and  $\text{Ca}^{2+}$  currents during depolarization.

Excess membrane cholesterol also increases basal and noradrenaline-activated  $^{86}\text{Rb}$  efflux through  $\text{K}^+$  channels in cultured aortic SMCs (Tulenکو et al., 1990). In vascular SMCs, hypercholesterolemia regulates the gene expression levels of several  $\text{K}^+$  channels, upregulating Kir6.2 mRNA expression (the pore-forming  $\alpha$ -subunit of cardiac ATP-sensitive  $\text{K}^+$  ( $\text{K}_{\text{ATP}}$ ) channels) and downregulating Kir3.1 mRNA expression (the  $\alpha$ -subunit of a G-protein coupled  $\text{K}^+$  channel), while Kir2.1 (inward rectifier  $\text{K}^+$  channel), Kir6.1 and SUR2B ( $\text{K}_{\text{ATP}}$   $\beta$ -subunits) mRNA expression are unchanged (Ren et al., 2001). Enrichment of bovine aortic endothelial cells with cholesterol decreases the inward rectifier  $\text{K}^+$  (Kir2) current density, whereas depleting the cells of cholesterol increases the density of the current (Romanenko et al., 2002). Since neither single channel properties nor membrane capacitance are affected, cholesterol modulates  $\text{K}^+$  conductance by changing the number of active channels. Moreover, increasing the cholesterol concentration reduces (70-80%) the mean open time and the open probability (80%) of the high-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel (BK) reconstituted into lipid bilayers from rat brain homogenates (Chang et al., 1995). These data are consistent with a model in which an increase in cholesterol concentration in the cell membrane increases intrabilayer structural stress and lateral elastic stress energy (Chang et al., 1995; Mason et al., 2004). As a result, channel opening events may be compromised resulting in a reduced probability of finding the channel in the open state, i.e, cholesterol affects the transitions between different conformational states of the channel protein (Chang et al., 1995). Hypercholesterolemia also impairs acidosis-induced dilation of rabbit coronary arterioles, an effect mediated via the activation of  $\text{K}_{\text{ATP}}$  channels (Shioiri et al., 2003). Interestingly, cholesterol depletion produces a marked hyperpolarization shift in the steady-state inactivation properties of Kv2.1 channels and alters Kv1.5 channel function by inducing a hyperpolarizing shift in the voltage dependence of activation and inactivation (Martens et al., 2004). In addition, it produced a dramatic redistribution of Kv2.1 channels from small clusters into large patches, while surface distribution of Kv1.4 and Kv1.3 was not altered (O'Connell & Tamkun, 2005).

**6.2.7.2. Role of statins on transmembrane ion fluxes.** From the previous results, it seems clear that, by reducing cholesterol content in cardiac membranes, statins could modulate the fatty acid composition and the physicochemical properties of the sarcolemma and regulate the activity of membrane integral proteins. In addition, statins could modulate ion channels through other mechanisms independent on the inhibition of cholesterol synthesis and even interacting directly with the channel proteins similarly to conventional antiarrhythmic drugs.

Independently of the mechanism involved, changes in transmembrane ion fluxes induced by statins may affect the electrophysiological properties of the cardiac muscle (Lamers et al., 1987; Pound et al., 2001)

In hypercholesterolemic patients, pravastatin reduces the erythrocyte and platelet membrane cholesterol content and this effect is accompanied by an increase in activity of the  $\text{Na}^+\text{-K}^+$  pump and a decrease in cellular  $\text{Na}^+$  concentration (Lijnen et al., 1994). Simvastatin and atorvastatin, but not pravastatin, inhibit the contractions induced by phenylephrine in isolated aorta from normocholesterolemic rats incubated in normal and in  $\text{Ca}^{2+}$ -free media as well as the increase in  $[\text{Ca}^{2+}]_i$  induced by angiotensin II in cultured rat aortic SMCs (Tesfamariam et al., 1999). These effects suggest that simvastatin and atorvastatin directly relax VSMCs by inhibiting  $\text{Ca}^{2+}$  influx via voltage-gated channels and  $\text{Ca}^{2+}$  release from intracellular store sites. Very recently, Vaquero et al (2006) have demonstrated that at the nanomolar range simvastatin, but not atorvastatin, inhibits the  $I_{\text{Ca,L}}$  recorded in mouse ventricular myocytes (Figure 2). The inhibition was frequency-dependent and displayed fast onset and washout kinetics (3-6 minutes), suggesting that it was the consequence of the direct interaction of the drug with the channel proteins. Thus, simvastatin at the same concentration at which it inhibits the HMG-CoA reductase, it also produces a marked frequency-dependent block of the  $I_{\text{Ca,L}}$  (40% at 1 Hz).

Experimental studies demonstrated that, simvastatin prevented AF promotion in dogs by attenuating the APD and refractoriness abbreviation and the downregulation of the  $\text{Ca}_v1.2\alpha$ -subunit expression produced after 7 days of atrial tachypacing (Shiroshita-Takeshita et al., 2004). It has been proposed that the control of intracellular  $\text{Ca}^{2+}$ , which is likely altered in the face of rapid rates, would help to prevent the  $\text{Ca}^{2+}$ -dependent gene transcription responsible for the electrical remodeling (Nattel, 2002). Therefore, it is possible that the frequency-dependent reduction in  $I_{\text{Ca,L}}$  produced by SVA might be responsible for the attenuation in  $\text{Ca}_v1.2$  downregulation and for the prevention of the electrical remodeling observed with simvastatin in the dog model of AF (Shiroshita-Takeshita et al., 2004).

Moreover, simvastatin dose-dependently delays anoxia-induced necrosis of cultured neonatal cardiomyocytes and attenuates the rise of  $[\text{Ca}^{2+}]_i$  upon incubation in  $\text{Na}^+$ -free buffer (Bastiaanse et al., 1994). These protective effects are attributed to the inhibition of the sarcolemmal  $\text{Na}^+/\text{Ca}^{2+}$  exchanger and of the rise in  $[\text{Ca}^{2+}]_i$  during anoxia (Bastiaanse et al., 1994). L- $\alpha$ -lysophosphatidylcholine (LPC) is a metabolite of phosphatidylcholine found in high concentrations in ischemic hearts (Sobel et al., 1978). In isolated cardiac myocytes LPC increases  $[\text{Ca}^{2+}]_i$  and is thought to be one of the causes of  $\text{Ca}^{2+}$  overload and

arrhythmias during ischemia and reperfusion (Ver Donk et al., 1992). LPC induces a nonselective cation current ( $I_{NSC}$ ) in human endothelial cells and guinea pig ventricular myocytes via a Gi/Go-coupled receptor and Rho-mediated pathway (Li et al., 2002). Fluvastatin prevents the GTP exchange of RhoA and its membrane translocation from the cytoplasm caused by LPC and inhibits the increase in  $[Ca^{2+}]_i$  in human endothelial cells (Yokoyama et al., 2002) and the  $I_{NSC}$  current induced by LPC in guinea pig ventricular myocytes (Li et al., 2002). The effects of fluvastatin are reversed by GGPP or mevalonic acid.

Statins also modulate  $K^+$  currents. In smooth muscle rabbit aortic cells, lovastatin produces to a nine-fold increase in the open probability of the  $Ca^{2+}$ -activated  $K^+$  channels presumably due to changes in plasma membrane fluidity, even when the elementary conductance of the channels does not change (Bolotina et al., 1989). In rats treated with the NOS inhibitor L-nitro-arginine methyl ester (L-NAME) for 4 weeks, pitavastatin inhibits the rise in systolic blood pressure and microvascular remodeling (medial thickening and perivascular fibrosis of coronary arterioles), these effects being independent of cholesterol lowering; in addition, pitavastatin abolishes the upregulation of intermediate-conductance  $Ca^{2+}$ -activated  $K^+$  channels (ImK) in vascular SMCs (Terata et al., 2003). Cerivastatin and fluvastatin enhance endothelium-dependent relaxations in the rat aorta, an effect that is abolished by the NOS inhibitor  $N^G$ -nitro-L-arginine and a PI3 kinase inhibitor (Mukai et al., 2003). At higher concentrations, both lipophilic statins, but not pravastatin, produce an endothelium-independent relaxation that is inhibited by 4-aminopyridine, a voltage-gated  $K^+$  ( $K_v$ ) channel blocker. Since  $K^+$  channels play a role in the regulation of VSMC membrane potential, opening of  $K_v$  channels would be expected to cause membrane hyperpolarization that closes voltage-gated L-type  $Ca^{2+}$  channels, leading to smooth muscle relaxation. These results indicate that multiple mechanisms are involved in the vasodilator effects of statins, including augmentation of NO-dependent relaxations through the PI3 kinase/Akt pathway and endothelium-independent mechanisms by opening  $K_v$  channels (cerivastatin and fluvastatin) or inhibiting  $Ca^{2+}$  influx (simvastatin and atorvastatin) (Furukawa et al., 1991). Opening of  $K_{ATP}$  channels, particularly mitochondrial  $K_{ATP}$  (mito $K_{ATP}$ ), is a primary mechanism of cardioprotection against ischemia (Seino & Miki, 2003; Tamargo et al., 2004b). Suppression of  $K_{ATP}$  channel opening during ischemia/reperfusion may underlie the hypercholesterolemia-induced extension of no-reflow, which delays infarct healing. NO directly activates mito $K_{ATP}$  channels (Sasaki et al., 2000). Thus, the cardioprotective effects of statins against ischemic injury can be related to the activation of  $K_{ATP}$  channels, an

effect that can be indirectly mediated by the release of NO or by increasing adenosine levels (Lee et al., 2004), an important mediator to trigger opening of  $K_{ATP}$  channels. In a model of global ischemia in rabbit hearts pravastatin exhibits a protective effect on cardiac metabolism during ischemia, inhibiting the decrease in ATP and intracellular pH and its effects are suppressed with the  $K_{ATP}$  channel blocker glibenclamide and L-NAME (Kawabata et al., 2001). These results suggest that the cardioprotection of pravastatin may be caused by the  $K_{ATP}$  channels that are mediated by the release of NO. In fact, pravastatin opens cardiac  $K_{ATP}$  channels, through the activation of the NOS3 and its cardioprotective effects were inhibited by glibenclamide, suggesting that activation of  $K_{ATP}$  channel is involved (Lee et al., 2004).

On the other hand, at nanomolar concentrations, atorvastatin, but not simvastatin, inhibited the atrioselective ultrarapid component of the delayed rectifier  $K^+$  current ( $I_{Kur}$ ) and the rapidly activating and inactivating transient outward current ( $I_{to}$ ) in mouse ventricular myocytes (Gómez et al, 2005; Vaquero et al, 2005). The blockade induced was concentration-, voltage- and frequency-dependent and was not modified by the antioxidant probucol or L-NAME. These results suggest that neither the antioxidant actions nor NO are involved in the effects of both statins. As a consequence of its effects on these  $K^+$  currents, atorvastatin prolonged the action potential duration in mouse atria, without modifying the amplitude and maximum upstroke velocity ( $V_{max}$ ) of the action potential or the resting membrane potential. These results suggest that atorvastatin does not modify the  $I_{Na}$  or the inward rectifier  $K^+$  ( $I_{K1}$ ) currents. In these experiments, atorvastatin-induced block also displays fast onset and washout kinetics and the drug does not modify the membrane capacitance, indicating that the effects are due to a direct interaction of the drug with the channel proteins without the implication of the inhibition in membrane cholesterol synthesis or putative changes in the membrane properties. It has been proposed that  $I_{to}/I_{Kur}$  block could be a promising target for the treatment of atrial fibrillation (Wijffels & Crijns, 2003). Indeed, in the human atria, the maintenance of the duration and the height of the plateau phase not only determines the excitation-contraction coupling, but is critical for the prevention of the occurrence of the periodic activity of sustained, high frequency, functional reentrant sources (rotors) that can be responsible for the AF (Pandit et al., 2005). Moreover, it has been demonstrated that the combined inhibition of  $I_{Kur}$  and  $I_{to}$  prolonged the atrial plateau duration and is critical for the termination of rotors. Therefore, atorvastatin could produce a cessation of fast reentrant sources by inhibiting the  $I_{Kur}$  and  $I_{to}$ . Therefore, atorvastatin and simvastatin may modify human atrial plateau currents, producing a mild

prolongation of the plateau phase, in the case of atorvastatin, and a decrease in the  $\text{Ca}^{2+}$  entry into the myocyte, in the case of SVA. However, it is difficult to assert whether these effects may contribute to the prevention of development and recurrence of AF since they appeared at concentrations somewhat higher than their free plasma concentrations.

Lovastatin also inhibits the rapid component of the delayed rectifier ( $I_{Kr}$ ) in stably transfected human embryonic kidney cells in a dose-dependent manner ( $\text{IC}_{50} = 7.4 \mu\text{M}$ ) that showed no use dependence (Katchman et al., 2006). However, lovastatin had no appreciable effect on either action potential duration or  $I_{Kr}$  in guinea pig ventricular myocytes.

## 7. Conclusions

In recent years, experimental and clinical evidence have demonstrated that statin therapy exerts antiarrhythmic effects. Clinical trials have shown that statins reduce the recurrences of AF as well as the episodes of recurrent VT and VF in patients with CAD treated with an ICD. In addition, some results from secondary prevention trials indicated that statins might reduce SCD associated to lethal ventricular arrhythmias. Interestingly, these antiarrhythmic effects have not been observed in some studies with other lipid-lowering drugs. Although the mechanisms underlying their antiarrhythmic properties are unknown, they can be attributed to their lipid-lowering effects that reduced the ischemic burden, to their pleioropic actions (including improvement of endothelial dysfunction and autonomic function, stabilization of the atherosclerotic plaques and antioxidant, anti-inflammatory, antithrombotic and cardioprotective properties) and to direct effects on transmembrane ion fluxes. However, further investigations are needed to clarify the potentially underlying mechanisms involved in the antiarrhythmic effects of statins and more prospective studies are necessary to assess the risks and benefits of routine statin therapy in clinical practice, particularly in high-risk CAD patients

## Acknowledgements

This work was supported by CICYT (SAF2005-04609), Red HERACLES and Mutua Madrileña Grants

**Figure legends**

**Figure 1.** Molecular pathway for cholesterol synthesis. Inhibition of 3-hydroxy-3-methyl glutaryl coenzyme A (HMG CoA) reductase inhibits cholesterol synthesis and isoprenoid production. BMP-2: bone morphogenetic protein. NADPH: nicotinamide adenine dinucleotide phosphate. PP: pyrophosphate. NOS3: endothelial nitric oxide synthase. PAI-1: plasminogen activator inhibitor-1. PI3: phosphatidylinositol 3-kinase/protein. t-PA: tissue-type plasminogen activator.

**Figure 2.** Calcium currents recorded in mouse ventricular myocytes by applying trains of 16 150 ms-pulses from  $-80$  to  $0$  mV (with a prepulse of 50 ms to  $-30$  mV to inactivate the  $I_{Na}$ ) at a frequency of stimulation of 1 Hz under control conditions and in the presence of 10 nM atorvastatin (panel A) and 10 nM of the acid form of simvastatin (SVA, panel B)

## References

- Aikawa, M., Rabkin, E., Sugiyama, S., Voglic, S. J., Fukumoto, Y., Furukawa, Y., et al. (2001). An HMG-CoA reductase inhibitor, cerivastatin, suppresses growth of macrophages expressing matrix metalloproteinases, and tissue factor *in vivo* and *in vitro*. *Circulation* 103, 276-283.
- Albert, M. A., Danielson, E., Rifai, N., & Ridker, P.M; PRINCE Investigators. (2001). Effect of statin therapy on C-reactive protein levels: the pravastatin inflammation/CRP evaluation (PRINCE): a randomized trial and cohort study. *JAMA* 286, 64-70.
- Albarte, C., & Zipes, D. P. (2003). Use of nonantiarrhythmic drugs for prevention of sudden cardiac death. *J Cardiovasc Electrophysiol* 14(9 Suppl), S87-S95.
- Amar, D., Zhang, H., Heerdt, P. M., Park, B., Fleisher, M., & Thaler, H. T. (2005). Statin use is associated with a reduction in atrial fibrillation after noncardiac thoracic surgery independent of C-reactive protein. *Chest* 128, 3421-3427.
- Anderson, K. P. (2003). Lipid-lowering therapy for prevention of ventricular tachyarrhythmias. *J Am Coll Cardiol* 42, 88-92.
- Anderson, J. L., Allen Maycock, C. A., Lappe, D. L., Crandall, B. G., Horne, B. D., Bair, T. L., et al. (2004). Frequency of elevation of C-reactive protein in atrial fibrillation. *Am J Cardiol* 94, 1255-1259.
- Auer, J., Weber, T., Berent, R., Lamm, G., Ng, C. K., Hartt, P., et al. (2004). Use of Hmg-coenzyme a-reductase inhibitors (statins) and risk reduction of atrial fibrillation after cardiac surgery: results of the SPPAF study: a randomised placebo-controlled-trial. *Eur Heart J* 25(Suppl), 353.
- Avilés, R. J., Martín, D. O., Apperson-Hansen, C., Hoghtaling, P. L., Rautaharju, P., Kronmal, R. A., et al. (2003). Inflammation as a risk factor for atrial fibrillation. *Circulation* 108, 3006-3010.
- Bastiaanse, E. M., Atsma, D. E., Kuijpers, M. M., & Van der Laarse, A. (1994). Simvastatin-sodium delays cell death of anoxic cardiomyocytes by inhibition of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *FEBS Lett* 343, 151-154.
- Bauersachs, J., Galuppo, P., Fraccarollo, D., Christ, M., & Ertl, G. (2001). Improvement of left ventricular remodeling and function by hydroxymethylglutaryl coenzyme a reductase inhibition with cerivastatin in rats with heart failure after myocardial infarction. *Circulation* 104, 982-985.

- Bellosta, S., Via, D., Canavesi, M., Pfister, P., Fumagalli, R., & Paoletti, R. (1998). HMG-CoA reductase inhibitors reduce MMP-9 secretion by macrophages. *Arterioscler Thromb Vasc Biol* 18, 1671-1678.
- Bettoni, M., & Zimmermann, M. (2002). Autonomic tone variations before the onset of paroxysmal atrial fibrillation. *Circulation* 105, 2753-2759.
- Bialecki, R. A., & Tulenko, T. N. (1989). Excess membrane cholesterol alters calcium channels in arterial smooth muscle. *Am J Physiol* 257, C306-C314.
- Bolotina, V., Omelyanenko, V., Heyes, B., Ryan, U., & Bregestovski, P. (1989). Variations of membrane cholesterol alter the kinetics of Ca<sup>2+</sup>-dependent K<sup>+</sup> channels and membrane fluidity in vascular smooth muscle cells. *Pflugers Arch* 415, 262-268.
- Bonetti, P. O., Wilson, S. H., Rodríguez-Porcel, M., Holmes, D. R. Jr., Lerman, L. O., & Lerman, A. (2002). Simvastatin preserves myocardial perfusion and coronary microvascular permeability in experimental hypercholesterolemia independent of lipid lowering. *J Am Coll Cardiol* 40, 546-554.
- Boos, C. J., Anderson, R. A., & Lip, G. Y. H. (2006). Is atrial fibrillation an inflammatory disorder? *Eur Heart J* 27, 136-149.
- Brown, M., & Goldstein, J. (1998). Sterol regulatory element binding proteins (SREBPs): controllers of lipid synthesis and cellular uptake. *Nutr Rev* 56, S1-S6.
- Bruins, P., te Velthuis, H., Yazdanbakhsh, A. P., Jansen, P. G., van Hardevelt, F. W., de Beaumont, E. M., et al. (1997). Activation of the complement system during and after cardiopulmonary bypass surgery: postsurgery activation involves C-reactive protein and is associated with postoperative arrhythmia. *Circulation* 96, 3542-3548.
- Bunch, T.J., Kottke, T.E., Lopez-Jimenez, F., Mahapatra, S., Elesber, A.A., & White, R.D. (2006). A comparative analysis of short- and long-term outcomes after ventricular fibrillation out-of-hospital cardiac arrest in patients with ischemic and nonischemic heart disease. *Am J Cardiol* 98, 857-860.
- Bustos, C., Hernández-Presa, M., Ortego, M., Tuñón, J., Ortega, L., Pérez F, et al. (1998). HMG-CoA reductase inhibition by atorvastatin reduces neointimal inflammation in a rabbit model of atherosclerosis. *J Am Coll Cardiol* 32, 2057-2064.
- Cai, H., Li, Z., Goette, A., Mera, F., Honeycutt, C., Feterik, K., Wilcox, J. N. et al. (2002). Downregulation of endocardial nitric oxide synthase expression and nitric oxide production in atrial fibrillation: potential mechanisms for atrial thrombosis and stroke. *Circulation* 106, 2854-2858.

- Cao, J. M., Chen, L. S., KenKnight, B. H., Ohara, T., Lee, M. H., Tsai, J., et al. (2000a). Nerve sprouting and sudden cardiac death. *Circ Res* 86, 816-821.
- Cao, J. M., Fishbein, M. C., Han, J. B., Lai, W. W., Lai, A. C., Wu, T. J., et al. (2000b). Relationship between regional cardiac hyperinnervation and ventricular arrhythmia. *Circulation* 101, 1960-1969.
- Caouette, D., Dongmo, C., Berube, J., Fournier, D., & Daleau P. (2003). Hydrogen peroxide modulates the Kv1.5 channel expressed in a mammalian cell line. *Naunyn Schmiedeberg's Arch Pharmacol* 368, 479-486.
- Carnes, C. A., Chung, M. K., Nakayama, T., Nakayama, H., Baliga, R. S., Piao, S., et al. (2001). Ascorbate attenuates atrial pacing-induced peroxynitrite formation and electrical remodeling and decreases the incidence of postoperative atrial fibrillation. *Circ Res* 89, E32-E38.
- Chang, H. M., Reitstetter, R., Mason, R. P., & Gruener, R. (1995). Attenuation of channel kinetics and conductance by cholesterol: an interpretation using structural stress as a unifying concept. *J Membr Biol* 143, 51-63.
- Chen, J., Nagasawa, Y., Zhu, B. M., Ohmori, M., Harada, K., Fujimura, A., et al. (2003). Pravastatin prevents arrhythmias induced by coronary artery ischemia/reperfusion in anesthetized normocholesterolemic rats. *J Pharmacol Sci* 93, 87-94.
- Cheung, B. M., Lauder, I. J., Lau, C-P., & Kumana, C. R. (2004). Meta-analysis of large randomized controlled trials to evaluate the impact of statins on cardiovascular outcomes. *Br J Clin Pharmacol* 57, 640-651.
- Cheruku, K. K., Ghani, A., Ahmad, F., Pappas, P., Silverman, P. R., Zelinger, A., et al. (2004). Efficacy of nonsteroidal anti-inflammatory medications for prevention of atrial fibrillation following coronary artery bypass graft surgery. *Prev Cardiol* 7, 13-18.
- Chiu, J. H., Abdelhadi, R. H., Chung, M. K., Gurm, H. S., Marrouche, N. F., Saliba, W. I., et al. (2005). Effect of statin therapy on risk of ventricular arrhythmia among patients with coronary artery disease and an implantable cardioverter-defibrillator. *Am J Cardiol* 95, 490-491.
- Cholesterol Treatment Trialists' Collaborators. (2005). Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins. *Lancet* 366, 1267-1278.
- Christensen, J. H., Toft, E., Christensen, M. S., & Schmidt, E. B. (1999). Heart rate variability and plasma lipids in men with and without ischaemic heart disease. *Atherosclerosis* 145, 181-186.

- Chung, M. K., Martin, D. O., Sprecher, D., Wazni, O., Kanderian, A., Carnes, C. A., et al. (2001). C-reactive protein elevation in patients with atrial arrhythmias: inflammatory mechanisms and persistence of atrial fibrillation. *Circulation* 104, 2886-2891.
- Cinca, J., Blanch, P., Carreño, A., Mont, L., García-Burillo, A., & Soler-Soler, J. (1997). Acute ischemic ventricular arrhythmias in pigs with healed myocardial infarction: comparative effects of ischemia at a distance and ischemia at the border zone. *Circulation* 96, 653-658.
- Colivicchi, F., Guido, V., Tubaro, M., Ammirati, F., Montefoschi, N., Varvery, A., et al. (2002). Effects of atorvastatin 80 mg daily early after onset of unstable angina pectoris or non-Q-wave myocardial infarction. *Am J Cardiol* 90, 872-874.
- Colivicchi, F., Guido, V., Ficili, S., Mocini, D., Ammirati, F., & Santini, M. (2004). Effects of statin therapy on the recurrence of persistent atrial fibrillation in patients with essential hypertension: a propensity score-adjusted analysis. *Eur Heart J* 25(Suppl), S353.
- Conti, C. R. (2001). Updated pathophysiologic concepts in unstable coronary artery disease. *Am Heart J* 141(2 Suppl), S12-S14.
- Crisby, M., Nordin-Fredriksson, G., Shah, P. K., Yano, J., Zhu, J., & Nilsson, J. (2001). Pravastatin treatment increases collagen content and decreases lipid content, inflammation, metalloproteinases, and cell death in human carotid plaques: implications for plaque stabilization. *Circulation* 103, 926-933.
- Davignon, J. (2004). Beneficial cardiovascular pleiotropic effects of statins. *Circulation* 109(23 Suppl 1), III39-III43.
- De Sutter, J., De Bacquer, D., Jordaens, L., et al. (2006). Intensive lipid-lowering therapy and ventricular arrhythmias in patients with coronary artery disease and internal cardioverter defibrillators. Late Breaking Clinical Trials. Presented in the Heart Rhythm Society 2006 Annual Scientific Sessions; May 17-20, 2006; Boston, Massachusetts.
- De Sutter, J., Tavernier, R., De Buyzere, M., Jordaens, L., & De Backer, G. (2000). Lipid lowering drugs and recurrences of life-threatening ventricular arrhythmias in high-risk patients. *J Am Coll Cardiol* 36, 766-772.
- Dechend, R., Fiebeler, A., Park, J. K., Muller, D. N., Theuer, J., Mervaala, E., et al. (2001). Amelioration of angiotensin II-induced cardiac injury by a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor. *Circulation* 104, 576-581.

- Dernellis, J., & Panaretou, M. (2004). Relationship between C-reactive protein concentrations during glucocorticoid therapy and recurrent atrial fibrillation. *Eur Heart J* 25, 1100-1107.
- Dernellis, J., & Panaretou, M. (2005). Effect of C-reactive protein reduction on paroxysmal atrial fibrillation. *Am Heart J* 150, 1064.
- Dotani, M. I., Elnicki, D. M., Jain, A. C., & Gibson, C. M. (2000). Effect of preoperative statin therapy and cardiac outcomes after coronary artery bypass grafting. *Am J Cardiol* 86, 1128-1130.
- Dupuis, J., Tardif, J. C., Cernacek, P., & Theroux, P. (1999). Cholesterol reduction rapidly improves endothelial function after acute coronary syndromes. The RECIFE (reduction of cholesterol in ischemia and function of the endothelium) trial. *Circulation* 99, 3227-3233.
- Elvan, A., Pride, H. P., Eble, J. N., & Zipes, D.P. (1995). Radiofrequency catheter ablation of the atria reduces inducibility and duration of atrial fibrillation in dogs. *Circulation* 91, 2235-2244.
- Feron, O., Dessy, C., Desager, J. P., & Balligand, J. L. (2001). Hydroxy-Methylglutaryl-Coenzyme A Reductase Inhibition Promotes Endothelial Nitric Oxide Synthase activation Through a Decrease in Caveolin Abundance. *Circulation* 103, 113-118.
- Folkerling, R.J., Van Kraaij, D.J., Tieleman, R.G., Nieman, F.H., Pinto, Y.M., & Crijns, H.J. (2006). Statins associated with reduced mortality in patients admitted for congestive heart failure. *J Card Fail* 12, 134-138.
- Fonarow, G. C., Wright, R. S., Spencer, F. A., Frederick, P. D., Dong, W., Every, N., et al., for the National Registry of Myocardial Infarction 4 Investigators. (2005). Effect of statin use within the first 24 hours of admission for acute myocardial infarction on early morbidity and mortality. *Am J Cardiol* 96, 611-616.
- Fromer, M., Genton, C., Schlaepfer, J., Goy, J. J., & Kappenberger, L. (1990). Is there an isolated arrhythmogenic right atrial myocarditis? *Eur Heart J* 11, 566-571.
- Frustaci, A., Chimenti, C., Bellocci, F., Morgante, E., Russo, M. A., & Maseri, A. (1997). Histological substrate of atrial biopsies in patients with lone atrial fibrillation. *Circulation* 96, 1180-1184.
- Fukuda, K., Davies, S.S., Nakajima, T., Ong, B.H., Kupersmidt, S., Fessel, J., et al. (2005). Oxidative mediated lipid peroxidation recapitulates proarrhythmic effects on cardiac sodium channels. *Circ Res.* 97,1262-1269.

- Fukuta, H., Sane, D.C., Brucks, S., & Little, W.C. (2005). Statin therapy may be associated with lower mortality in patients with diastolic heart failure: a preliminary report. *Circulation* 112, 357-363.
- Furukawa, T., Moroe, K., Mayrovitz, H. N., Sampsel, R., Furukawa, N., & Myerburg, R. J. (1991). Arrhythmogenic effects of graded coronary blood flow reductions superimposed on prior myocardial infarction in dogs. *Circulation* 84, 368-377.
- Fuster, V., Rydén, L. E., Cannom, D. S., Crijns, H. J., Curtis, A. B., Ellenbogen, K. A., et al. (2006). ACC/AHA/ESC 2006 Guidelines for the Management of Patients with Atrial Fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. *Circulation* 114, e257-354.
- Galis, Z. S., Muszynski, M., Sukhova, G. K., Simon-Morrissey, E., Unemori, E. N., Lark, M. W., et al. (1994). Cytokine-stimulated human vascular smooth muscle cells synthesize a complement of enzymes required for extracellular matrix digestion. *Circ Res* 75, 181-189.
- Gaudino, M., Andreotti, F., Zamparelli, R., Di Castelnuovo, A., Nasso, G., Burzotta, F., et al. (2003). The -174G/C interleukin-6 polymorphism influences postoperative interleukin-6 levels and postoperative atrial fibrillation. Is atrial fibrillation an inflammatory complication?. *Circulation* 108(Suppl 1), II195-II199.
- Gleason, M. M., Medow, M. S., & Tulenko, T. N. (1991). Excess membrane cholesterol alters calcium movements, cytosolic calcium levels, and membrane fluidity in arterial smooth muscle cells. *Circ Res* 69, 216-227.
- Goldberger, J. J., Subacius, H., Schaechter, A., Howard, A., Berger, R., Shalaby, A., et al. for the DEFINITE Investigators. (2006). Effects of statin therapy on arrhythmic events and survival in patients with nonischemic dilated cardiomyopathy. *J Am Coll Cardiol* 48, 1228-1233.
- Gómez, R., Núñez, L., Vaquero, M., Caballero, R., Delpón, E., & Tamargo, J. (2005). Effects of atorvastatin on hKv1.5 and Kv4.3 potassium currents. *Eur Heart J* 26 (Suppl), 452.

- Gottlieb, S. S., McCarter, R. J., & Vogel, R. A. (1998). Effect of beta-blockade on mortality among high-risk and low-risk patients after myocardial infarction. *N Engl J Med* 339, 489-497.
- Haigney, M. C., Zareba, W., Gentlesk, P. J., Goldstein, R. E., Illovsky, M., McNitt, S., et al., Multicenter Automatic Defibrillator Implantation Trial II investigators. (2004). QT interval variability and spontaneous ventricular tachycardia or fibrillation in the Multicenter Automatic Defibrillator Implantation Trial (MADIT) II patients. *J Am Coll Cardiol* 44, 1481-1487.
- Hanna, I.R., Heeke, B., Bush, H., Brosius, L., King-Hageman, D., Dudley, S.C. Jr., et al. (2006). Lipid-lowering drug use is associated with reduced prevalence of atrial fibrillation in patients with left ventricular systolic dysfunction. *Heart Rhythm* 3, 881-886.
- Hasin, Y., Shimoni, Y., Stein, O., & Stein, Y. (1980). Effect of cholesterol depletion on the electrical activity of rat heart myocytes in culture. *J Mol Cell Cardiol* 12, 675-683.
- Hayashidani, S., Tsutsui, H., Shiomi, T., Suematsu, N., Kinugawa, S., Ide, T., et al. (2002). Fluvastatin, a 3-hydroxy-3-methylglutaryl coenzyme a reductase inhibitor, attenuates left ventricular remodeling and failure after experimental myocardial infarction. *Circulation* 105, 868-873.
- Holdgate, G. A., Ward, W. H., & McTaggart, F. (2003). Molecular mechanism for inhibition of 3-hydroxy-3-methylglutaryl CoA (HMG-CoA) reductase by rosuvastatin. *Biochem Soc Trans* 31, 528-531.
- Horwich, T. B., MacLellan, W. R., & Fonarow, G. C. (2004). Statin therapy is associated with improved survival in ischemic and non-ischemic heart failure. *J Am Coll Cardiol* 43, 642-648.
- Huikuri, H. V., Jokinen, V., Syvanne, M., Nieminen, M. S., Airaksinen, K. E., Ikaheimo, M. J., et al. (1999). Heart rate variability and progression of coronary atherosclerosis. *Arterioscler Thromb Vasc Biol* 19, 1979-1985.
- Indolfi, C., Di Lorenzo, E., Perrino, C., Stingone, A. M., Curcio, A., Torella, D., et al. (2002). Hydroxymethylglutaryl coenzyme A reductase inhibitor simvastatin prevents cardiac hypertrophy induced by pressure overload and inhibits p21ras activation. *Circulation* 106, 2118-2124.
- Jain, M. K., & Ridker, P. M. (2005). Anti-inflammatory effects of statins: clinical evidence and basic mechanisms. *Nat Rev Drug Discov* 4, 977-987.

- Jialal, I., Stein, D., Balis, D., Grundy, S. M., Adams-Huet, B., & Devaraj, S. (2001). Effect of hydroxymethyl glutaryl coenzyme a reductase inhibitor therapy on high sensitive C-reactive protein levels. *Circulation* 103, 1933-1935.
- Kannel, W.B. (1983). Prevalence and natural history of electrocardiographic left ventricular hypertrophy. *Am J Med* 75(3A): 4-11.
- Karagueuzian, H. S., Pennec, J. P., Deroubaix, E., de Leiris, J., & Coraboeuf, E. (1982). Effects of excess free fatty acids on the electrophysiological properties of ventricular specialized conducting tissue: a comparative study between the sheep and the dog. *J Cardiovasc Pharmacol* 4, 462-468.
- Katchman, A., Koerner, J., Tosak, A. T., Woosley, R., & Ebert, S. (2006). Comparative evaluation of HERG currents and QT intervals following challenge with suspected torsadogenic and nontorsadogenic drugs. *J Pharmacol Exp Ther* 316, 1098-1106.
- Kawabata, H., Ryomoto, T., & Ishikawa, K. (2001). Role of cardiac ATP-sensitive K<sup>+</sup> channels induced by HMG CoA reductase inhibitor in ischemic rabbit hearts. *Hypertens Res* 24, 573-577.
- Kayikcioglu, M., Can, L., Evrengul, H., Payzin, S., & Kultursay, H. (2003). The effect of statin therapy on ventricular late potentials in acute myocardial infarction. *Int J Cardiol* 90, 63-72.
- Kim, Y.M., Guzik, T.J., Zhang, Y.H., Zhang, M.H., Kattach, H., Ratnatunga, C., et al. (2005). A myocardial Nox2 containing NAD(P)H oxidase contributes to oxidative stress in human atrial fibrillation. *Circ Res*. 97: 629-636.
- Kjekshus, J., Pedersen, T. R., Olsson, A. G., Faergeman, O., & Pyorala, K. (1997). The effects of simvastatin on the incidence of heart failure in patients with coronary heart disease. *J Card Fail* 3, 249-254.
- Kolvekar, S., D'Souza, A., Akhtar, P., Reek, C., Garratt, C., & Spyt, T. (1997). Role of atrial ischemia in development of atrial fibrillation following coronary artery bypass surgery. *Eur J Cardiothorac Surg* 11, 70-75.
- Krum, H., & McMurray, J. J. (2002). Statins and chronic heart failure: do we need a large-scale outcome trial? *J Am Coll Cardiol* 39, 1567-1573.
- Kumagai, K., Nakashima, H., & Saku, K. (2004). The HMG-CoA reductase inhibitor atorvastatin prevents atrial fibrillation by inhibiting inflammation in a canine sterile pericarditis model. *Cardiovasc Res* 62, 105-111.

- Kutryk, M. J., & Pierce, G. N. (1988). Stimulation of sodium-calcium exchange by cholesterol incorporation into isolated cardiac sarcolemmal vesicles. *J Biol Chem* 263, 13167-13172.
- Lamers, J. M., Hartog, J. M., Verdouw, P. D., & Hulsmann, W. C. (1987). Dietary fatty acids and myocardial function. *Basic Res Cardiol* 82(Suppl 1), 209-221.
- Landmesser, U., & Drexler, H. (2005). The clinical significance of endothelial dysfunction. *Curr Opin Cardiol* 20, 547-551.
- Laufs, U., Custodis, F., & Böhm, M. (2006). HMG-CoA reductase inhibitors in chronic heart failure. *Drugs* 66,146-154.
- Laufs, U., Wassmann, S., Hilgers, S., Ribaldo, N., Bohm, M., & Nickenig, G. (2001). Rapid effects on vascular function after initiation and withdrawal of atorvastatin in healthy, noncholesterolemic men. *Am J Cardiol* 88, 1306-1307.
- Lee, T.M., Chou, T.F., & Tsai, C.H. (2002). Association of pravastatin and left ventricular mass in hypercholesterolemic patients: role of 8-iso-prostaglandin f2alpha formation. *J Cardiovasc Pharmacol* 40, 868-874.
- Lee, T. M., Chou, T. F., & Tsai, C. H. (2003). Effects of pravastatin on cardiomyocyte hypertrophy and ventricular vulnerability in normolipidemic rats after myocardial infarction. *J Moll Cell Cardiol* 35, 1449-1459.
- Lee, T. M., Lin, M. S., Chou, T. F., Tsai, C. H., & Chang, N. C. (2004). Effect of pravastatin on left ventricular mass by activation of myocardial K ATP channels in hypercholesterolemic rabbits. *Atherosclerosis* 176, 273-278.
- Lee, T. M., Lin, M. S., Chou, T. F., & Chang, N. C. (2005a). Effect of simvastatin on left ventricular mass in hypercholesterolemic rabbits. *Am J Physiol Heart Circ Physiol* 288, H1352-H1358.
- Lee, T. M., Lin, M. S., Chou, T. F., Tsai, C. H., & Chang, N. C. (2005b). Effect of pravastatin on development of left ventricular hypertrophy in spontaneously hypertensive rats. *Am J Physiol Heart Circ Physiol* 289, H220-H227.
- Lefler, A. M., Campbell, B., Shin, Y. J., Scalia, R., Hayward, R., & Lefler, D. J. (1999). Simvastatin preserves the ischemic-reperfused myocardium in normocholesterolemic rat hearts. *Circulation* 100, 178-184.
- Lefler, D. J., Scalia, R., Jones, S. P., Sharp, B. R., Hoffmeyer, M. R., Farvid, A. R., et al. (2001). HMG-CoA reductase inhibition protects the diabetic myocardium from ischemia-reperfusion injury. *FASEB J* 15, 1454-1456.

- Levy, D., Garrison, R.J., Savage, D.D., Kannel, W.B., & Castelli, W.P. (1990). Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N Engl J Med* 322, 1561–1566.
- Li, L., Matsuoka, I., Suzuki, Y., Watanabe, Y., Ishibashi, T., Yokoyama, K., et al. (2002). Inhibitory effect of fluvastatin on lysophosphatidylcholine-induced nonselective cation current in Guinea pig ventricular myocytes. *Mol Pharmacol* 62, 602-607.
- Liao, J. K., & Laufs, U. (2005). Pleiotropic effects of statins. *Annu Rev Pharmacol Toxicol* 45, 89-118.
- Libby, P. (2001). Current concepts of the pathogenesis of the acute coronary syndromes. *Circulation* 104, 365-372.
- Lijnen, P., Celis, H., Fagard, R., Staessen, J., & Amery, A. (1994). Influence of cholesterol lowering on plasma membrane lipids and cationic transport systems. *J Hypertens* 12, 59-64.
- Lijnen, P., & Petrov, V. (1995). Cholesterol modulation of transmembrane cation transport systems in human erythrocytes. *Biochem Mol Med* 56, 52-62.
- Liu, Y. B., Wu, C. C., Lu, L. S., Su, M. J., Lin, C. W., Lin, S. F., et al. (2003). Sympathetic nerve sprouting, electrical remodeling, and increased vulnerability to ventricular fibrillation in hypercholesterolemic rabbits. *Circ Res* 92, 1145-1152.
- Locher, R., Neyses, L., Stimpfle, M., Kuffer, B., & Vetter, W. (1984). The cholesterol content of the human erythrocyte influences calcium influx through the channel. *Biochem Biophys Res Commun* 124, 822-828.
- López-Sendón, J., Swedberg, K., McMurray, J., Tamargo, J., Maggioni, A. P., Dargie, H., et al., Task Force On Beta-Blockers of the European Society of Cardiology. (2004). Expert consensus document on beta-adrenergic receptor blockers. *Eur Heart J* 25, 1341-1362.
- Lorenz, H., Junger, C., Seidl, K., Gitt, A., Schneider, S., Schiele, R., et al. (2005). Do statins influence the prognostic impact of non-sustained ventricular tachycardia after ST-elevation myocardial infarction? *Eur Heart J* 26, 1078-1085.
- Luan, Z., Chase, A. J., & Newby, A. C. (2003). Statins inhibit secretion of metalloproteinases-1,-2,-3, and -9 from vascular smooth muscle cells and macrophages. *Arterioscler Thromb Vasc Biol* 23, 769-775.
- Maack, C., Kartes, T., Kilter, H., Schafers, H.J., Nickenig, G., Bohm, M., et al. (2003). Oxygen free radical release in human failing myocardium is associated with increased activity of rac1-GTPase and represents a target for statin treatment. *Circulation* 108, 1567-1574.

- Maixent, J.M., Paganelli, F., Scaglione, J., & Levy, S. (1998). Antibodies against myosin in sera of patients with idiopathic paroxysmal atrial fibrillation. *J Cardiovasc Electrophysiol* 9, 612-617.
- Marín, F., Pascual, D. A., Roldán, V., Arribas, J. M., Ahumada, M., Tornel, P. L., et al. (2006). Statins and postoperative risk of atrial fibrillation following coronary artery bypass grafting. *Am J Cardiol* 97, 55-60.
- Mark, L., & Katona, A. (2000). Effect of fluvastatin on QT dispersion: a new pleiotropic effect? *Am J Cardiol* 85, 919-920.
- Martens, J. R., O'Connell, K., & Tamkun, M. (2004). Targeting of ion channels to membrane microdomains: localization of Kv channels to lipid rafts. *Trends Pharmacol Sci* 25, 16-21.
- Mason, R. P., Walter, M. F., & Jacob, R. F. (2004). Effects of HMG-CoA reductase inhibitors on endothelial function: role of microdomains and oxidative stress. *Circulation* 109(21 Suppl 1): II34-II41.
- Mehta, D., Curwin, J., Gomes, J. A., & Fuster, V. (1997). Sudden death in coronary artery disease: acute ischemia versus myocardial substrate. *Circulation* 96, 3215-3223.
- Mendes, L. A., Connelly, G. P., McKenney, P. A., Podrid, P. J., Cupples, L. A., Shemin, R. J., et al. (1995). Right coronary artery stenosis: an independent predictor of atrial fibrillation after coronary artery bypass surgery. *J Am Coll Cardiol* 25, 198-202.
- Merckx, L., Tieleman, R. G., Folkeringa, R. J., Pinto, Y. M., Niemann F, H., Cheriex, E. C., et al. (2004). Use of statins is associated with reduced incidence of atrial fibrillation in patients with left ventricular hypertrophy and left atrial dilatation. *Heart Rhythm* 1 (1 Suppl 1), S105.
- Mihm, M. J., Yu, F., Carnes, C. A., Reiser, P. J., McCarthy, P. M., Van Wagoner, D. R., et al. (2001). Impaired myofibrillar energetics and oxidative injury during human atrial fibrillation. *Circulation* 104, 174-180.
- Mitchell, L. B., Powell, J. L., Gillis, A. M., Kehl, V., & Hallstrom, A. P.; AVID Investigators. (2003). Are lipid-lowering drugs also antiarrhythmic drugs? An analysis of the Antiarrhythmics versus Implantable Defibrillators (AVID) trial. *J Am Coll Cardiol* 42, 81-87.
- Mukai, Y., Shimokawa, H., Matoba, T., Hiroki, J., Kunihiro, I., Fujiki, T., et al. (2003). Acute vasodilator effects of HMG-CoA reductase inhibitors: involvement of PI3-kinase/Akt pathway and Kv channels. *J Cardiovasc Pharmacol* 42, 118-124.
- Nattel, S. (2002). New ideas about atrial fibrillation 50 years on. *Nature* 415, 219-226.

- Node, K., Fujita, M., Kitakaze, M., Hori, M., & Liao, J. K. (2003). Short-term statin therapy improves cardiac function and symptoms in patients with idiopathic dilated cardiomyopathy. *Circulation* 108, 839-843.
- O'Connell, S., & Tamkun, M. M. (2005). Targeting of voltage-gated potassium channel isoforms to distinct cell surface microdomains. *J Cell Sci* 118, 2155-2166.
- Oi, S., Haneda, T., Osaki, J., Kashiwagi, Y., Nakamura, Y., Kawabe, J., et al. (1999). Lovastatin prevents angiotensin II-induced cardiac hypertrophy in cultured neonatal rat heart cells. *Eur J Pharmacol* 376,139-148.
- Ortega, A., & Mas-Oliva, J. (1984). Cholesterol effect on enzyme activity of sarcolemmal (Ca<sup>2+</sup>+Mg<sup>2+</sup>)-ATPase from cardiac muscle. *Biochim Biophys Acta* 773, 231-236.
- Ozaydin, M., Varol, E., Aslan, S. M., Kucuktepe, Z., Dogan, A., Ozturk, M., et al. (2006). Effect of atorvastatin on the recurrence rates of atrial fibrillation after electrical cardioversion. *Am J Cardiol* 97, 1490-1493.
- Pandit, S.V., Berenfeld, O., Anumonwo, J. M., Zaritski, R. M., Kneller, J., Nattel, S., et al. (2005). Ionic determinants of functional reentry in a 2-D model of human atrial cells during simulated chronic atrial fibrillation. *Biophys J* 88, 3806-3821.
- Patti, G., Chello, M., Candura, D., Pasceri, V., D'Ambrosio, A., Covino, E., et al. (2006). Randomized trial of atorvastatin for reduction of postoperative atrial fibrillation in patients undergoing cardiac surgery: results of the ARMYDA-3 (Atorvastatin for Reduction of MYocardial Dysrhythmia After cardiac surgery) study. *Circulation* 114, 1455-1461.
- Pehlivanidis, A. N., Athyros, V. G., Demetriadis, D. S., Papageorgiou, A. A., Bouloukos, V. J., & Kontopoulos, A. G. (2001). Heart rate variability after long-term treatment with atorvastatin in hypercholesterolaemic patients with or without coronary artery disease. *Atherosclerosis* 157, 463-469.
- Pelat, M., Dessy, C., Massion, P., Desager, J. P., Feron, O., & Balligand, J. L. (2003). Rosuvastatin decreases caveolin-1 and improves nitric oxide-dependent heart rate and blood pressure variability in apolipoprotein E<sup>-/-</sup> mice *in vivo*. *Circulation* 107, 2480-2486.
- Pfeffer, M. A., & Braunwald, E. (1990). Ventricular remodeling after myocardial infarction. Experimental observations and clinical implications. *Circulation* 81, 1161-1172.
- Pliquett, R. U., Cornish, K. G., Peuler, J. D., & Zucker, I. H. (2003). Simvastatin normalizes autonomic neural control in experimental heart failure. *Circulation* 107, 2493-2498.

- Pound, E. M., Kang, J. X., & Leaf, A. (2001). Partitioning of polyunsaturated fatty acids, which prevent cardiac arrhythmias, into phospholipid cell membranes. *J Lipid Res* 42, 346-351.
- Pratt, C. M., Waldo, A. L., & Camm, A. J. (1998). Can antiarrhythmic drugs survive survival trials? *Am J Cardiol* 81, 24D-34D.
- Quinn, P. J. (1981). The fluidity of cell membranes and its regulation. *Prog Biophys Mol Biol* 38, 1-104.
- Quintana, M., Storck, N., Lindblad, L. E., Lindvall, K., & Ericson, M. (1997). Heart rate variability as a means of assessing prognosis after acute myocardial infarction. A 3-year follow-up study. *Eur Heart J* 18, 789-797.
- Raina, A., Pickering, T., & Shimbo, D. (2006). Statin use in heart failure: A cause for concern? *Am Heart J* 152, 39-49.
- Ray, K. K., & Cannon, C. P. (2005). The potential relevance of the multiple lipid-independent (pleiotropic) effects of statins in the management of acute coronary syndromes. *J Am Coll Cardiol* 46, 1425-1433.
- Ren, Y. J., Xu, X. H., Zhong, C. B., Feng, N., & Wang, X. L. (2001). Hypercholesterolemia alters vascular functions and gene expression of potassium channels in rat aortic smooth muscle cells. *Acta Pharmacol Sin* 22, 274-278.
- Riahi, S., Christensen, J. H., Toft, E., Skou, H. A., & Schmidt, E. B. (2002). HMG-CoA reductase inhibitors improve heart rate variability in patients with a previous myocardial infarction. *Pharmacol Res* 45, 479-483.
- Ridker, P. M., Rifai, N., Pfeffer, M. A., Sacks, F., & Braunwald, E. (1999). Long-term effects of pravastatin on plasma concentration of C-reactive protein. The Cholesterol and Recurrent Events (CARE) Investigators. *Circulation* 100, 230-235.
- Ridker, P. M. (2001a). High-sensitivity C-reactive protein: potential adjunct for global risk assessment in the primary prevention of cardiovascular disease. *Circulation*. 103,1813-1818.
- Ridker, P. M., Rifai, N., Clearfield, M., Downs, J. R., Weis, S. E., Miles, J. S., et al., for the Air Force/Texas Coronary Atherosclerosis Prevention Study Investigators. (2001b). Measurement of C-reactive protein for the targeting of statin therapy in the primary prevention of acute coronary events. *N Engl J Med* 344, 1959-1965.
- Rikitake, Y., Kawashima, S., Takeshita, S., Yamashita, T., Azumi, H., Yasuhara, M., et al. (2001). Anti-oxidative properties of fluvastatin, an HMG-CoA reductase inhibitor,

- contribute to prevention of atherosclerosis in cholesterol-fed rabbits. *Atherosclerosis* 154, 87-96.
- Romanenko, V. G., Rothblat, G. H., & Levitan, I. (2002). Modulation of endothelial inward-rectifier K<sup>+</sup> current by optical isomers of cholesterol. *Biophys J* 83, 3211-3222.
- Rubart, M., & Zipes, D. P. (2005). Mechanisms of sudden cardiac death. *J Clin Invest* 115, 2305-2315.
- Sacks, F. M., Pfeffer, M. A., Moye, L. A., Brown, L. E., Hamm, P., Cole, T. G., et al. (1996). The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med* 335, 1001-1009.
- Sasaki, N., Sato, T., Ohler, A., O'Rourke, B., & Marban, E. (2000). Activation of mitochondrial ATP-dependent potassium channels by nitric oxide. *Circulation* 101, 439-445.
- Scalia, R., Gooszen, M. E., Jones, S. P., Hoffmeyer, M., Rimmer, D. M. 3rd., Trocha, S. D., et al. (2001). Simvastatin exerts both anti-inflammatory and cardioprotective effects in apolipoprotein E-deficient mice. *Circulation* 103, 2598-2603.
- Scandinavian Survival Study Group. (1994). Randomized trial of cholesterol lowering in 4,444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 344, 1383-1389.
- Schauerte, P., Scherlag, B. J., Pitha, J., Scherlag, M. A., Reynolds, D., Lazzara, R, et al. (2000). Catheter ablation of cardiac autonomic nerves for prevention of vagal atrial fibrillation. *Circulation* 102, 2774-2780.
- Schonbeck, U., & Libby, P. (2004). Inflammation, immunity, and HMG-CoA reductase inhibitors: statins as antiinflammatory agents? *Circulation* 109(21 Suppl 1), II18-II26.
- Schwartz, G., Olsson, A., Ezekowitz, M., Ganz, P., Oliver, M. F., Waters, D., et al. (2001). Effects of atorvastatin on early recurrent ischemic events in acute coronary syndromes. The MIRACL Study: a randomized trial. *JAMA* 285, 1711-1718.
- Seino, S. & Miki, T. (2003). Physiological and pathophysiological roles of ATP-sensitive K<sup>+</sup> channels. *Prog Biophys Mol Biol* 81, 133-176.
- Shioiri, H., Komaru, T., Sato, K., Takahashi, K., Takeda, S., Kanatsuka, H., et al. (2003). Impact of hypercholesterolemia on acidosis-induced coronary microvascular dilation. *Basic Res Cardiol* 98, 76-83.
- Shiroshita-Takeshita, A., Schram, G., Lavoie, J., & Nattel, S. (2004). Effect of simvastatin and antioxidant vitamins on atrial fibrillation promotion by atrial-tachycardia remodeling in dogs. *Circulation* 110, 2313-2319.

- Shusterman, V., Aysin, B., Gottipaty, V., Weiss, R., Brode, S., Schwartzman, D., et al. (1998). Autonomic nervous system activity and the spontaneous initiation of ventricular tachycardia. *J Am Coll Cardiol* 32, 1891-1899.
- Sinno, H., Derakhchan, K., Libersan, D., Merhi, Y., Leung, T. K., & Nattel, S. (2003). Atrial ischemia promotes atrial fibrillation in dogs. *Circulation* 107, 1930-1936.
- Siu, C. W., Lau, C. P., & Tse, H. F. (2003). Prevention of atrial fibrillation recurrence by statin therapy in patients with lone atrial fibrillation after successful cardioversion. *Am J Cardiol* 92, 1343-1345.
- Sobel, B. E., Corr, P. B., Robinson, A. K., Goldstein, R. A., Witkowski, F. X., & Lein, M. S. (1978). Accumulation of lysophosphoglycerides with arrhythmogenic properties in ischemic myocardium. *J Clin Invest* 62, 546-553.
- Sola, S., Mir, M. Q., Rajagopalan, S., Helmy, T., Tandon, N., & Khan, B. V. (2005). Statin therapy is associated with improved cardiovascular outcomes and levels of inflammatory markers in patients with heart failure. *J Card Fail* 11, 607-612.
- Sola, S., Mir, M. Q., Lerakis, S., Tandon, N., & Khan, B. V. (2006). Atorvastatin improves left ventricular systolic function and serum markers of inflammation in nonischemic heart failure. *J Am Coll Cardiol* 47, 332-337.
- Sorescu, D., & Griendling, K. K. (2002). Reactive oxygen species, mitochondria, and NAD(P)H oxidases in the development and progression of heart failure. *Congest Heart Fail* 8, 132-140.
- Sukhova, G. K., Williams, J. K., & Libby, P. (2002). Statins reduce inflammation in atheroma of nonhuman primates independent of effects on serum cholesterol. *Arterioscler Thromb Vasc Biol* 22, 1452-1458.
- Sundstrom, J., Lind, L., Vessby, B., Andren, B., Aro, A., & Lithell, H. (2001). Dyslipidemia and an unfavorable fatty acid profile predict left ventricular hypertrophy 20 years later. *Circulation* 103, 836-841.
- Suzumura, K., Yasuhara, M., Tanaka, K., & Suzuki, T. (1999). Protective effect of fluvastatin sodium (XU-62-320), a 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitor, on oxidative modification of human low-density lipoprotein in vitro. *Biochem Pharmacol* 57, 697-703.
- Szmitko, P.E., Wang, C.H., Weisel, R.D., Jeffries, G.A., Anderson, T.J., & Verma, S. (2003). Biomarkers of vascular disease linking inflammation to endothelial activation: Part I. *Circulation*. 108, 1917-1923.

- Takemoto, M., Node, K., Nakagami, H., Liao, Y., Grimm, M., Takemoto, Y., et al. (2001). Statins as antioxidant therapy for preventing cardiac myocyte hypertrophy. *J Clin Invest* 108, 1429-1437.
- Tamargo, J., Caballero, R., & Delpón, E. (2004a). Pharmacological approach in the treatment of atrial fibrillation. *Curr Med Chem* 11, 13-28.
- Tamargo, J., Caballero, R., Gomez, R., Valenzuela, C., & Delpón, E. (2004b). Pharmacology of cardiac potassium channels. *Cardiovasc Res* 62, 9-33.
- Terata, Y., Saito, T., Fujiwara, Y., Hasegawa, H., Miura, H., Watanabe, H., et al. (2003). Pitavastatin inhibits upregulation of intermediate conductance calcium-activated potassium channels and coronary arteriolar remodeling induced by long-term blockade of nitric oxide synthesis. *Pharmacology* 68, 169-176.
- Tesfamariam, B., Frohlich, B. H., & Gregg, R. E. (1999). Differential effects of pravastatin, simvastatin, and atorvastatin on  $Ca^{2+}$  release and vascular reactivity. *J Cardiovasc Pharmacol* 34, 95-101.
- Tomaselli, G. F., & Zipes, D. P. (2004). What causes sudden death in heart failure? *Circ Res* 95, 754-763.
- Tomita, T., Takei, M., Saikawa, Y., Hanaoka, T., Uchikawa, S., Tsutsui, H., et al. (2003). Role of autonomic tone in the initiation and termination of paroxysmal atrial fibrillation in patients without structural heart disease. *J Cardiovasc Electrophysiol* 14, 559-564.
- The Long-term Intervention with Pravastatin in Ischemic Disease (LIPID) Study Group. (1998). Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. *N Engl J Med* 339, 1349-1357.
- Tveit, A., Grundtvig, M., Gundersen, T., Vanberg, P., Semb, A. G., Holt, E., et al. (2004). Analysis of pravastatin to prevent recurrence of atrial fibrillation after electrical cardioversion. *Am J Cardiol* 93, 780-782.
- Tulenko, T. N., Bialecki, R., Gleason, M., & D'Angelo, G. (1990). Ion channels, membrane lipids and cholesterol: a role for membrane lipid domains in arterial function. *Prog Clin Biol Res* 334, 187-203.
- Ueda, Y., Kitakaze, M., Komamura, K., Minamino, T., Asanuma, H., Sato, H., et al. (1999). Pravastatin restored the infarct size-limiting effect of ischemic preconditioning blunted by hypercholesterolemia in the rabbit model of myocardial infarction. *J Am Coll Cardiol* 34, 2120-2125.

- Van Boven AJ, Jukema JW, Crijns HJ, Lie KI. (1995). Heart rate variability profiles in symptomatic coronary artery disease and preserved left ventricular function: relation to ventricular tachycardia and transient myocardial ischemia. Regression Growth Evaluation Statin Study (REGRESS). *Am Heart J* 130, 1020-1025.
- Van Boven, A. J., Jukema, J. W., Zwindermann, A. H., Crijns, H. J., Lie, K. I., & Bruschke, A. V. (1996). Reduction of transient myocardial ischemia with pravastatin in addition to the conventional treatment in patients with angina pectoris. REGRESS Study Group. *Circulation* 94(7), 1503-1505.
- Vaquero, M., Núñez, L., Gómez, R., Caballero, R., Delpón, E., & Tamargo, J. (2005). Efectos de la atorvastatina sobre las corrientes de potasio responsables de la repolarización auricular humana, hKv1.5 y Kv4.3. *Rev Esp Cardiol* 58(Suppl 1), 130.
- Vaquero, M., Núñez, L., Gómez, R., Caballero, R., Delpón, E., & Tamargo, J. (2006). Effects of simvastatin on cardiac hKv1.5, Kv4.3, and calcium currents. *Meth and Findings in Exp Clin Pharmacol*. In press.
- Ver Donck, L., Verellen, G., Geerts, H., & Borgers, M. (1992). Lysophosphatidylcholine-induced  $Ca^{2+}$ -overload in isolated cardiomyocytes and effect of cytoprotective drugs. *J Mol Cell Cardiol* 24, 977-988.
- Wolk, R. (2000). Arrhythmogenic mechanisms in left ventricular hypertrophy. *Europace* 2, 216-223.
- Vrtovec, B., Okrajsek, R., Golicnik, A., Ferjan, M., Starc, V., & Radovancevic, B. (2005). Atorvastatin therapy increases heart rate variability, decreases QT variability, and shortens QTc interval duration in patients with advanced chronic heart failure. *J Card Fail* 11, 684-690.
- Vyas, A., Guo, H., Moss, A. J., Olshansky, B., McNitt, S. A., Hall, W. J., et al., for the MADIT-II Research Group. (2006). Reduction in ventricular tachyarrhythmias with statins in the Multicenter Automatic Defibrillator Implantation Trial (MADIT)-II. *J Am Coll Cardiol* 47, 769-773.
- Wagner, A. H., Köhler, T., Rückschloss, U., Just, I., & Hecker, M. (2000). Improvement of nitric oxide-dependent vasodilatation by HMG-CoA reductase inhibitors through attenuation of endothelial superoxide anion formation. *Arterioscler Thromb Vasc Biol* 20, 61-69.
- Wassmann, S., Laufs, U., Muller, K., Konkol, C., Ahlbory, K., Baumer, A. T., et al. (2002a). Cellular antioxidant effects of atorvastatin in vitro and in vivo. *Arterioscler Thromb Vasc Biol* 22(2), 300-305.

- Wassmann, S., & Nickenig, G. (2002b). Improvement of endothelial function by HMG-CoA reductase inhibitors. *Drug News Perspect* 15, 85-92.
- Welzig, C. M., Shin, D. G., Park, H. J., Kim, Y.J., Saul, J. P., & Galper, J. B. (2003). Lipid lowering by pravastatin increases parasympathetic modulation of heart rate: Galpha(i2), a possible molecular marker for parasympathetic responsiveness. *Circulation* 108, 2743-2746.
- Wijffels, M., & Crijns, H.J. (2003). Recent advances in drug therapy for atrial fibrillation. *J Cardiovasc Electrophysiol* 14(9 Suppl), S40-S47.
- Williams, J. K., Sukhova, G. K., Herrington, D. M., & Libby, P. (1998). Pravastatin has cholesterol-lowering independent effects on the artery wall of atherosclerotic monkeys. *J Am Coll Cardiol* 31, 684-691.
- Wolf, P. A., Abbott, R. D., & Kannel, W. B. (1987). Atrial fibrillation: a major contributor to stroke in the elderly. The Framingham Study. *Arch Intern Med* 147, 1561-1564.
- Wu, C. C., Su, M. J., Chi, J. F., Chen, W. J., Hsu, H. C., & Lee, Y. T. (1995). The effect of hypercholesterolemia on the sodium inward currents in cardiac myocyte. *J Mol Cell Cardiol* 27, 1263-1269.
- Yamazaki, M., Suzuki, H., Hanano, M., & Sugiyama, Y. (1993). Different relationships between cellular ATP and hepatic uptake among taurocholate, cholate, and organic anions. *Am J Physiol* 264, G693-G701.
- Yeagle, P. L., Young, J., & Rice, D. (1988). Effects of cholesterol on (Na<sup>+</sup>,K<sup>+</sup>)-ATPase ATP hydrolyzing activity in bovine kidney. *Biochemistry* 27, 6449-6452.
- Yeagle, P. L. (1991). Modulation of membrane function by cholesterol. *Biochimie* 73, 1303-1310.
- Yokoyama, K., Ishibashi, T., Ohkawara, H., Kimura, J., Matsuoka, I., Sakamoto, T., et al. (2002). HMG-CoA reductase inhibitors suppress intracellular calcium mobilization and membrane current induced by lysophosphatidylcholine in endothelial cells. *Circulation* 105, 962-967.
- Young-Xu, Y., Jabbour, S., Goldberg, R., Blatt, C. M., Graboys, T., Bilchik, B., et al. (2003). Usefulness of statin drugs in protecting against atrial fibrillation in patients with coronary artery disease. *Am J Cardiol* 92, 1379-1383.
- Zipes, D. P., & Wellens, H. J. (1998). Sudden cardiac death. *Circulation* 98, 2334-2351.

**Table 1. Studies of statins to prevent ventricular tachyarrhythmias**

Study	Design/n	Medication/ subjects	Odds ratio/RRR (95% CI)	Comments
<b>Patients with ICD</b>				
Lipid lowering therapy prevents recurrences of life-threatening VA (De Sutter et al, 2000)	Observational n = 78	Lipid-lowering users vs non users. Patients with CAD and VA treated with an ICD	0.61 (0.34-0.77) (P = 0.004)	Lipid-lowering drugs is associated with a reduction of recurrences of VA in patients with CAD and ICD implants
On VA recurrence (Mitchell et al, 2003)	Prospective, multicenter, R n = 237	Lipid-lowering users vs non users. Patients with CAD and near-fatal VA with an ICD	0.40 (0.15-0.58) (P = 0.04)	Lipid-lowering therapy decreases the risk of VA recurrence
On recurrent VA (Chiu et al, 2005)	Prospective n = 281	Statin users vs non users. Patients with CAD treated with a ICD	0.60 (0.41-0.89) (P = 0.01)	Statin use decreases the risk of VA that would require ICD
On first recurrence VA (De Sutter et al., 2006)	Prospective, R, PC n = 106	Atorvastatin 80 mg/day vs no drug. Patients with life-threatening VA requiring an ICD	21% vs 38% 0.47 (0.22-0.98) (P = 0.04)	Atorvastatin reduces the first recurrence of VA after one year treatment
On patients with nonischemic dilated cardiomyopathy (Goldberger et al, 2006)	Prospective, R n = 458	Statin users vs non users. Patients with nonischemic dilated cardiomyopathy at risk for VA	0.22 (0.09-0.58) (P = 0.001)	Statin therapy reduces the rate of total mortality and arrhythmic sudden death
On patients with an ICD (Vyas et al,	R, open, parallel-groups	Statin users vs non users. Patients with a	0.65 (0.49-0.87) (P < 0.01)	Statins reduce the risk of cardiac death or VT/VF

2006)	n = 654	previous MI (EF = 30%) receiving an ICD		
<b>Patients without ICD</b>				
On post CABG arrhythmias (Dotani et al, 2000)	Retrospective, nonrandomized n = 323	Lipid-lowering users vs non users. Patients undergoing CABG	0.23 (0.08-0.65) (P = 0.006)	Preoperative statin use decreases the incidence of death, unstable angina and cardiac arrhythmias (sustained AF, sustained or nonsustained VT and VF)
On acute MI (Kayikcioglu et al, 2003)	Prospective n = 72	Pravastatin 40 mg/day vs no drug. Patients with acute MI	27% vs 63% (P = 0.006)	Statins reduce the incidence of in-hospital VA in acute MI
In patients with systolic heart failure (Horwich et al, 2004)	n = 551	Statin users vs non users.	0.41 (0.18-0.94) (P = 0.01)	Statin use decreases mortality and the need of urgent cardiac transplantation
In the acute phase of MI (Fonarow et al., 2005)	Prospective, observational n = 174,635	Statin users vs non users. Patients with acute MI	3.7% vs 4.7% 0.23 (0.22-0.25) (P < 0.001)	Statins given within the first 24 hours of hospitalization for acute MI reduce the rate of VT/VF
On NSVT after ST-elevation MI (Lorenz et al., 2005)	Prospective, multicenter, observational n = 346	Statin users vs non users. Patients with ST- elevation MI	0.39 (0.15-0.98) (P < 0.047)	In patients with ST-elevation MI under statin treatment NSVT is not associated with an adverse long-term prognosis
Outcomes after VF out-of-hospital cardiac arrest (Bunch et al., 2006)	n = 208	Statin users s non-users	0.68 (0.17-0.73) (P = 0.001)	Statin use was associated with decreased mortality

CABG: coronary artery bypass grafting. CAD: coronary artery disease. CI: 95% confidence interval. CRP: C-reactive protein. EF: ejection fraction. ICD: implantable cardioverter defibrillator MI: myocardial infarction. NSVT: non-sustained ventricular tachycardia. PC: placebo-controlled. R: randomized. RRR: relative risk reduction. VA/VF/VT: ventricular arrhythmias/fibrillation/tachycardia.

**Table 2. Studies of statins to prevent atrial fibrillation**


<b>Study</b>	<b>Design/n</b>	<b>Medication/ subjects</b>	<b>OR/RRR (95% CI)</b>	<b>Comments</b>
Prevention of AF recurrence after cardioversion (Siu et al, 2003)	Retrospective n = 62	Statin users vs non users. Patients with persistent lone AF (> 3 months)	0.31 (CI 0.10-0.90) (P = 0.032)	Patients on statins had higher cholesterol and were older than nonusers. Statins decrease recurrent AF after cardioversion
Statins in protecting against AF (Young-Xu et al, 2003)	Retrospective n = 449	Statin users vs non users. Patients with CAD in SR at high risk of AF	0.48 (CI 0.28-0.83) (P = 0.01)	Effect independent of changes in LDL-C. Dose-response relation between length of statin use and reduction of AF
Reduction of AF after cardiac surgery (Auer et al., 2004).	n = 253	Statin user vs non users. Patients undergone major cardiac surgery	45.9% vs 32.8% (P < 0.05)	Statins decrease the rate of postoperative AF in patients undergoing cardiac surgery
Recurrences of AF after ECV (Colivicchi et al, 2004)	n = 476	Statin users vs non users.	0.72 (0.54-0.97) (P = 0.034)	Statins user had a reduction of recurrence at 1-year follow-up
Reduction of new-onset AF (Merckx et al, 2004)	n = 667	Statin users vs non users. Patients with CAD	(7.3% vs 11.6%)	Lower rate of new-onset AF during a mean follow-up of 6.5 years
Prevent recurrences of AF after ECV (Tveit et al., 2004)	Prospective, OL, R, multicenter n = 114	Pravastatin, 40 mg/day, vs no drug.	35% vs 33% (CI not available)	Pravastatin does not reduce the recurrence of AF after ECV at 6-weeks follow-up
AF after noncardiac thoracic surgery	Prospective n = 130	Statin user vs non users. Patients undergone major lung or	0.26 (0.08-0.82) (P = 0.022)	Preoperative use of statins reduces postoperative AF independent of CRP levels

(Amar et al, 2005)		esophageal resection		
Statins on paroxysmal AF (Dernellis & Panaretou, 2005)	SB, PC, R, prospective, parallel design n= 80	Atorvastatin (20-40 mg/day) users vs non users. Patients with paroxysmal AF	10% vs 65% (P < 0.001)	Atorvastatin reduces the incidence of paroxysmal AF
Reduction of AF after CABG (Marín et al, 2006)	n = 234	Statin users vs nonusers. Patients undergoing CABG	0.52 (0.28-0.96) (P = 0.038)	Statins protect against AF after CABG
Recurrence rates of AF after ECV (Ozaydin et al, 2006)	Prospective, R n = 99	Atorvastatin (10 mg/day) vs no drug. Persistent AF	0.23 (0.064-0.82) (P = 0.024)	Atorvastatin decreases the recurrence rate of AF after ECV
Patients with reduced LVEF at high risk of subsequent AF (Hanna et al., 2006)	National Registry n = 25,268 in SR	Lipid-lowering drugs users vs non users	0.69 (0.64-0.74)	In patients with reduced LVEF statin use is associated with a significant reduction in the prevalence of AF
Patients undergoing cardiac surgery with cardiopulmonary bypass (Patti et al., 2006)	Prospective, R, DB, PC N = 200	Atorvastatin (40 mg/day) initiated 7 days before surgery vs placebo	35% vs 57% 0.39 (0.18-0.85) (P = 0.017)	Atorvastatin reduces the incidence of postoperative AF and shortens the length of stay

AF: atrial fibrillation. CABG: coronary artery bypass grafting. CAD: coronary artery disease. CI: 95% confidence interval. CRP: C reactive protein. DB: double-blind. ECV: electrical cardioversion. LDL-C: low density lipoprotein-cholesterol. LVEF: left ventricular ejection fraction. OL: open-label. OR: odds ratio. PC: placebo-controlled. R: randomized. RRR: relative risk reduction. SB: single-blind. SR: sinus rhythm.

Table 3. Pleiotropic actions related to the antiarrhythmic effects of statins

1. Anti-inflammatory properties
  - ↓ macrophage number and activation
  - ↓ expression of adhesion molecules and leukocyte-endothelial interactions
  - ↓ inflammatory cytokine secretion by macrophages
  - Switch Th-1 to Th-2 type phenotype in lymphocytes and ↑ anti-inflammatory cytokine production
2. Stabilization of vulnerable plaques
  - ↑ SMC and collagen
  - ↓ MMPs in atheroma plaque
  - ↓ LDL uptake and oxidation by macrophages
3. Improvement of endothelial dysfunction
  - ↑ NO levels by stabilizing mRNA
  - ↑ NO bioavailability
  - ↑ expression and activity of NOS3
  - ↓ endothelin-1 expression
4. Antioxidant effects
  - ↓ oxidative stress and reactive oxygen species
5. Cardioprotective effects
  - ↓ cardiac and smooth muscle hypertrophy and cardiac fibrosis
6. Improvement of autonomic function
  - ↓ HRV and sympathetic function
7. Antithrombotic effects
  - ↓ platelet activation
  - ↓ coagulation factors
  - ↓ tissue factor expression
  - ↑ fibrinolytic activity (tPA/PAI-1 ratio)
  - ↑ thrombomodulin expression
8. Effects on cardiac ion channels and transporters
  - ↓ Na<sup>+</sup>-pump and Na<sup>+</sup>/Ca<sup>2+</sup> exchanger
  - ↓ I<sub>Ca,L</sub> and I<sub>Kr</sub>
  - ↓ hKv1.5, Kv4.3 and ImK channels
  - ↑ K<sub>ATP</sub> channels






Figure 2