

JOURNAL OF PALEONTOLOGY

DIMORPHINITES (AMMONOIDEA, JURASSIC, UPPER BAJOCIAN) IN THE PRECORDILLERA OF NORTHERN CHILE

SIXTO R. FERNÁNDEZ-LÓPEZ¹ AND GUILLERMO B. CHONG DIAZ²

¹Departamento de Paleontología, Facultad de Ciencias Geológicas (UCM) and Instituto de Geología Económica (CSIC-UCM), Madrid 28040, Spain, <sixto@geo.ucm.es>; and ²Departamento de Ciencias Geológicas, Universidad Católica del Norte, Antofagasta, Chile, <gchong@ucn.cl>



This pdf file is licensed for distribution in the form of electronic reprints and by way of personal or institutional websites authorized by the author(s).

A copyright license has been purchased to make this possible.

JOURNAL OF PALEONTOLOGY

The image features three ammonite fossils against a solid dark green background. One ammonite is positioned at the top right, another at the bottom left, and a third is partially visible behind the bottom one. All three fossils exhibit a distinct siphuncular canal and radial septa.

VOL 85 □ NO 3 □ MAY 2011 □ THE PALEONTOLOGICAL SOCIETY

DIMORPHINITES (AMMONOIDEA, JURASSIC, UPPER BAJOCIAN) IN THE PRECORDILLERA OF NORTHERN CHILE

SIXTO R. FERNÁNDEZ-LÓPEZ¹ AND GUILLERMO B. CHONG DIAZ²

¹Departamento de Paleontología, Facultad de Ciencias Geológicas (UCM) and Instituto de Geología Económica (CSIC-UCM), Madrid 28040, Spain, <sixto@geo.ucm.es>; and ²Departamento de Ciencias Geológicas, Universidad Católica del Norte, Antofagasta, Chile, <gchong@ucn.cl>

ABSTRACT—A new upper Bajocian ammonite assemblage containing the morphoceratids *Dimorphinites dimorphus* (d'Orbigny) and *Vigoriceras defrancei* (d'Orbigny) is reported from the circum-Pacific area. These ammonites were found at the top of the Torcazas Formation, in the Quebrada San Pedro area, Precordillera of northern Chile. Taphonomic, systematic, and paleobiogeographic data confirm these dimorphs were part of indigenous populations within the Tarapaca Basin, belonging to the same demic biospecies: *Dimorphinites defrancei* (d'Orbigny). The West Tethyan and East-Pacific distribution of *D. defrancei* corroborates the availability of the migratory seaway, the so-called Hispanic Corridor, between the western Tethys and the eastern Pacific Ocean during the upper Bajocian Parkinsoni Zone. The discovery of this upper Bajocian *Dimorphinites* assemblage provides a new biochronostratigraphic horizon in the Tarapaca Basin and an interoceanic correlation point for the latest Bajocian.

INTRODUCTION

SEVERAL AMMONITE assemblages containing morphoceratids have been discovered in the Quebrada San Pedro area, northern Chile, Región de Antofagasta, Comuna Sierra Gorda, approximately 40 km ESE of Sierra Gorda town and 75 km south of Calama city (Fig. 1). These assemblages were collected from three localities (12P, 13P and 14P) along an isolated outcrop that is exposed for 600 m. The outcrop is located 1 km north of the composite section described as Quebrada San Pedro and 10 km south of the classic localities of the Caracoles Mining District previously studied by numerous authors (Gottschke, 1878; Steinmann, 1881; Möricke, 1894; Westermann, 1967, 1981; Hillebrandt, 1970, 1973, 1977, 2001; Westermann and Riccardi, 1979, 1980; Jensen and Quinzio, 1979, 1981; Gröschke and Hillebrandt, 1985, 1994; Hillebrandt et al., 1986; Gröschke et al., 1988; Riccardi et al., 1990a, 1990b; Riccardi and Westermann, 1991a, 1991b; Fernandez-Lopez et al., 1994; Fernandez-Lopez and Chong Diaz, 2010).

This paper provides a systematic description of the morphoceratids from Quebrada San Pedro (Antofagasta, Chile), which includes two available European morphospecies *Dimorphinites dimorphus* (d'Orbigny, 1846) [M=macroconch] and *Vigoriceras defrancei* (d'Orbigny, 1845) [m=microconch], now regarded as dimorphs of *Dimorphinites defrancei* (d'Orbigny, 1845) [M + m] new combination. The specimens found in Chile provide important taphonomic, systematic, paleobiogeographic and biochronological information, and they have important paleogeographic and geochronologic implications.

GEOLOGICAL SETTING

The localities yielding the morphoceratids described in this paper occur in Jurassic strata that extend along the western edge of the Domeyko Range ($S\ 23^{\circ}30'-27^{\circ}$, Fig. 1). These deposits were originally part of the marine, back-arc Tarapaca Basin ($S\ 12^{\circ}-29^{\circ}$; Mordojoyich, 1981; Westermann and Riccardi, 1985; Mpodozis and Ramos, 1989; Carlotto et al., 2005; Vicente, 2005, 2006; Parent, 2006) or Domeyko Basin (Flint et al., 1993; Ardill et al., 1998; Hillebrandt et al., 2000).

In the Caracoles and Quebrada San Pedro areas, a succession of Jurassic, fossiliferous deposits belonging to the

Caracoles Group unconformably overlies Triassic volcanic and metamorphic rocks (Harrington, 1961; Garcia, 1967; Montaño, 1976; Ardill et al., 1998; Marinovic and Garcia, 1999). The lowest marine deposits of this group are brown, calcareous sandstones and fine-grained conglomerates belonging to the Coronado Formation (Fig. 2). This unit is overlain by shallow-marine, yellow and brown shales with intercalations of limestones and calcareous sandstones of the Torcazas Formation. These are in turn overlain by offshore, yellow and green shales with intercalations of limestones of the Mina Chica Formation. The Coronado, Torcazas and Mina Chica formations span the Bajocian through the Bathonian and display maximum thicknesses of 40, 100 and 75 m, respectively.

Lithostratigraphically, the new morphoceratid assemblages were collected from the top of the Torcazas Formation, below the Mina Chica Formation, in Quebrada San Pedro area. The sequence of the Torcazas Formation at this location consists of 10–15 m of gray and yellow silty limestones, wackestones to packstones and marls that are very fossiliferous in the upper part and are the source of several successive ammonite assemblages. The limestone beds vary in thickness from 15 to 40 cm, and they extend laterally several tens of meters. Most of the beds contain abundant resedimented and silicified macrofossils and display sharp and irregular bases, as well as normal grading of bioclasts, suggesting that carbonate inputs were affected by traction currents and scouring. The most abundant macrofossils are costate brachiopods, bivalves, ammonites, belemnites and gastropods. Isolated and fragmentary bones of decimetric or centimetric size are locally common. The most abundant ammonites are *Megasphaeroceras* Imlay, 1961 [M + m] spp.

At the uppermost beds of the Torcazas Formation, in this outcrop, Stephanoceratidae, Perisphinctidae and Oppeliidae are scarce, but *Cadomites* Munier-Chalmas, 1892 [M]-*Polyplectites* Mascke, 1907 [m], *Vermisphinctes* Buckman, 1920 [m]-*Prorsisphinctes* Buckman, 1921 [M] and *Oxycerites* Rollier, 1909 [M]-*Paroecotraustes* Spath, 1928 [m] are well represented. Morphoceratidae, Lissoceratidae and Strigoceratidae are very scarce, respectively represented by *Dimorphinites* Buckman, 1923 [M]-*Vigoriceras* Rioult, 1994 [m], *Lissoceras* Bayle, 1879 [M]-*Microlissoceras* Sturani, 1971 [m]




FIGURE 1—Geographic location of the Caracoles and Quebrada San Pedro Bajocian and Bathonian marine deposits.

and *Strigoceras* Quenstedt, 1886 [M]. Besides Ammonitina, Phylloceratina and Lytoceratina are extremely scarce. Among the morphoceratids of this stratigraphic interval, macroconchs of *Dimorphinites dimorphus* (d'Orbigny) and microconchs of

Vigoriceras defrancei (d'Orbigny) have been identified and are described below.

TAPHONOMIC ANALYSIS OF THE MORPHOCERATID AMMONITES

The morphoceratids described and figured in this paper (Figs. 3–5) were collected by one of the authors (F.-L.) from the uppermost limestone beds of the Torczas Formation during several field seasons, between 2006 and 2010.

The total number of morphoceratids found is nine (five macroconchs and four microconchs). Most of these specimens are preserved as calcareous internal molds of unflattened ammonites. They are filled with relatively homogeneous mudstones to wackestones in the outer preserved whorls, whereas the inner whorls are partially filled with macrocrystalline calcite. They are resedimented elements (displaced on the sea-bottom before their burial), showing structural continuity between the sedimentary infill and the enclosing sedimentary rock, without evidence of reworking (exhumation and displacement on the sea-bottom before their final burial; cf. Fernandez-Lopez, 2007). Fragmentary shells, lacking peristome borders, are dominant, generally without traces of rounding, encrustation, or bioerosion. Shells bearing signs of intrathalamic encrusting by organisms (cf. Tintant, 1984), such as serpulids, bryozoans, oysters and crinoids, are absent. Body chambers without apical sedimentary infilling, internal molds with a local infill channel on the ventral region, or siliciclastic pseudomorphosis of the shells are absent, too. Phragmocones with incomplete sedimentary infilling (i.e., hollow ammonites) and calcitic internal molds of phragmocones with recrystallized septa represent the dominant mode of preservation. No signs of biostratinomic or synsedimentary dissolution of the aragonitic remains have been recognized. Aragonitic shells were dissolved during later diagenesis. Empty ammonite camera, with incomplete sedimentary infilling and moldic porosity resulting from the dissolution of walls and septa were partially filled by spar cement giving rise to calcitic pseudomorphs of shells during late diagenesis.

Age	Lithostratigraphy	Column	Lithology	Setting
Middle Jurassic	Caracoles Formation	18 m	Grey and blue limestones with intercalations of green shales	Shallow marine
	Mina Chica Formation	75 m	Yellow and green shales with intercalations of limestones	Offshore
	Torczas Formation	100 m	Yellow and brown shales with intercalations of limestones and calcareous sandstones	Shallow marine
	Coronado Formation	40 m	Brown calcareous sandstones and fine-grained conglomerates	Shallow marine
Trias.	Agua Dulce Formation		Violet and red silicified volcanic rocks, including lava flows, tuffs and breccias	Volcanic

FIGURE 2—Age, stratigraphic column, lithology and depositional setting of the lithostratigraphic units mentioned in text (from Fernandez-Lopez and Chong Diaz, 2010).




FIGURE 3—*Dimorphinites dimorphus* (d'Orbigny) [macroconch] from Quebrada San Pedro (Antofagasta, Chile). Black asterisk marks the last septum of the phragmocone. The specimens were whitened with magnesium oxide prior to photography. 1–4, 13P84/4, ventral, left side, oral and right side views of complete specimen of pre-adult, showing peristome simple, sedimentary internal mold of outer whorls mostly without shell; 5, 6, 13P84/9, ventral and left side views of incomplete specimen of pre-adult, with part of the bodychamber, internal mold of fragmentary bodychamber, without shell, inner whorls not preserved; 7–10, 12P84/4, ventral, left side, oral and right side views of incomplete specimen, with septate whorls, internal mold of fragmentary phragmocone, with recrystallized inner whorls without shell.

These taphonomic observations are indicative of both a high rate of sediment accumulation and a low rate of sedimentation during biostratigraphic processes, due to tractive bottom-currents and sedimentary starvation in relatively deep-water environments, according to the model of Fernandez-Lopez (1997, 2000, 2007, 2008).

These morphoceratid internal molds do not exceed 50 mm in diameter (mean size = 25.1 mm). Most of them represent pre-adult individuals, lacking evidence of juveniles. Both macroconchs and microconchs are represented by populations dominated by pre-adults (taphonomic-populations of type-2 in Fernandez-Lopez, 1995; Fernandez-Lopez et al., 1999), suggesting autochthonous production of shells by indigenous populations of the marine basin (Fernandez-Lopez and Melendez, 1996). If the accumulation of shells had been produced by immigrants, or resulted from necroplanktic drift of shells from more open marine or exotic oceanic areas, it would probably be dominated by adults (taphonomic-populations of type-3 in Fernandez-Lopez, 1995; Fernandez-Lopez et al., 1999) and, in the case of drifted shells, the specimens would bear signs of intrathalamic encrusting by organisms (cf. Tintant, 1984; Callomon, 1985; Fernandez-Lopez and Melendez, 1996).




FIGURE 4—Septal suture of *Dimorphinites dimorphus* (d'Orbigny) [M], specimen 13P84/4 (cf. Figure 3.1–3.4) showing ontogenetic simplification at 70° of the last septum and 155° from the peristome. L=lateral lobe, U₂=second umbilical lobe.

SYSTEMATIC PALEONTOLOGY

Order AMMONITIDA Zittel, 1884
Superfamily ?PERISPINCTOIDEA Steinmann, 1890
Family MORPHOCERATIDAE Hyatt, 1900

The family Morphoceratidae is known in West Tethys, from the late Bajocian to the late Bathonian. In Europe, it is a common component of assemblages of the lower Bathonian Zigzag Zone (Mangold, 1970, 1997; Westermann, 1993; Galacz, 1995, 1999; Dietze et al., 1997, 2002). This family is known in the Andes from the lower Bathonian to upper Bathonian as a relatively scarce component of certain ammonite assemblages (Westermann and Hillebrandt, 1995; Riccardi and Westermann, 1999).

Morphoceratids comprise sphaerocone and planulate forms, with constrictions and irregular bundling or furcation of the ribs, and most have a smooth ventral band or sulcus. Septal sutures are relatively simple, with an unretracted suspensive lobe (cf. Arkell et al., 1957).

Evidence of sexual dimorphism in morphoceratids, with minute, evolute microconchs bearing lateral lappets, and larger, involute macroconchs with a contracted bodychamber and a simple peristome, has been proposed by diverse authors. In particular, the following taxonomic groups are regarded as dimorphs: *Morphoceras* Douville, 1881 [M]-*Ebrayiceras* Buckman, 1920 [m], *Polysphinctites* Buckman, 1922 [m]-*Asphinctites* Buckman, 1924 [M], *Dimorphinites* Buckman, 1923 [M]-*Vigoriceras* Rioult, 1994a [m], *Berbericeras* Roman, 1933 [M]-*Microberbericeras* Geczy and Galacz, 1998 [m] and *Pseudodimorphinites* Seyed-Emami (in Seyed-Emami et al., 1989) [M]-“*Polysphinctites*” [m].

Morphoceratids have most commonly been classified with the perisphinctaceans, due to the presence of distinct constrictions (Arkell, 1955; Westermann, 1956, 1958a; Mangold, 1970, 1997; Galacz, 1980, 2008; Donovan et al., 1981; Westermann and Callomon, 1988; Seyed-Emami et al., 1989;




FIGURE 5—*Vigoriceras defranceti* (d'Orbigny) [microconch] from Quebrada San Pedro (Antofagasta, Chile). Black asterisk marks the last septum of the phragmocone. The specimens were whitened with magnesium oxide prior to photography. Internal molds of bodychambers mostly without shell. Phragmocones with recrystallized calcitic whorls and mostly with calcitic pseudomorphic shell. 1–5, 13P84/7, nearly complete pre-adult specimen, with phragmocone and most of the bodychamber, without peristome; 6–10, 13P84/6, incomplete specimen of pre-adult, with phragmocone and most of the bodychamber, last portion of the bodychamber and peristome are missing; 11–15, 14P82/2, nearly complete specimen showing partial impression of peristome with lappets (at the right side).

TABLE 1—Measurements and abbreviations of the morphoceratid specimens from Quebrada San Pedro. Stratigraphic information is indicated by locality number, section letter and stratigraphic level: sections 12P, 13P and 14P; stratigraphic levels 82 and 84; /N=number of specimen. [M] and [m] indicate macroconch and microconch. Measurements (in mm) include maximum shell diameter at which the following measurements were made (D), whorl-height (H), whorl-height/diameter ratio (H/D), whorl-width (W), whorl-width/diameter ratio (W/D), umbilical diameter (U), umbilicus/diameter ratio (U/D), whorl-width/whorl-height ratio (W/H), primary ribs per half whorl (Ni/2), secondary ribs per half whorl (Ne/2), secondary ribs/primary ribs ratio (i) and number of constrictions per whorl (C).

Specimens		D	H	H/D	W	W/D	U	U/D	W/H	Ni/2	Ne/2	i	C
13P84/4	[M]	44.0	16.6	0.37	16.2	0.37	11.0	0.25	0.97	11	47	4.3	4
		35.7	16.2	0.45	20.6	0.58	2.0	0.06	1.27	11	36	3.3	—
12P84/4	[M]	18.5	7.5	0.40	10.3	0.56	4.2	0.23	1.37	—	23	—	3
		14.0	6.7	0.48	8.9	0.64	2.7	0.19	1.32	—	25	—	—
14P82/2	[m]	30.0	9.0	0.30	8.9	0.30	14.7	0.49	0.98	16	30	1.9	4
		22.5	7.7	0.34	8.6	0.38	9.1	0.40	1.11	—	—	—	4
13P84/8	[m]	25.0	9.3	0.37	9.0	0.36	8.4	0.34	0.96	—	—	—	5
		19.0	8.7	0.46	9.3	0.49	5.4	0.28	1.06	—	—	—	—
13P84/6	[m]	26.0	9.2	0.35	8.9	0.34	9.3	0.36	0.97	13	32	2.5	6
		20.0	8.1	0.40	8.7	0.43	5.9	0.30	1.07	12	33	2.8	4
13P84/7	[m]	26.0	8.5	0.33	8.2	0.31	11.0	0.42	0.96	—	37	—	4
		20.0	7.6	0.38	8.0	0.40	6.0	0.30	1.05	—	—	—	—

Besnosov and Mikhailova, 1991; Page, 1993; Rioult, 1994a, 1994b, 1994c; Geczy and Galacz, 1998; Riccardi and Westermann, 1999). However, studies of the septal suture have shown the presence of a dorsal Un lobe in the morphoceratids, which is characteristic of the Stephanocerataceae (Schindewolf, 1957, 1965; Westermann, 1958b; Hahn, 1970; Kullmann and Wiedmann, 1970; Tintant and Mouterde, 1981; Sandoval, 1983; Schlegelmilch, 1985; O'Dogherty et al., 2006). On the other hand, a possible sphaeroceratinid ancestor from the late Bajocian in Indonesia (Westermann, 1993) and transitional Tethyan forms from the stephanoceratid genus *Orthogarantiana* Bentz, 1928, or from a perisphinctid genus such as *Leptosphinctes* Buckman, 1920, or *Prorsosphinctes* Buckman, 1921, have been mentioned by Sandoval (1983) and Sturani (1964b, 1971) respectively.

Genus DIMORPHINITES Buckman, 1923

Type species.—“*Ammonites dimorphus*” d’Orbigny (1846, p. 410, pl. 141, figs. 1, 2, 5–8) by original designation (Buckman, 1923, pl. 377). This species is now regarded as a junior synonym of “*Ammonites defrancei*” (d’Orbigny, 1845), described and interpreted below.

Diagnosis.—Sphaeroconic shells, from small (~10 mm) to medium-size (~80 mm), with involute inner and intermediate whorls. Umbilicus from moderately narrow or occluded to wide. Venter rounded without smooth ventral band or groove. Aperture simple. Irregular bundling or furcation of the ribs, that pass over the venter rectiradiately or with a gentle forward inclination. Constrictions from the innermost whorls to the adult peristome. Sutures relatively simple, with suspensive lobe not retracted.

Occurrence.—*Dimorphinites* is known from Europe, northern Africa and southern Transcaucasus: England (Buckman, 1923; Donovan et al., 1981), France (d’Orbigny, 1845; Rioult, 1994a, 1994b, 1994c), Algeria (Roman, 1930), northern Italy (Sturani, 1964a, 1964b; Clari et al., 1984), Sicily (Wendt, 1964; Pavia et al., 2002; Galacz, 2008), Alps (Krystyn, 1972), Hungary (Galacz, 1980), Betic Basin (Sandoval, 1983, 1990; Sandoval et al., 2001; O’Dogherty et al., 2006), Iberian Basin (Fernandez-Lopez, 1985), Transcaucasus (Rostovtsev, 1985), Atlas (Enay et al., 1987), Carpathian (Wierzbowski et al., 1999; Schlögl et al., 2005), Tunisia (Soussi et al., 2000), Portugal (Fernandez-Lopez et al., 2006, 2009a) and Germany (Dietze and Bernt, 2009).

In the Tethyan areas, the *Dimorphinites* seems to be restricted to the uppermost Bajocian Parkinsoni Zone. Nevertheless, possible ancestral members of this taxonomic

group could be represented in the upper Bajocian Garantiana or Niortense zones of Venetian Prealps (Sturani, 1971) and Betic Basin (Sandoval, 1983). This genus also could be represented in the lower Bathonian Zigzag Zone of Sicily (Wendt, 1964; Pavia et al., 2002).

Discussion.—*Dimorphinites* has been regarded as macroconchiate with simple peristome, and dimorphic microconchs with lappets currently included in the (sub)genus *Vigoriceras* (Douville, 1881; Glangeaud, 1897; Makowski, 1963; Mangold, 1970, 1997; Galacz, 1980; Sandoval, 1983, 1990; Rioult, 1994a, 1994b; Dietze et al., 2002; Dietze and Bernt, 2009).

At least three nominal species of the genus *Dimorphinites* are available from Tethyan areas: 1) *D. dimorphus* (d’Orbigny, 1846, p. 410, pl. 141, figs. 1, 2, 5–8; lectotype designated by Rostovtsev, 1985 and Rioult, 1994b) from the “Oolite ferrugineuse of Bayeux” Formation, middle part of the Parkinsoni Zone (Densicosta-Subzone) of Bayeux or Port-en-Bessin (Calvados, France); 2) *D. compressus* Galacz (1980, p. 102, pl. 23, figs. 4, 6) from the upper part of the Parkinsoni Zone, of Gyenespuszta, Northern Bakony (Hungary); and 3) *D. centriglobus* Rioult (1994c, p. 131, pl. 50, fig. 8) from the middle part of the Parkinsoni Zone (Densicosta Subzone; cf. fig. 6) of Sainte-Honorine-des-Pertes (Calvados, France).

Dimorphinites differs from *Morphoceras* Douville (1881, p. 242; type species: “*Ammonites polymorphus*” d’Orbigny, 1846; renamed “*M. multiforme*” Arkell, 1951) by the absence of smooth ventral band. Other macroconchiate morphoceratids, such as *Asphinctites* Buckman (1924; type species: *A. recinctus* Buckman, 1924, TA-5, pl. 484), *Berbericeras* Roman (1933, p. 67; type species: *B. sekikense* Roman, 1933, pl. 2, fig. 15) and *Pseudodimorphinites* Seyed-Emami (in Seyed-Emami et al., 1989, type species: “*Morphoceras pingue*” Grossouvre, 1919, p. 391, pl. 14, fig. 7), are similar to *Dimorphinites* based on the lack of a ventral groove, but they show wider umbilicus and more evolute inner whorls. “*Dimorphinites nodifer*” Wendt (1964, pl. 21, fig. 5) may belong to the genus *Berbericeras*, as suggested by Dietze et al. (2002).

DIMORPHINITES DIMORPHUS (d’Orbigny, 1846) [M] Figures 3, 4, Table 1

Ammonites dimorphus d’ORBIGNY, 1845, p. 410, pl. 141, figs. 1, 2 (lectotype), 5–8.

Dimorphinites dimorphus d’Orbigny sp. BUCKMAN, 1923, pl. 377, figs. 1–3.

Morphoceras dimorphum d’Orbigny. ROMAN, 1930, p. 17, pl. 7, figs. 1, 2.

Dimorphinites dimorphus (d'Orbigny). STURANI, 1964b, p. 26, pl. 3, figs. 3, 4, text-fig. 22; KRYSTYN, 1972, p. 262, pl. 8, fig. 4; ROSTOVSEV, 1985, p. 162, pl. 44, figs. 7–10; SANDOVAL, 1990, p. 161, pl. 4, fig. 5; RIOUT, 1994b, p. 130, pl. 50, fig. 7; PAVIA et al., 2002, p. 47, fig. 26 (right); DIETZE AND BERNT, 2009, p. 6, figs. 1a–e, h–k.

Dimorphinites (Dimorphinites) dimorphus (d'Orbigny). GALACZ, 1980, p. 100, pl. 23, figs. 1–3, 5; SANDOVAL, 1983, p. 330, pl. 27, figs. 1, 2; WIERZBOWSKI et al., 1999, p. 48, figs. 17 (4, 5), 18 (1); SCHLÖGL et al., 2005, p. 344, pl. 13, fig. 2; GALACZ, 2008, p. 62, pl. 2, figs. 11, 12.

Description.—Inner and intermediate whorls are sphaeroconic, with nearly occluded umbilicus, convex flanks and venter. The coiling becomes evolute, with contracted body-chamber and flattened whorl-sides, at the last whorl of the phragmocone. The aperture is constricted, with simple peristome. Ribbing is blunt, irregular and dense, without smooth ventral band or groove, and weaker in the outer whorl. Constrictions, three to four per whorl, are narrow and deep commencing in the innermost whorls. In the largest specimen that shows umbilical egression, the bodychamber encompasses 125° (Fig. 3.1–3.4). All the specimens display recrystallized septa, although locally preserved. Sutures are relatively simple, with suspensive lobe not retracted (Fig. 4).

Material.—A relatively well-preserved (13P84/4), two incomplete (13P84/9, 12P84/4), and two fragmentary and crushed specimens (12P84/5–6), from 10 to 50 mm in diameter. The largest specimen (Fig. 3.1–3.4) show remains of the last two septa, lacking evidence of sutural crowding, at 125° of the peristome in the left side. The smallest specimen (Fig. 3.7–3.10) is completely septate.

Occurrence.—*Dimorphinites dimorphus* is the most common and widely distributed species of the genus, characterizing the middle part of the Bajocian Parkinsoni Zone from: England (Truellei Subzone cf. fig. 6; Buckman, 1923; Dietze and Bernt, 2009), France (Densicosta Subzone; Rioult, 1994b) and Venetian Prealps (Sturani, 1964a, 1964b). Certain problems of these chronostratigraphic assignments, however, are due to the diverse criteria of subzonation applied in the European basins within condensed assemblages (cf. Pavia, 1994, 2000). On the basis of the presence of *Parkinsonia rarecostata* (Buckman) in the same assemblages, the range of *D. dimorphus* have been extended into the lower part of the Parkinsoni Zone at least in Morocco (Enay et al., 1987), England (Page, 1993) and Germany (Ditze and Bernt, 2009). Moreover, this species has been mentioned in the upper part of the Parkinsoni Zone (Bomfordi Subzone) in the Venetian Prealps (Italy, Sturani, 1971), type area (France, Pavia and Martire, 2010) and Monte Kumeta (Sicily, Galacz, 2008). The species characterizes a Dimorphus Biohorizon (or subzone) in the uppermost Parkinsoni Zone of diverse Mediterranean basins (Galacz, 1980; Sandoval, 1983, 1990; Ouahhabi, 1994; Wierzbowski et al., 1999; O'Dogherty et al., 2006). It is known also with condensed assemblages at the Bajocian/Bathonian transition in Sicily (Wendt, 1964; Pavia et al., 2002).

Discussion.—The largest macroconch of Quebrada San Pedro (Fig. 3.1–3.4) belongs to a pre-adult individual. It shows possible signs of maturity, such as umbilical egression, flattening of the bodychamber, variocostation and simplification of the septal sutures (Fig. 4). Crowding of the last sutures and collared aperture, however, are missing and the bodychamber is extremely short. Within Tethyan *Dimorphinites*, the variability in adult size and bodychamber is extreme too, surpassing values of 80 mm and 360° respectively. Therefore, *Dimorphinites* developed umbilical egression during the late

pre-adult growth period, before maturity and cessation of growth, while the bodychamber length initially decreased and subsequently increased.

Dimorphinites compressus Galacz is distinguished from *D. dimorphus* by its less inflated morphology. *Dimorphinites centriglobus* Rioult displays a sphaeroconic phragmocone and a planulate bodychamber more pronounced. Smaller, adult *Dimorphinites* have been figured by Wendt (1964, p. 133, pl. 21, fig. 3) and Pavia et al. (2002, p. 48, fig. 26, bottom left).

Genus VIGORICERAS Rioult, 1994

Type species.—“*Ammonites defrancii*” d'Orbigny (1845, p. 239, pl. 129, figs. 7, 8) by original designation (Rioult, 1994a, p. 115).

Diagnosis.—Planulate shells, small-size (15–35 mm), with moderately evolute whorls. Umbilicus moderately narrow to wide. Venter rounded, without smooth ventral band or groove. Aperture with lateral spatulate lappets. Irregular bundling or furcation of the ribs, that pass over the venter rectiradiately or with a gentle forward inclination. Constrictions from the innermost whorls. Sutures relatively simple, with suspensive lobe not retracted.

Occurrence.—The (sub)genus *Vigoriceras* is known from several localities of Europe: France (d'Orbigny, 1845; Rioult, 1994a), England (Buckman, 1924) and Hungary (Galacz, 1980). These records are late Bajocian in age, exactly Parkinsoni Zone. However, microconchs of this taxonomic group could be represented in the Bajocian Garantiana or Niortense zones of Venetian Prealps (Sturani, 1971) and in the lower Bathonian Zigzag Zone of Sicily (Wendt, 1964; Pavia et al., 2002) and Iberian Range (Fernandez-Lopez, 1985).

Discussion.—At present, two nominal species of the genus *Vigoriceras* are available from the Western Tethys: 1) *V. defrancei* (d'Orbigny, 1845, p. 239, pl. 129, figs. 7, 8), from the “Oolite ferrugineuse of Bayeux” Formation, middle part of the Parkinsoni Zone (Densicosta Subzone) of Saint-Vigor (Calvados, France) and the type species of the genus; and 2) *V. dimorphoides* (Parona, 1896, p. 21, pl. 2, fig. 8; figured by Sturani, 1971, pl. 14, fig. 19) from the Monte Meletta assemblage (Niortense Zone, Venetian Prealps) but showing diverse sedimentary infill than other fossils from the same assemblage (according to Sturani, 1971, p. 175).

Vigoriceras differs from *Ebrayiceras* Buckman (1920, p. 23; type species: *E. ocellatum* Buckman, 1920, TA-3, pl. 173) by the absence of smooth ventral band. Other microconchiate morphoceratids, such as *Polysphinctites* Buckman (1922; type species: *P. polysphinctus* Buckman, 1922, TA-4, pl. 322), *Microberbericeras* Geczy and Galacz (1998; type species: “*Berbericeras (Microberbericeras) kopeki*” Geczy and Galacz, p. 499, pl. 2, figs. 17, 18) and the dimorphic counterpart of *Pseudodimorphinites* are similar to *Vigoriceras*, lacking of ventral groove, but they show wider umbilicus, more evolute inner whorls and more discoidal shell.

VIGORICERAS DEFRAUCEI (d'Orbigny, 1845) [m]

Figure 5, Table 1

Ammonites Defrancii d'ORBIGNY, 1845, p. 389, pl. 129, figs. 7, 8.
Ammonites Defrancei d'Orbigny. GLANGEAUD, 1897, p. 100, pl. 3, figs. 1–3.

Dimorphinites defrancii d'Orbigny sp. BUCKMAN, 1924, pl. 510.

Dimorphinites (unnamed subg.) *defrancii* (d'Orbigny). GALACZ, 1980, p. 103, pl. 22, fig. 3.

Dimorphinites (*Vigoriceras*) *defrancii* (d'Orbigny). RIOUT, 1994a, p. 115, pl. 50, figs. 5 (holotype) and 6.

European zones and subzones				South American (sub)zones and horizons	
NW European Province		Submediterranean Province		Mediterranean Province	
England, Normandy, Boulonnais, Lorraine, Alsace, northern Germany		Portugal, Iberian Basin, Aquitaine, Causses, Central-west France, Nièvre southern Jura, Maconnais, Ardeche, southern Germany		Betic Basin	
	TENUPLICATUS	AURIGERUS	Postpollubrum		
	Yeovilensis	Recinctus	Yeovilensis	Cadomites - Tilitidae	
	ZIGZAG	ZIGZAG	ZIGZAG	Morphoceras gulisanoi	?Zigzagiceras
	Macrescens	Macrescens	Macrescens		
	Convergens	Parvum	Dimorphitiformis		
Lower Bathonian	PARKINSONI	Bomfordi	Dimorphus		"Cobbanites"
	PARKINSONI	Truellei			
	PARKINSONI	Acris			
	GARANTIANA	Tetragona			
	GARANTIANA	Subgaranti			
	GARANTIANA	Dichotoma			
	NIORTENSE	Baculata			
	NIORTENSE	Polygyralis			
	NIORTENSE	Banksi			
Upper Bajocian	PARKINSONI				
	PARKINSONI				
	PARKINSONI				
	GARANTIANA				
	GARANTIANA				
	GARANTIANA				
	NIORTENSE				
	NIORTENSE				
	NIORTENSE				

FIGURE 6—Standard zonation for the upper Bajocian and lower Bathonian, showing correlation of European zones and subzones with South America zones, subzones and horizons (cf. Rioult et al., 1997; Mangold and Rioult, 1997; Hillebrandt, 2001; Callomon, 2003; O'Dogherty et al., 2006; Riccardi, 2008; Fernandez-Lopez et al., 2009b).

Vigoriceras defrancei (d'Orbigny) [m]. PAVIA et al., 2002, p. 47, figs. 26 (top left);

Vigoriceras defrancii (d'Orbigny). DIETZE AND BERNT, 2009, p. 6, figs. 1f, g.

Description.—Planulate, with moderately evolute whorls. Whorl-section ovaly shaped. Umbilicus moderately narrow to wide. Venter rounded, without smooth ventral band or groove. Aperture with lateral lappets. The ribbing is blunt, irregular and dense, weaker in the adult outer whorls. The constrictions, four to six per whorl, with forward inclination, are narrow and deep. Bodychamber ranges between 190 and

280°. The septa are commonly recrystallized. Sutures relatively simple, with suspensive lobe not retracted.

Material.—Three relatively well-preserved (13P84/7, 13P84/6, 14P82/2) and a fragmentary specimen (13P84/8), from 20 to 30 mm in diameter. All these specimens preserve phragmocone and bodychamber, but only one displays septal approximation (Fig. 5.6–5.10).

Occurrence.—*Vigoriceras defrancei* has been recorded in the middle part of the Bajocian Parkinsoni Zone from: England (Buckman, 1923; Dietze and Bernt, 2009), France (Rioult, 1994a) and northern Italy (Sturani, 1964a, 1964b). It has also been mentioned at the lower part of the Bajocian Parkinsoni Zone in Hungary (Galacz, 1980), in the upper part of the Parkinsoni Zone in Normandy (Pavia and Martire, 2010) and within condensed assemblages at the Bajocian/Bathonian transition in the Iberian Range (Fernandez-Lopez, 1985).

Discussion.—Only one of the four specimens from Quebrada San Pedro (Fig. 5.6–5.10) shows crowding and simplification of the last septal sutures, but associated with the shortest bodychamber. Most of these specimens, therefore, belong to pre-adult individuals.

Vigoriceras dimorphoides (Parona) display smaller size, with more globose shape and more involute coiling. Jaworski (1926, p. 262, pl. 2, fig. 4) determined an Argentine specimen as *Morphoceras* n. sp. cf. *M. defrancei* (d'Orbigny), from the Aalenian/Bajocian transition beds of Cerro Tricolor (Argentine). However, this specimen differs of the Morphoceratidae by the strong retraction of the oblique umbilical lobes and the more complex septal suture. It was proposed as holotype of a new species of Perisphinctidae: *Praeleptosphinctes jaworskii* Westermann (1956, p. 268; Westermann and Riccardi, 1982, p. 39, text-fig. 7a–e).

SYSTEMATIC, PALEOBIOGEOGRAPHIC AND BIOCHRONOSTRATIGRAPHIC INTERPRETATION

Most ammonite species of the late Bajocian and early Bathonian from South America are endemic to the Andean




FIGURE 7—Principal paleobiogeographic units during Bajocian/Bathonian transition, with location of *Dimorphinites* (asterisk) and *Vigoriceras* (circle) occurrences (paleogeography based on Moyne and Neige, 2007; Fernandez-Lopez et al., 2009b).

Province, or subprovinces, and the Eastern Pacific Subrealm (Westermann and Riccardi, 1980; Westermann, 1981, 1993, 2000, 2001; Riccardi et al., 1990a, 1990b, 1992, 1994; Riccardi and Westermann, 1991a, 1991b, 1999; Hillebrandt et al., 1992a, 1992b; Fernandez-Lopez et al., 1994, 2009b; Gröschke and Hillebrandt, 1993, 1994; Hillebrandt, 2001; Gröschke, 1996; Page, 2008; Riccardi, 2008). Upper Bajocian and lower Bathonian biozones and chronozone are, therefore, commonly defined by the regional, stratigraphic range of Pacific taxa (Fig. 6). The Andean Province was characterized by Eurycephalitinae, *Megasphaeroceras* being the most common member of the late Bajocian ammonite assemblages. Nevertheless, these eurycephalitins are associated with diverse taxa of West Tethyan affinity belonging to the families Stephanoceratidae (*Cadomites* [M]-*Polyplectites* [m], *Strenoceras* Hyatt, 1900 [m], *Orthogarantiana*? [M]), Spiroceratidae (*Spiroceras* Quenstedt, 1858), Perisphinctidae (*Leptosphinctes* Buckman, 1929 [M], *Prorsiphinctes* Buckman, 1921 [M], *Lobosphinctes* Buckman, 1923 [M]) and Lissoceratidae (*Lissoceras* Bayle, 1879 [M]).

In the Tarapaca Basin, the latest Bajocian *Dimorphinites defrancei* (d'Orbigny) [M + m] is represented by indigenous populations indistinguishable from its Tethyan counterparts, as far as can be seen from the scarce material. This species ranged throughout the Northwest European, Sub-Mediterranean and Mediterranean provinces of the West Tethyan Subrealm, the north-eastern Tethyan border and, according to this new record in the north Chile, the southeastern Pacific areas of the East-Pacific Subrealm (Fig. 7). Since *Dimorphinites defrancei* (d'Orbigny) [M + m] has only been known from the upper Bajocian Parkinsoni Zone in the western and north-eastern Tethyan border, this standard zone is now extended to the SE Pacific. Some associated taxa, such as *Vermiphinctes* [m]-*Prorsiphinctes* [M], *Strigoceras* [M] and *Oxycerites* [M] - *Paroecotraustes* [m] also indicate the uppermost Bajocian age and Tethyan affinities. Judging from work in progress, a fragmentary specimen of *Strigoceras* from the Dimorphinites assemblage of Quebrada San Pedro belongs to the species *S. truellei* (d'Orbigny) and corresponds also to the Bajocian Parkinsoni Zone.

The West Tethyan and East-Pacific distribution of *Dimorphinites defrancei* (d'Orbigny) [M + m] lend further credence to the presence of a marine connection between the western Tethys and the eastern Pacific Ocean, the hypothesized Hispanic Corridor (cf. Smith, 1983; Cantú-Chapa, 2001; Iturralde-Vinent, 2006). Moreover, the Mediterranean origin of the Morphoceratidae becomes uncertain, as the possibility exists of an origin along the southeastern Pacific margin.

ACKNOWLEDGMENTS

We thank T. Poulton (Geological Survey of Canada), an anonymous referee, P. Harries and B. Huber for their constructive comments and suggestions on the manuscript. One of us (F.L.) wishes to acknowledge the field assistance given by A. Hernandez Aravena (Antofagasta, Chile). This work was supported by the CGL2008-01273 (MICINN) project.

REFERENCES

- ARDILL, J., S. FLINT, G. CHONG, AND H. WILKE. 1998. Sequence stratigraphy of the Mesozoic Domeyko Basin, northern Chile. Journal of the Geological Society, London, 155:71–88.
- ARKELL, W. J. 1951–1959. A monograph of English Bathonian ammonites. Palaeontographical Society, London, 264 p.
- ARKELL, W. J., B. KUMMEL, AND C. W. WRIGHT. 1957. Mesozoic Ammonoidea, p. L80–L465. In R. C. Moore, (ed.), Treatise on Invertebrate Paleontology. Part L. Mollusca 4. Cephalopoda, Ammonoidea. Geological Society of America and University of Kansas Press, Lawrence.
- BAYLE, E. 1878–1879. Fossiles principaux des terrains jurassiques. Mémoire explicative de la Carte géologique de France, 4:23–99.
- BENTZ, A. 1928. Über Strenoceraten und Garantianen insbesondere aus dem Mittleren Dogger von Bielefeld. Jahrbuch der Preussischen Geologischen Landesanstalt, 49:138–206.
- BESNOV, N. V. AND I. A. MIKHAJOVA. 1991. Higher taxa of the Jurassic and Cretaceous Ammonitida. Paleontological Journal, 4:3–18. (In Russian)
- BUCKMAN, S. S. 1909–1930. Yorkshire Type Ammonites & Type Ammonites. Wesley & Son–Wheldon & Wesley, London, 790 pl.
- CALLOMON, J. H. 1985. The evolution of the Jurassic ammonite family Cardioceratidae. Special Papers in Paleontology, 33:49–90.
- CALLOMON, J. H. 2003. The Middle Jurassic of western and northern Europe: its subdivisions, geochronology and correlations. Geological Survey of Denmark and Greenland Bulletin, 1:61–73.
- CANTÚ-CHAPA, A. 2001. Mexico as the western margin of Pangea based on biogeographic evidence from the Permian to the Lower Jurassic, p. 1–27. In C. Bartolini, R. T. Buffler, and A. Cantú-Chapa (eds.), The western Gulf of Mexico Basin: Tectonics, sedimentary basins, and petroleum systems. American Association of Petroleum Geologists, Memoir, 75.
- CARLOTTO, V., L. CERPA, J. CÁRDENAS, J. QUISPE, AND G. CARLIER. 2005. Paleogeographic, structural and magmatic evidences for the existence of different lithospheric blocks in the Central Andes: Samples from southern Peru and northern Chile. 6th International Symposium on Andean Geodynamics, Barcelona, Abstracts, 146–149.
- CLARI, P. A., P. MARINI, M. PASTORINI, AND G. PAVIA. 1984. Il Rosso Ammonitico Inferiore (Baiociano-Calloviano) nei Monti Lessini settentrionali (Verona). Rivista Italiana di Paleontologia e Stratigrafia, 90:15–86.
- DIETZE, V. AND E. BERNT. 2009. Erstnachweis der Gattung *Dimorphinites* Buckman, 1923 (Morphoceratidae) aus dem Ober-Bajocium (Mittlerer Jura) Süddeutschlands. Jahreshefteder Gesellschaft für Naturkunde in Würtemberg, 165:5–11.
- DIETZE, V., T. KRIEGER, AND G. SCHWEIGERT. 1997. Über *Oeocoptychius subrefractus* (BUCKMAN), *Asaphinctites tenuiplicatus* (BRAUNS) und *Polysiphinctes secundus* (WETZEL) (Ammonoidea) aus dem Unter-Bathonium (Mittlerer Jura) der Oberpfalz (Nordost-Bayern, Süddeutschland). Stuttgarter Beiträge zur Naturkunde, 245:1–25.
- DIETZE, V., C. MANGOLD, AND R. B. CHANDLER. 2002. Two new species of *Berbericeras* Roman, 1933 (Morphoceratidae, Ammonoidea) from the Zigzag Bed (lower Bathonian, Zigzag Zone) of Waddan Hill (Broadwindsor, Dorset, Southern England). Stuttgarter Beiträge zur Naturkunde, 324:1–11.
- DONOVAN, D. T., J. H. CALLOMON, AND M. K. HOWARTH. 1981. Classification of the Jurassic Ammonitina. The Systematics Association, 18:101–155.
- DOUVILLE, H. 1881. Note sur l'*Ammonites pseudo-anceps* et sur la forme de son ouverture. Bulletin de la Société Géologique de France 3, 8, 1890:239–246.
- ENAY, R., C. MANGOLD, R. DU DRESNAY, AND M. RAKUS. 1987. Arrivals of Arabian origin among the ammonite faunas of Morocco during the Bajocian–Bathonian. Palaeogeography, Palaeoclimatology, Palaeoecology, 61:107–120.
- FERNANDEZ-LOPEZ, S. R. 1985. El Bajociense en la Cordillera Ibérica. Thesis Doctoral, Departamento de Paleontología, Universidad Complutense de Madrid, 850 p.
- FERNANDEZ-LOPEZ, S. R. 1995. Taphonomie et interprétation des paléoenvironnements. Geobios, M.S., 18:137–154.
- FERNANDEZ-LOPEZ, S. R. 1997. Ammonites, taphonomic cycles and stratigraphic cycles in carbonate epicontinental platforms. Cuadernos de Geología Ibérica, 23:95–136.
- FERNANDEZ-LOPEZ, S. R. 2000. Ammonite taphocycles in carbonate epicontinental platforms, p. 293–300. In R. L. Hall and P. L. Smith (eds.), Advances in Jurassic Research 2000. GeoResearch Forum, 6.
- FERNANDEZ-LOPEZ, S. R. 2007. Ammonoid taphonomy, palaeoenvironments and sequence stratigraphy at the Bajocian/Bathonian boundary on the Bas Auran area (Subalpine Basin, SE France). Lethaia, 40:377–391.
- FERNANDEZ-LOPEZ, S. R. 2008. Distinction between fill channels and abrasion furrows on ammonoid internal moulds. Geobios, 41:5–13.
- FERNANDEZ-LOPEZ, S. R. AND G. MELENDEZ. 1996. Phylloceratina ammonoids in the Iberian Basin during the Middle Jurassic: a model of biogeographical and taphonomic dispersal related to relative sea-level changes. Palaeogeography, Palaeoclimatology, Palaeoecology, 120:291–302.
- FERNANDEZ-LOPEZ, S. R. AND G. B. CHONG DIAZ. 2010. First circum-Pacific record of *Dimorphinites* (Ammonoidea, upper Bajocian) in the Precordillera of northern Chile. Earth Science Frontiers, 17:73–75.

- FERNANDEZ-LOPEZ, S. R., G. CHONG DIAZ, L. A. QUINZIO SINN, AND H.-G. WILKE. 1994. The upper Bajocian and Bathonian in the Cordillera de Domeyko, North-Chilean Precordillera: sedimentological and biostratigraphical results. *Geobios*, M.S., 17:187–201.
- FERNANDEZ-LOPEZ, S. R., M. H. HENRIQUES, A. LINARES, J. SANDOVAL, AND M. S. URETA. 1999. Aalenian *Tmetoceras* (Ammonoidea) from Iberia: taphonomy and palaeobiogeography, p. 395–417. In F. Olóriz, F. and F. J. Rodríguez Tovar (eds), *Cephalopods—Present and Past*, Kluwer Academic/Plenum Publishers, New York.
- FERNANDEZ-LOPEZ, S. R., M. H. HENRIQUES, AND C. MANGOLD. 2006. Ammonite succession at the Bajocian/Bathonian boundary in the Cabo Mondego region (Portugal). *Lethaia*, 39:253–264.
- FERNANDEZ-LOPEZ, S. R., G. PAVIA, E. ERBA, M. GUIOMAR, M. H. HENRIQUES, R. LANZA, C. MANGOLD, D. OLIVERO, AND D. TIRABOSCHI. 2009a. Formal proposal for the Bathonian GSSP (Middle Jurassic) in the Ravin du Bès Section (Bas-Auran, SE France). *Swiss Journal of Geosciences*, 102:271–295.
- FERNANDEZ-LOPEZ, S. R., G. PAVIA, E. ERBA, M. GUIOMAR, M. H. HENRIQUES, R. LANZA, C. MANGOLD, N. MORTON, D. OLIVERO, AND D. TIRABOSCHI. 2009b. The Global Boundary Stratotype Section and Point (GSSP) for base of the Bathonian Stage (Middle Jurassic), Ravin du Bès Section, SE France. *Episodes*, 32:222–248.
- FLINT, S., P. TURNER, E. J. JOLLEY, AND A. J. HARTLEY. 1993. Extensional tectonics in convergent margin basins: An example from the Salar de Atacama, Chilean Andes. *Geological Society of America Bulletin*, 105:603–617.
- GALACZ, A. 1980. Bajocian and Bathonian ammonites of Gyenespuszta, Bakony Mts., Hungary. *Geologia Hungarica*, Serie Palaeontologica, 39:1–227.
- GALACZ, A. 1995. Revision of the Middle Jurassic ammonite fauna from Csóka-hegy, Vértes Hills (Transdanubian Hungary). *Hantkeniana*, 1:119–125.
- GALACZ, A. 1999. A lower Bathonian ammonite fauna from Erice (Western Sicily). *Annales Universitatis Scientiarum Budapestinensis de Rolando Eotvos Nominaeae. Sectio Geologica*, 32:149–168.
- GALACZ, A. 2008. *Parkinsonia parkinsoni* Zone (upper Bajocian, Middle Jurassic) ammonites from Monte Kumeta (Western Sicily). *Bulletin della Società Paleontologica Italiana*, 47:51–69.
- GARCIA, F. 1967. Geología del Norte Grande de Chile. Sociedad Geológica de Chile. Simposio sobre el Geosinclinal Andino, 1–138.
- GECZY, B. AND A. GALACZ. 1998. Bathonian ammonites from the classic Middle Jurassic locality of Villany, South Hungary. *Revue de Paléobiologie*, 17: 479–511.
- GLANGEAUD, P. 1897. Sur la forme de l'ouverture de quelques ammonites. *Bulletin de la Société Géologique de France*, 25:99–106.
- GOTTSCHE, C. 1878. Ueber jurassische Versteinerungen aus der argentinischen Cordillere. *Palaeontographica*, supplement III, Lieferung II, Heft, 2:1–50.
- GRÖSCHKE, M. 1996. Zwei neue Ammoniten der Gattung *Iniskinites* aus dem Bathonium (Mitteljura) von Nordchile. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1996:140–146.
- GRÖSCHKE, M. AND A. VON HILLEBRANDT. 1985. Trias und Jura in der mittleren Cordillera Domeyko von Chile ($22^{\circ}30'$ – $24^{\circ}30'$). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 170:129–166.
- GRÖSCHKE, M. AND A. VON HILLEBRANDT. 1993. Die paläobiogeographischen Beziehungen zwischen Nordchile und Europa im Mittleren Jura. *Geologische Blätter für Nordost-Bayern*, 43:57–72.
- GRÖSCHKE, M. AND A. VON HILLEBRANDT. 1994. The Bathonian in Northern Chile. *Geobios*, M.S., 17:255–264.
- GRÖSCHKE, M., A. VON HILLEBRANDT, P. PRINZ, L. A. QUINZIO, AND H. G. WILKE. 1988. Marine Mesozoic Paleogeography in Northern Chile between S 21° – 26° . *Lecture Notes Earth Sciences*, 17:105–117.
- GROSSOUVRE A. DE. 1919. Bajocien-Bathonien dans la Nièvre. *Bulletin de la Société géologique de France*, sér. 4, 18, fasc. 8, 9:337–459.
- HAHN, W. 1970. Die Parkinsoniidae S. BUCKMAN und Morphoceratidae HYATT (Ammonoidea) des Bathoniums (Brauner Jura epsilon) im südwestdeutschen Jura. *Jahreshefte des Geologischen Landesamtes in Baden-Württemberg*, 12:7–62.
- HARRINGTON, H. L. 1961. Geology of parts of Antofagasta and Atacama Provinces, Northern Chile. *The American Association of Petroleum Geologists Bulletin*, 45:169–197.
- HILLEBRANDT, A. VON. 1970. Zur Biostratigraphie und Ammoniten-Fauna des südamerikanischen Jura (insbes. Chile). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 136:166–211.
- HILLEBRANDT, A. VON. 1973. Neue Ergebnisse über den Jura in Chile und Argentinien. *Münstersche Forschungen zur Geologie und Paläontologie*, 31/32:167–199.
- HILLEBRANDT, A. VON. 1977. Ammoniten aus dem Bajocien (Jura) von Chile (Südamerika). Neue Arten der Gattungen *Stephanoceras* und *Domeykoceras* n. gen. (Stephanoceratidae). *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie*, 17:35–69.
- HILLEBRANDT, A. VON. 2001. Ammonite stratigraphy of the Bajocian in Northern Chile. *Hantkeniana*, 3:49–87.
- HILLEBRANDT, A. VON, M. GRÖSCHKE, P. PRINZ, AND H.-G. WILKE. 1986. Marines Mesozoikum in Nordchile zwischen 21° und S 26° . *Berliner Geowissenschaftliche Abhandlungen*, 66:169–190.
- HILLEBRANDT, A. VON, P. SMITH, G. E. G. WESTERMANN, AND J. H. CALLOMON. 1992a. Ammonites zones of the circum-Pacific region, p. 247–272. In E. G. E. Westermann (ed.), *The Jurassic of the Circum-Pacific*. Cambridge University Press, Cambridge.
- HILLEBRANDT, A. VON, G. E. G. WESTERMANN, J. H. CALLOMON, AND R. L. DETTERMAN. 1992b. Ammonites of the circum-Pacific region, p. 342–359. In E. G. E. Westermann (ed.), *The Jurassic of the Circum-Pacific*. Cambridge University Press, Cambridge.
- HILLEBRANDT, A. VON, V. BARTSCH, A. BEBIOLKA, A. KOSSLER, W. KRAMER, H. G. WILKE, AND S. WITTMANN. 2000. The Paleogeographic Evolution in a Volcanic Arc/Back-Arc Setting during the Mesozoic in Northern Chile. *Zeitschrift für Angewandte Geologie*, Sonderheft SH 1, 87–93.
- HYATT, A. 1900. Cephalopoda, p. 502–592. In K. A. von Zittel, *Text-book of Palaeontology*. 1st English edition, translated by C.R. Eastman. Macmillan, London & New York.
- IMLAY, R. W. 1961. New genera and subgenera of Jurassic (Bajocian) ammonites from Alaska. *Journal of Paleontology*, 35:467–474.
- ITURRALDE-VINENT, M. A. 2006. Meso-Cenozoic Caribbean paleogeography: Implications for the historical biogeography of the region. *International Geology Review*, 48:791–827.
- JAWORSKI, K. 1926. La Fauna del Lias y Dogger de la Cordillera Argentina en la parte meridional de la Provincia de Mendoza. *Actas de la Academia Nacional de Ciencias de Córdoba*, 9:135–317.
- JENSEN, A. AND L. A. QUINZIO. 1979. Geología del Área de Pampa Elvira y contribución al conocimiento del Jurásico marino entre los $23^{\circ}00'$ y $23^{\circ}30'$ Latitud Sur y los $68^{\circ}45'$ y $69^{\circ}03'$ Longitud Oeste. II Región de Antofagasta, Chile. *Memoria Universidad Católica del Norte*, Departamento de Geociencias, Antofagasta, 141 p.
- JENSEN, A. AND L. A. QUINZIO. 1981. Hallazgo de Batoniano y descripción de facies sedimentarias marinas del Dogger en Quebrada San Pedro, Caracoles, II Región—Antofagasta (Chile). *Estudios geológicos*, 37:209–214.
- KRYSTYN, L. 1972. Die Oberbajocium- und Bathonium-Ammoniten der Klaus-Schichten des Steinbruches Neumühle bei Wien (Österreich). *Annalen des Naturhistorischen Museums in Wien*, 76:195–310.
- KULLMANN, J. AND J. WIEDMANN. 1970. Significance of sutures in phylogeny of Ammonoidea. *The University of Kansas, Paleontological Contributions*, 47:1–32.
- MAKOWSKI, H. 1963. Problem of sexual dimorphism in ammonites. *Palaeontologia Polonica*, 12, 1962:1–92.
- MANGOLD, C. 1970. Morphoceratidae (Ammonoidea, Perisphinctidae) bathoniens du Jura méridional, de la Nièvre et du Portugal. *Geobios*, 3:43–130.
- MANGOLD, C. 1997. Tendances évolutives chez les Morphoceratidés (Perisphinctacés, Ammonitina). *Cahiers de l'Université Catholique de Lyon*, 10:93–101.
- MANGOLD, C. AND M. RIOULT. 1997. Bathonian. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires*, 17:55–62.
- MARINOVIC, N. AND M. GARCIA. 1999. Hoja Pampa Unión, región de Antofagasta. 1:100.000. *Mapas Geológicos*, N° Mapa: M33. Servicio Nacional de Geología y Minería, Santiago de Chile.
- MASCKE, E. 1907. Die Stephanoceras-Verwandten in den Coronenschichten von Norddeutschland. *Inaugural-Dissertation*, Universität Göttingen, 38 p.
- MORDJOVICH, C. 1981. Sedimentary basins of the Chilean Pacific offshore. *American Association of Petroleum Geologists, Studies in Geology*, 12:732–749.
- MÖRICKE, W. 1894. Versteinerungen des Lias und Unteroolith von Chile. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, 9:1–100.
- MOYNE, S. AND P. NEIGE. 2007. The space-time relationship of taxonomic diversity and morphological disparity in the Middle Jurassic ammonite radiation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 248:82–95.
- MONTAÑO, J. M. 1976. Estudio geológico de la zona de Caracoles y áreas vecinas, con énfasis en el Sistema Jurásico, provincia de Antofagasta, II Región, Chile. *Tesis*, Universidad de Chile, Departamento de Geología, Santiago, 182 p.
- MPODOZIS, C. AND V. RAMOS. 1989. The Andes of Chile and Argentina. *Circum-Pacific Council for Energy and Mineral Resources, Earth Science Series*, 11:59–90.
- MUNIER-CHALMAS, E. C. P. A. 1892. Sur la possibilité d'admettre un dimorphisme sexuel chez les Ammonitidés. *Comptes redues de la Société Géologique de France*, 20, Sér. 3:170–174.

- O'DOGHERTY, L., J. SANDOVAL, A. BARTOLINI, S. BRUCHEZ, M. BILL, AND J. GUEX. 2006. Carbon-isotope stratigraphy and ammonite faunal turnover for the Middle Jurassic in the Southern Iberian palaeomargin. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 239:311–333.
- ORBIGNY, A. d'. 1842–1850. *Paléontologie française. Description zoologique et géologique de tous les animaux mollusques et rayonnés fossiles de France. Terrains oolitiques ou jurassiques*, T. 1. Masson, Paris, 642 p.
- OUAHABI, B. 1994. Le Lias et le Dogger inférieur des Beni Snassen orientaux (Maroc). Stratigraphie, paléontologie et dynamique du bassin. Comparaison avec les Monts d'Oujda et avec les régions limitrophes. Thesis University of Oujda, 496 p.
- PAGE, K. N. 1993. Mollusca: Cephalopoda (Ammonoidea: Phylloceratina, Lytoceratina, Ammonitina, Ancyloceratina), p. 213–228. In M. J. Benton (ed.), *The Fossil Record 2*. Chapman & Hall, London.
- PAGE, K. N. 2008. The evolution and geography of Jurassic ammonoids. *Proceedings of the Geologist's Association*, 119:35–57.
- PARENT, H. 2006. Oxfordian and late Callovian ammonite faunas and biostratigraphy of the Neuquén-Mendoza and Tarapacá basins (Jurassic, Ammonoidea, western South America). *Boletín del Instituto de Fisiografía y Geología*, 76:1–70.
- PARONA, C. E. 1896. Nuove osservazioni sopra la fauna e l'età degli strati con *Posidonia alpina* nei Sette Comuni. *Palaeontographia Italica*, 1:1–42.
- PAVIA, G. 1994. Taphonomic remarks on d'Orbigny's type-Bajocian (Bayeux, west France). *Miscellanea del Servizio Geologico Nazionale*, 5:93–111.
- PAVIA, G. 2000. Biostratigraphic framework and problems of the *P. parkinsoni* Zone, latest Bajocian. In A. Galacz (ed.), *Bajocian- Bathonian Working Groups Meeting*, Budapest, Abstracts, 31–32.
- PAVIA, G. AND L. MARTIRE. 2010. Indirect biostratigraphy in condensed successions: a case history from the Bajocian of Normandy (NW France). *Volumina Jurassica*, 7:67–76.
- PAVIA, G., L. MARTIRE, V. CANZONERI, AND C. D'ARPA. 2002. Stop 3—Rocca Chi Parra Quarry; a condensed Rosso Ammonitico succession: depositional and erosional geometries, neptunian dykes and ammonites assemblages. In P. di Stefano (coord.), *Pre-Symposium Field Trip A* (12–15 September 2002). An Introduction to the Jurassic geology of Western Sicily. In M. Santantonio (ed.), *6th International Symposium on the Jurassic System*, General Field Trip Guidebook, 42–48.
- QUENSTEDT, F. A. 1858. *Der Jura*. Laupp, Tübingen, 843 p.
- QUENSTEDT, F. A. 1883–1888. *Die Ammoniten des Schwäbischen Jura*. Schweizerbart, Stuttgart, 815 p.
- RICCARDI, A. C. 2008. El Jurásico de la Argentina y sus ammonites. *Revista de la Asociación Geológica Argentina*, 63:625–643.
- RICCARDI, A. C. AND G. E. G. WESTERMANN. 1991a. Middle Jurassic ammonite fauna and biochronology of the Argentine-Chilean Andes. Part III: Bathonian-Callovian *Eurycephalitinae*, *Stephanocerataceae*. *Palaeontographica*, 216:1–110.
- RICCARDI, A. C. AND G. E. G. WESTERMANN. 1991b. Middle Jurassic ammonite fauna and biochronology of the Argentine-Chilean Andes. Part IV: Bathonian-Callovian *Reineckeiiidae*. *Palaeontographica*, 216: 111–145.
- RICCARDI, A. C. AND G. E. G. WESTERMANN. 1999. An early Bathonian Tethyan ammonite fauna from Argentina. *Palaeontology*, 42:193–209.
- RICCARDI, A. C., G. E. G. WESTERMANN, AND S. ELMI. 1990a. The Middle Jurassic Bathonian-Callovian Ammonite Zones of the Argentine-Chilean Andes. *Geobios*, 22:553–597.
- RICCARDI, A. C., G. E. G. WESTERMANN, AND S. E. DAMBORENEA. 1990b. Middle Jurassic of South America and Antarctic Peninsula. *Newsletters on Stratigraphy*, 21:105–128.
- RICCARDI, A. C., C. A. GULISANO, J. MOJICA, O. PALACIOS, C. SCHUBERT, AND M. R. A. THOMSON. 1992. Western South America and Antarctica, p. 122–161. In E. G. E. Westermann (ed.), *The Jurassic of the Circum-Pacific*. Cambridge University Press, Cambridge.
- RICCARDI, A. C., S. E. DAMBORENEA, M. O. MANCENÍDO, AND S. C. BALLENT. 1994. Middle Jurassic biostratigraphy of Argentina. *Geobios M.S.*, 17:423–430.
- RIOUT, M. 1994a. *Dimorphinites (Vigoriceras) defrancii*, p. 115. In J.-C. Fischer (ed.), *Révision critique de la Paléontologie française d'Alcide d'Orbigny, 1. Céphalopodes jurassiques*. Milan & Barcelona (Masson), Paris.
- RIOUT, M. 1994b. *Dimorphinites dimorphus*, p. 130–131. In J.-C. Fischer (ed.), *Révision critique de la Paléontologie française d'Alcide d'Orbigny, 1. Céphalopodes jurassiques*. Milan & Barcelona (Masson), Paris.
- RIOUT, M. 1994c. *Dimorphinites centriglobus* n. sp., p. 131. In J.-C. Fischer (ed.), *Révision critique de la Paléontologie française d'Alcide d'Orbigny, 1. Céphalopodes jurassiques*. Milan & Barcelona (Masson), Paris.
- RIOUT, M., D. CONTINI, S. ELMI, AND J. GABILLY. 1997. Bajocien. *Bulletin du Centre de Recherches Elf Exploration Production, Mémoires*, 17:41–53.
- ROLLIER, L. 1909. Phylogénie des principaux genres d'ammonoïdes de l'Oolithique (Dogger) et de l'Oxfordien. *Archives des Sciences Physiques et Naturelles de Genève*, 28:611–623.
- ROMAN, F. 1930. Céphalopodes du Lias et du Jurassique moyen des régions d'Oujda et du Grand Atlas (Maroc Oriental). Protectorat de la République Française au Maroc. Direction générale des travaux publics. Service des Mines et de la Carte Géologique. Notes et mémoires, 22 p.
- ROMAN, F. 1933. Note sur le Bathonien inférieur du Djebel-es-Sekika près Nemours (Départ. d'Oran). *Bulletin de la Société Géologique de France*, 5. Sér., 3:59–73.
- ROSTOVTEW, K. O. 1985. The Jurassic deposits of the south part of the Transcaucasus. Interdepartmental Stratigraphic Commission of the USSR, *Transactions*, 15:1–188. (In Russian).
- SANDOVAL, J. 1983. *Bioestratigrafía y Paleontología (Stephanocerataceae y Perisphinctaceae) del Bajocense y Bathonense en las Cordilleras Béticas*. Thesis Doctoral, Universidad Granada, 613 p.
- SANDOVAL, J. 1990. A revision of the Bajocian divisions in the Subbetic Domain (southern Spain). *Memorie descrittive della Carta geologica d'Italia*, 40:141–162.
- SANDOVAL, J., L. O'DOGHERTY, AND J. GUEX. 2001. Evolutionary rates of Jurassic ammonites in relation to sea-level fluctuations. *Palaios*, 16:311–335.
- SCHINDEWOLF, O. H. 1957. Die Lobenlinie im System der Ammonoidea. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 1957:433–443.
- SCHINDEWOLF, O. H. 1961–1968. Studien zur Stammgeschichte der Ammoniten. *Abhandlungen der Akademie der Wissenschaften und der Literature, Mainz, Mathematisch-naturwissenschaftliche Klasse*, 1:1–901.
- SCHLEGELMILCH, R. 1985. *Die Ammonites des süddeutschen Doggers*. Gustav Fischer Verlag, Stuttgart, 284 p.
- SCHLÖGL, J., M. RAKÚS, C. MANGOLD, AND S. ELMI. 2005. Bajocian–Bathonian ammonite fauna of the Czorsztyn Unit, Pieniny Klippen Belt (Western Carpathians, Slovakia); its biostratigraphical and palaeobiogeographical significance. *Acta Geologica Polonica*, 55/4:339–359.
- SEYED-EMAMI, K., G. SCHAIRER, AND M. ALAVI-NAINI. 1989. Ammoniten aus der unteren Dalichai-Formation (Unterbathon) östlich von Semnan (SE-Alborz, Iran). *Müncher geowissenschaftliche Abhandlungen*, 15:79–91.
- SMITH, P. L. 1983. The Pliensbachian ammonite *Dayiceras dayiceroides* and Early Jurassic paleogeography. *Canadian Journal of Earth Science*, 20:86–91.
- SOUSSI, M., C. MANGOLD, R. ENAY, M. BOUGHDIRI, AND M. H. BEN ISMAIL. 2000. Le Jurassique inférieur et moyen de la Tunisie septentrionale; corrélations avec l'axe Nord-Sud et paléogéographie. *Geobios*, 33:437–446.
- SPATH, F. 1927–1933. Revision of the Jurassic cephalopod fauna of Kachh (Cutch). *Palaeontologia Indica*, N.S., 9, 2:1–945.
- STEINMANN, G. 1881. Zur Kenntnis der Jura- und Kreideformation von Caracoles (Bolivia). *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, 49:239–301.
- STEINMANN, G. 1890. Cephalopoda, p. 344–475. In G. Steinmann and L. Döderlein (eds.), *Elemente der Paläontologie*. Engelmann, Leipzig.
- STURANI, C. 1964a. La successione delle faune ad Ammoniti nelle formazioni mediogiurassiche delle Prealpi Venete occidentali (regione tra il Lago di Garda e la Valle del Brenta). *Memorie degli Istituti di Geologia e Mineralogia dell' Università di Padova*, 24:1–63.
- STURANI, C. 1964b. Ammoniti mediogiurassiche del Veneto. Faune del Bajociano terminale (Zona a Garantiana e a Parkinsoni). *Memorie degli Istituti di Geologia e Mineralogia dell' Università di Padova*, 24:1–43.
- STURANI, C. 1971. Ammonites and stratigraphy of the "Posidonia alpina" beds of the Venetian Alps (Middle Jurassic, mainly Bajocian). *Memorie degli Istituti di Geologia e Mineralogia dell' Università di Padova*, 28:1–190.
- TINTANT, H. 1984. Autochtonie ou allochtonie chez les Céphalopodes. *Mémoires de Géologie Université Dijon* 7, 1982:257–271.
- TINTANT, H. AND R. MOUTERDE. 1981. Classification et phylogénèse chez les ammonites jurassiques, p. 85–101. In J. Martinell (ed.), *International Symposium on "Concept and Method in Paleontology," Departament de Paleontologia. Universitat de Barcelona*.
- VICENTE, J. C. 2005. Dynamic paleogeography of the Jurassic Andean Basin: pattern of transgression and localisation of main straits through the magmatic arc. *Revista de la Asociación Geológica Argentina*, 60:221–250.

- VICENTE, J. C. 2006. Dynamic paleogeography of the Jurassic Andean Basin: pattern of regression and general considerations of main features. *Revista de la Asociación Geológica Argentina*, 61:408–437.
- WENDT, J. 1964. Stratigraphisch-palaontologische Untersuchungen im Dogger Westsiziliens. *Boletino della Società Paleontologica Italiana*, 2:57–145.
- WESTERMANN, G. E. G. 1956. Phylogenie der Stephanocerataceae und Perisphinctaceae des Dogger. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 103:233–279.
- WESTERMANN, G. E. G. 1958a. Ammoniten-Fauna und Stratigraphie des Bathonien NW-Deutschlands. *Beihefte zum Geologischen Jahrbuch*, 32:1–103.
- WESTERMANN, G. E. G. 1958b. The significance of septa and sutures in Jurassic Ammonite systematics. *Geological Magazine*, 95, 6:441–455.
- WESTERMANN, G. E. G. 1967. Sucesión de ammonites del Jurásico medio en Antofagasta, Atacama, Mendoza y Neuquén. *Revista de la Asociación Geológica Argentina*, 22:65–73.
- WESTERMANN, G. E. G. 1981. Ammonite biochronology and biogeography of the Circum Pacific Middle Jurassic. *The Systematics Association*, 18:459–498.
- WESTERMANN, G. E. G. 1993. Global bio-events in mid-Jurassic ammonites controlled by seaways. *The Systematics Association, Special Vol.* 47:187–226.
- WESTERMANN, G. E. G. 2000. Marine faunal realms of the Mesozoic: review and revision under the new guidelines for biogeographic classification and nomenclature. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 163:49–68.
- WESTERMANN, G. E. G. 2001. Modes of extinction, pseudo-extinction and distribution in Middle Jurassic ammonites: terminology. *Canadian Journal of Earth Science*, 38:187–195.
- WESTERMANN, G. E. G. AND J.H. CALLOMON. 1988. The Macrocephalitinae and associated Bathonian and early Callovian (Jurassic) ammonoids of the Sula Islands and New Guinea. *Palaeontographica*, 203:1–90.
- WESTERMANN, G. E. G. AND A. VON HILLEBRANDT. 1995. A late Bathonian morphoceratid (Jurassic, Ammonitina) from Peru. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie*, 35:27–37.
- WESTERMANN, G. E. G. AND A. C. RICCARDI. 1979. Middle Jurassic Ammonoid fauna and biochronology of the Argentine-Chilean Andes, Part II, Bajocian Stephanocerataceae. *Palaeontographica*, 164:85–188.
- WESTERMANN, G. E. G. AND A. C. RICCARDI. 1980. The upper Bajocian ammonite *Sirenoceras* in Chile: first circum-Pacific record of the Subfurcatum Zone. *Newsletters on Stratigraphy*, 9:19–29.
- WESTERMANN, G. E. G. AND A. C. RICCARDI. 1982. Ammonoid fauna from the Early and Middle Jurassic of Mendoza Province, Argentina. *Journal of Paleontology*, 56:11–41.
- WESTERMANN, G. E. G. AND A. C. RICCARDI. 1985. Middle Jurassic Ammonite Evolution in the Andean Province and Emigration to Tethys. *Lecture Notes in Earth Sciences*, 1:6–34.
- WIERZBOWSKI, A., M. JAWORSKA, AND M. KROBICKI. 1999. Jurassic (Upper Bajocian-lowest Oxfordian) ammonítico rosso facies in the Pieniny Klippen Belt, Carpathians, Poland: its fauna, age, microfacies and sedimentary environment. *Studia Geologica Polonica*, 115:7–74.
- ZITTEL, K. A. VON. 1884. Cephalopoda, p. 329–522. In K. A. von Zittel, *Handbuch der Paläontologie*. 1. Abteilung. Band 2. Lieferung 3. R. Oldenbourg, München and Leipzig.

ACCEPTED 21 OCTOBER 2010