



# Socioeconomic patterns and COVID-19 outcomes before, during and after the lockdown in Italy (2020)

Alberto Mateo-Urdiales<sup>a,b</sup>, Massimo Fabiani<sup>a</sup>, Aldo Rosano<sup>c</sup>, Maria Fenicia Vescio<sup>a</sup>,  
Martina Del Manso<sup>a,b</sup>, Antonino Bella<sup>a</sup>, Flavia Riccardo<sup>a</sup>, Patrizio Pezzotti<sup>a</sup>,  
Enrique Regidor<sup>d,e,\*</sup>, Xanthi Andrianou<sup>a,f</sup>

<sup>a</sup> Department of Infectious Diseases, Istituto Superiore di Sanità, Rome, Italy

<sup>b</sup> European Programme for Intervention Epidemiology Training (EPIET), European Centre for Disease Prevention and Control (ECDC), Stockholm, Sweden

<sup>c</sup> Servizio Statistico, Istituto nazionale per l'analisi delle politiche pubbliche, Rome, Italy

<sup>d</sup> Department of Public Health & Maternal and Child Health, Faculty of Medicine, Universidad Complutense de Madrid, Madrid, Spain

<sup>e</sup> CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

<sup>f</sup> Cyprus International Institute for Environmental and Public Health, Cyprus University of Technology, Limassol, Cyprus

## ARTICLE INFO

### Keywords:

COVID-19

HEALTH Inequalities

Deprivation

SOCIAL Epidemiology

## ABSTRACT

The objective was to investigate the association between deprivation and COVID-19 outcomes in Italy during pre-lockdown, lockdown and post-lockdown periods using a retrospective cohort study with 38,534,169 citizens and 222,875 COVID-19 cases. Multilevel negative binomial regression models, adjusting for age, sex, population-density and region of residence were conducted to evaluate the association between area-level deprivation and COVID-19 incidence, case-hospitalisation rate and case-fatality. During lockdown and post-lockdown, but not during pre-lockdown, higher incidence of cases was observed in the most deprived municipalities compared with the least deprived ones. No differences in case-hospitalisation and case-fatality according to deprivation were observed in any period under study.

## 1. Introduction

Italy has been one of the most affected European countries by the coronavirus disease 2019 (COVID-19) pandemic which spread out of Hubei (China) in the early months of 2020 (Riccardo et al., 2020). By the December 15, 2020 over 1,500,000 people had been diagnosed with the disease and over 60,000 had died from it (Istituto Superiore di Sanità, 2020). In order to control the spread of the infection and safeguard the national health system, the Italian government implemented a series of social distancing measures. On the 4th of March, primary and secondary education centres were closed, followed by a national lockdown implemented on the March 10, 2020, by which citizens were only allowed outside their homes for work -if considered essential workers- and to acquire basic need items (Consiglio dei Ministri, D, 2020; Presidenza del Consiglio, 2020a). This measure was eased on the 18th of May, when non-essential work was resumed (Presidenza del Consiglio, 2020b). The full measure was lifted, including travel restrictions between regions, on the 3rd of June. Other measures implemented during

this period include the need to keep 1-m distance between people and the mandatory use of face masks indoors and in places where social distancing may not be possible (Presidenza del Consiglio, 2020a). Employers were asked to keep remote working where possible and, if not possible, to ensure the safety of employees by enforcing social distancing and providing face masks (Presidenza del Consiglio, 2020a).

These measures have caused severe social and economic disruption across the country. Yet, it is not yet known whether the different periods of the pandemic, and the measures implemented, could have modulated the risk exposure to SARS-CoV-2 across the different socioeconomic groups in Italy. Studies analysing the impact of previous pandemics on the different socioeconomic groups have found contradictory results. For example, some authors found higher illiteracy rates to be associated with an increased risk of mortality during the 1918 pandemic in the US (Grantz et al., 2016), but others have reported no differences by socioeconomic status in New Zealand during the same pandemic (Summers et al., 1918). Similarly, the impact of the 2009 pandemic influenza has been found to be higher in lower socioeconomic groups in England

\* Corresponding author. Hospital Clínico San Carlos, C/ Martín Lagos s/n, 5ª planta ala norte, 28040, Madrid, Spain.

E-mail address: [enriqueregidor@hotmail.com](mailto:enriqueregidor@hotmail.com) (E. Regidor).

<https://doi.org/10.1016/j.healthplace.2021.102642>

Received 4 March 2021; Received in revised form 15 July 2021; Accepted 19 July 2021

Available online 29 July 2021

1353-8292/© 2021 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

(Rutter et al., 2020), but not in France (Mansiaux et al., 2015). With regards to COVID-19, it has been suggested that those living in the most deprived areas could be at higher risk of morbidity and mortality from COVID-19 (Bambra et al., 2020; Di Girolamo et al., 2021). This increased risk could be the consequence of a greater exposure to the virus mediated by the working and living conditions of those who suffer deprivation (Bambra et al., 2020; The Health Foundation. Wi, 2020; Patel et al., 2020a). It has been proposed that low-paid workers and those in manual occupations may be at increased exposure to SARS-CoV-2 compared with other occupations given that they are less likely to be able to work remotely, more likely to suffer from poor working conditions and more likely to live in crowded housing, among other factors (The Health Foundation. Wi, 2020; Patel et al., 2020a). Besides the increased risk in exposure, there is evidence that Non-Communicable Diseases (NCD), such as diabetes, cardiovascular and chronic respiratory diseases, are associated with deprivation (Mamo et al., 2020). As these diseases are risk factors for hospitalisation and mortality from COVID-19, it is plausible that rates of these outcomes are higher in the most deprived areas (Bambra et al., 2020). Yet, the published literature shows inconsistent results (Wachtler et al., 2020). Ecological studies carried out in the UK and US have found a positive association between deprivation and incidence, hospitalisation and mortality from SARS-CoV-2 infection (Williamson et al., 2020; Niedzwiedz et al., 2020; Nayak et al., 2020), but other studies have not found such association (Pollán et al., 2020; Carrat et al., 2020); and others have found that it is actually the wealthier groups who have been hit harder by COVID-19 (Abedi et al., 2020; Mukherji, 2020).

It is likely that the association between COVID-19 outcomes and socioeconomic variables is influenced by different social, cultural, economic and policy factors; as well as by epidemic dynamics that vary from country to country and within countries. For example, in Italy, incidence has been particularly high in the northern areas, which are wealthier than the centre and south of the country, especially during the first periods of the pandemic.

In this study, we aimed to investigate the association between COVID-19 related outcomes and the level of deprivation of the municipality of residence in the Italian population; and how this association changed throughout the different epidemic periods.

## 2. Methods

### 2.1. Study design

We conducted a retrospective cohort study using a contextual approach to evaluate the association between deprivation and COVID-19 incidence, as well as between deprivation and the risk of hospitalisation and death among COVID-19 cases; across Italian municipalities in the different periods of the epidemic (pre-lockdown, lockdown and post-lockdown). The study was carried out by analysing individual data and using the Italian deprivation index of the municipality of residence as a contextual measure of deprivation.

We described the methods and presented findings according to the reporting guidelines for observational studies that are based on routinely collected health data (The RECORD statement –checklist of items extended from the STROBE statement) (Supplementary Material 1) (Benchimol et al., 2015).

### 2.2. Data sources

We obtained individual data on cases, hospitalisations and deaths from the Italian integrated epidemiological surveillance system of COVID-19, which collects demographic, clinical and epidemiological data on all PCR confirmed cases of COVID-19 in the country (National Health Institute, 2020). From every case, we extracted information on age, sex, vital status, history of hospitalisation, whether or not they were healthcare workers, and their municipality of residence. For this last

variable we used the 2020 list of Italian municipalities as reported by the national institute of statistics (ISTAT) (Istituto Nazionale di Sta, 2020). As a measure of deprivation, we used the Italian municipality index of deprivation (Rosano et al., 2020).

We obtained estimates of the Italian population (stratified by region, municipality, age and sex) as well as the population density of Italian municipalities updated on the January 1, 2020 through the Italian institute of statistics (ISTAT) (Istituto Nazionale di Sta, 2020), assuming these remained unchanged during the study period.

### 2.3. Exposure, outcomes and potential confounders

We analysed the association between deprivation (exposure) and COVID-19 incidence, case-hospitalisation rate and case-fatality (outcomes). We used the index of deprivation as a contextual measure of deprivation. This index was built using information from the 2011 census on unemployment, educational attainment, percentage of rented housing, house overcrowding and percentage of single-parent families (Rosano et al., 2020). In the analysis, we categorized the index according to quintiles of its distribution among municipalities, with “one” being the least deprived and “five” the most deprived.

We considered as COVID-19 cases those who were tested positive for SARS-CoV-2 infection by RT-PCR. Among these, we considered hospitalisations and deaths occurring within 40 days of the date of sampling/diagnosis.

We considered age, sex, population density and region of residence, as potential confounders of the associations between the exposure and outcomes in each epidemic period. Age was categorized into three groups (0–49, 50–69 and over 70 years old). We used these cut-offs based on the observed changes in age's case-fatality as reported by routine surveillance data (Istituto Superiore di San, 2020). Population density was categorized into three levels (<54 people per km<sup>2</sup>, 54–106 people per km<sup>2</sup> and >106 people per km<sup>2</sup>).

We used the date of sampling/diagnosis of cases to assign them to each period studied (pre-lockdown, lockdown and post-lockdown). The lockdown in Italy was implemented on the 10th of March and was lifted on the 18th of May. We added seven days to these dates to account for the median time between infection and diagnosis –four days of incubation period and three days between symptom onset and diagnosis (Guan et al., 2020)–. Therefore, cases were assigned to the pre-lockdown period if they had a date of sampling/diagnosis between the February 20, 2020 and the March 16, 2020, to the lockdown period if the date was between the March 17, 2020 and the 24th of May; and to the post-lockdown if between the May 25, 2020 and the 15th of October.

### 2.4. Statistical analysis

The analysis was conducted using surveillance data extracted on the December 9, 2020.

We excluded from the analysis individuals living in municipalities with a population larger than 50,000 people, as we considered that the social deprivation index could not represent the reality of large municipalities. The threshold of 50,000 was set up based on previous studies who have analysed data at the level of Italian municipalities (Minichilli et al., 2017). We also excluded healthcare workers because, as they have a greater risk of being exposed to the virus than the general population and they are less likely to suffer from socioeconomic deprivation, they could confound the associations tested in this study. Finally, we excluded cases with incomplete information for the analysis. At the end, we included 222,875 cases (see Fig. 1), which represented 54.1 % of the total cases (ranging from 33.6 % to 91.9 % across the different regions). Cases were aggregated by 7624 municipalities, which represented a population of 38,534,169 (64.0 % of the total Italian population).

We described the main demographic characteristics by level of deprivation of the area of residence with counts and percentages. We conducted a descriptive analysis of the distribution of deprivation and




Fig. 1. Flow chart showing the selection of cases included in the analysis.

COVID-19 related outcomes. We calculated age-adjusted rates for each outcome by deprivation quintile, stratifying the results by sex and epidemic period. To adjust rates by age we used direct standardisation using the European Standard Population 2013 as reference (Pace et al., 2013). To calculate rates, we included in the denominator the number of person-days at risk in each period. When calculating incidence, persons living in municipalities included in the study were considered at risk until they were diagnosed with the infection or until the end of the period under study, whichever came first. When calculating case-hospitalisation and case-fatality rates, cases were considered at risk until their recovery/death. If the event did not happen, they were considered as exposed for 40 days.

Then, we carried out a multivariable analysis using negative-binomial regression models for each outcome to measure its association with the level of deprivation of the municipality of residence. We deemed this as the most appropriate method given the significant level of overdispersion ( $>1$ ).

We calculated one model for each outcome and period, in which the number of cases/hospitalisations/deaths was included as the dependent variable. We included the independent variables in three steps. First, we calculated the models including deprivation of the municipality of residence (exposure of interest) together with sex and age group. Then, we added the level of population density of the municipality of residence and in the final step we added the region of residence. We also included in the model random effects accounting for clustering at municipality level (random intercept only). The offset was the person-days at risk. The Intraclass Correlation Coefficient (ICC) (i.e., the proportion of variance explained by random effects) was used to evaluate the need to use multilevel models. To this purpose, we used the formula suggested by Nakagawa et al. for negative binomial models (Nakagawa et al., 2017).

Estimates are presented with the 95 % confidence intervals (CI). The analysis was carried out in R (version 4.0.2), using Rstudio (version 1.3.959) (The, 2018; tudio Team.tudio: In, 2015). We used the package glmmTMB for the multivariable analysis. The formula used for the

calculation of the models alongside the full list of the R packages used can be found in the Supplementary Material 2.

### Ethical statement

This study was conducted using data from the Italian national integrated COVID-19 surveillance routinely collected and analysed within the mandate of the Italian National Institute of Health. The scientific dissemination of COVID-19 surveillance data was authorised by the Italian Presidency of the Council of Ministers on the February 27, 2020 (Ordinance n. 640).

## 3. Results

Table 1 shows the demographic characteristics of the included population according to the variables of interest, a map with the geographical position of each Italian region can be found in the Supplementary Material 3.

### 3.1. Distribution of COVID-19 outcomes according to deprivation

In Italy, deprivation follows a north-south gradient, with the south concentrating a larger number of municipalities in the most deprived quintiles compared to the north. On the contrary, incidence of COVID-19 was higher in the north of the country, particularly during the pre-lockdown period, spreading more widely during the lockdown and post-lockdown periods (See Fig. 2).

Table 2 summarises the number of cases, hospitalisations and deaths by municipalities' deprivation quintiles, with their respective age-adjusted rates; stratified by sex and epidemic period. Incidence peaked during the lockdown period and decreased afterwards. During pre-lockdown and lockdown periods, higher incidence was observed in the municipalities belonging to the least deprived quintile (Q1) compared with those in the most deprived ones (Q5), in both females and males. However, this gradient inverted during the post-lockdown period, when

**Table 1**

Distribution (column %) of the Italian population included in the study (38,534,169 population) according to the characteristics of interest.

	Index of deprivation				
	Q1 (least deprived)	Q2	Q3	Q4	Q5 (most deprived)
<b>Females</b>					
<b>Age group</b>					
0–49	701,820 (49.0 %)	1,493,434 (51 %)	2,080,147 (51.6 %)	2,475,434 (51.7 %)	3,562,867 (55.3 %)
50–69	419,544 (29.3 %)	840,618 (28.7 %)	1,139,603 (28.3 %)	1,356,866 (28.3 %)	1,787,085 (27.8 %)
70 and over	309,753 (21.6 %)	596,217 (20.3 %)	807,980 (20.1 %)	957,769 (20 %)	1,089,894 (16.9 %)
	1,431,117 (100.0 %)	2,930,269 (100.0 %)	4,027,730 (100.0 %)	4,790,069 (100.0 %)	6,439,846 (100.0 %)
<b>Population density</b>					
<54.6 ppkm2	266,933 (18.7 %)	299,537 (10.2 %)	274,120 (6.8 %)	319,201 (6.7 %)	330,982 (5.1 %)
54.6–106 ppkm2	243,517 (17 %)	386,368 (13.2 %)	465,022 (11.5 %)	498,064 (10.4 %)	603,154 (9.4 %)
>106 ppkm2	920,667 (64.3 %)	2,244,364 (76.6 %)	3,288,588 (81.6 %)	3,972,804 (82.9 %)	5,505,710 (85.5 %)
	1,431,117 (100.0 %)	2,930,269 (100.0 %)	4,027,730 (100.0 %)	4,790,069 (100.0 %)	6,439,846 (100.0 %)
<b>Regions' grouped by geographical area</b>					
North <sup>1</sup>	1,312,378 (91.7 %)	2,327,889 (79.5 %)	2,709,840 (67.2 %)	2,543,374 (53.0 %)	840,279 (13.0 %)
Centre <sup>2</sup>	63,446 (4.4 %)	432,369 (14.8 %)	876,548 (21.8 %)	870,667 (18.2 %)	875,738 (13.6 %)
South and Islands <sup>3</sup>	55,293 (3.9 %)	170,011 (5.8 %)	441,342 (11 %)	1,376,028 (28.7 %)	4,723,829 (73.4 %)
	1,431,117 (100.0 %)	2,930,269 (100.0 %)	4,027,730 (100.0 %)	4,790,069 (100.0 %)	6,439,846 (100.0 %)
<b>Males</b>					
<b>Age group</b>					
0–49	735,841 (52.5 %)	1,558,283 (54.9 %)	2,165,134 (55.8 %)	2,575,729 (56.2 %)	3,682,631 (59.3 %)
50–69	424,539 (30.3 %)	828,766 (29.2 %)	1,107,808 (28.6 %)	1,300,344 (28.3 %)	1,689,939 (27.2 %)
70 and over	239,932 (17.1 %)	453,733 (16 %)	606,423 (15.6 %)	711,007 (15.5 %)	835,029 (13.5 %)
	1,400,312 (100.0 %)	2,840,782 (100.0 %)	3,879,365 (100.0 %)	4,587,080 (100.0 %)	6,207,599 (100.0 %)
<b>Population density</b>					
<54.6 ppkm2	267,081 (19.1 %)	294,422 (10.4 %)	269,540 (6.9 %)	313,750 (6.8 %)	327,828 (5.3 %)
54.6–106 ppkm2	239,837 (17.1 %)	376,023 (13.2 %)	450,146 (11.6 %)	480,715 (10.5 %)	591,378 (9.5 %)
>106 ppkm2	893,394 (63.8 %)	2,170,337 (76.4 %)	3,159,679 (81.4 %)	3,792,615 (82.7 %)	5,288,393 (85.2 %)
	1,400,312 (100.0 %)	2,840,782 (100.0 %)	3,879,365 (100.0 %)	4,587,080 (100.0 %)	6,207,599 (100.0 %)
<b>Regions' grouped by geographical area</b>					
North <sup>1</sup>	1,285,354 (91.8 %)	2,261,555 (79.6 %)	2,628,251 (67.8 %)	2,435,911 (53.1 %)	808,554 (13.1 %)
Centre <sup>2</sup>	61,693 (4.4 %)	413,442 (14.6 %)	828,163 (21.3 %)	830,278 (18.1 %)	846,282 (13.6 %)
South and Islands <sup>3</sup>	53,265 (3.8 %)	165,785 (5.8 %)	422,951 (10.9 %)	1,320,891 (28.8 %)	4,552,763 (73.3 %)
	1,400,312 (100.0 %)	2,840,782 (100.0 %)	3,879,365 (100.0 %)	4,587,080 (100.0 %)	6,207,599 (100.0 %)

<sup>1</sup> Includes Regions of: Piemonte, Valle d'Aosta, Liguria and Lombardia, Trentino-Alto Adige, Veneto, Friuli-Venezia Giulia and Emilia-Romagna; <sup>2</sup>Includes Regions of: Toscana, Umbria, Marche and Lazio; <sup>3</sup>Includes Regions of: Abruzzo, Molise, Campania, Puglia, Basilicata, Calabria, Sicilia and Sardegna.

incidence was somewhat higher in municipalities belonging to the most deprived quintile than in the least deprived ones, in females and males.

Case-hospitalisations rates were higher during the pre-lockdown period, decreased during lockdown and reaching its lowest level during the post-lockdown period, in females and males and in all deprivation groups. In the pre-lockdown period, the most and least deprived quintiles (Q1 and Q5) had the lowest case-hospitalisation rates. During lockdown, case-hospitalisation rate was lowest in municipalities

belonging to the least deprived quintile, but no clear gradient was observed. In the post-lockdown, similar rates were observed across deprivation groups.

Case-fatality rates also peaked during the pre-lockdown period and decreased afterwards, reaching its lowest levels during post-lockdown. No clear socioeconomic gradient was observed in any period. During pre-lockdown, cases living in the least and most deprived municipalities had the lowest case-fatality rates. During lockdown and post-lockdown, case-fatality rates were similar across all groups.

#### 4. Results from the multivariable analysis

Table 3 shows the main results from multilevel model adjusted for sex, age, population density and region of residence. The full results of the models, including the ICC, can be found in the Supplementary Material 4. During the pre-lockdown period, there was not a clear socioeconomic gradient in the incidence of COVID-19. Incidence was 20 % lower in municipalities belonging to the second least deprived quintile (Q2) compared with the least deprived one (Q1, IRR 0.80, 95 % CI: 0.70 to 0.91); and it was 17 % higher in municipalities belonging to the most deprived quintile, but not statistically significant (Q5, IRR 1.17, 95 % CI: 0.98 to 1.41). During lockdown, incidence was significantly higher in the most deprived quintile (Q5, IRR: 1.14, 95 % CI: 1.03 to 1.27) and in the second most deprived quintile (Q4, IRR: 1.18, 95 % CI: 1.08 to 1.29) compared with the least deprived one. These differences increased during post-lockdown, when municipalities in the most deprived quintile had 47 % higher incidence compared with the least deprived one (Q5, IRR: 1.47, 95 % CI: 1.32 to 1.63).

The results of the models using case-hospitalisation as the dependent variable show no gradient according to deprivation after full adjustment. During the pre-lockdown, cases living in the most deprived municipalities had the lowest hospitalisation rate (IRR: 0.68, 95 % CI: 0.51 to 0.92). No statistically significant differences with cases living in least deprived municipalities were observed in any other group and in any other period studied.

No differences in case-fatality rates were observed across groups during the pre-lockdown or lockdown periods after full adjustment. During the post-lockdown, compared with cases living in least deprived municipalities, case-fatality rates were higher in cases from municipalities belonging to the third quintile (Q3, IRR: 1.26, 95 % CI: 0.96 to 1.66), as well as in those from the most deprived municipalities (Q4, IRR: 1.20, 95 % CI: 0.90 to 1.59; Q5, IRR: 1.02, 95 % CI: 0.73 to 1.41), but these differences were not statistically significant.

#### 5. Discussion


##### 5.1. Statement of principal findings

Incidence of COVID-19 did not vary according to deprivation of the municipality of residence in the pre-lockdown, but as the epidemic affected more regions of Italy during the lockdown and post-lockdown, the incidence increased more during the lockdown and decreased less during the post-lockdown in the municipalities with the greatest deprivation. On the other hand, we did not observe differences in case-hospitalisation or case-fatality in any period according to deprivation of the municipality of residence.

##### 5.2. Comparison with other studies and possible explanations

Several studies have analysed the correlation between incidence of COVID-19 and socioeconomic indicators. The majority of those using area-level deprivation indexes as the socioeconomic measure have found higher incidence in the most deprived areas (Niedzwiedz et al., 2020; Disparities in the risk a, 2020; Liu et al., 2020; Ho et al., 2020; Whittle and Diaz-Artiles, 2020; Baena-Díez et al., 2020). Our findings suggest that, in Italy, municipality-level deprivation was only associated





**Fig. 2.** Geographical distribution of the cumulative number of cases per 100,000 persons during the periods of the epidemic in Italian municipalities with less than 50,000 population ( $n = 7624$ ), and distribution of quantiles of the index of deprivation.

with incidence in the lockdown and post-lockdown periods. The finding that the association between deprivation and COVID-19 outcomes varied throughout the different epidemic periods might be explained by different epidemiological and policy factors. The first cases reported of SARS-CoV-2 infection in Europe were associated with clusters affecting, generally, young adults and linked to travel to East Asia for work related reasons (Spiteri et al., 2019). Infection likely spread through the social networks of these first cases during the pre-lockdown period affecting, at the initial stage, a series of municipalities in the northern region of Lombardy. None of the 11 municipalities which formed part of the first “red zone” in Italy belong to the most deprived quintile, with all but one belonging to the first 3 quintiles (Coronavirus and firmato il D, 2020). Although the epidemic spread outside this initial “red zone”, it remained contained in the north during the pre-lockdown period and did not spread widely to other parts of the country, which could explain the lack of socioeconomic gradient observed.

During lockdown, even if due to the blanket measures implemented the epidemic kept limited to the north and centre of the country, it started to spread to a wider area and population. During this period, we observed that incidence increased more in the most deprived municipalities than in the least deprived ones. This finding coincides two previous studies carried out in the Italian region of Emilia-Romagna (Di Girolamo et al., 2021; Stivanello et al., 2020). On the other hand, during March 2020, the relative differences in mortality according to citizens’ educational level showed greater magnitude than the differences observed in March 2019 (apporto annuale 2, 2020). This finding suggests a higher incidence of COVID-19 in people of low socioeconomic position. The higher incidence in these citizens could explain the findings observed here during the lockdown and post-lockdown, since the proportion of people of low socioeconomic position is higher in the municipalities with greater deprivation.

During the post-lockdown period the epidemic spread widely through the country, even though at a lower rate of infection in the

population and at a much lower rate of hospitalisation and death compared with the previous period. It was in this period when we observed the largest differences in incidence between the most deprived quintile and the least deprived. It is possible that, with the spread of the epidemic, the socio-economic risk factors showed the role they play and the impact of deprivation on the epidemiology of epidemic became more apparent. These findings coincide with those reported in Germany, where it was found that initially incidence was higher in less deprived areas, but that the gradient inverted overtime, with higher incidence in more deprived areas from April to June (Socioeconomic inequalities, 2020); and differ from what has been observed in the UK, where, during the second wave that started in early September, incidence increased more in the least deprived areas compared with the more deprived ones (Office for National Statistics, 2020a). It is likely that the socioeconomic pattern of COVID-19 incidence varies depending on the country. For example, seroprevalence studies in Spain and France have not found a clear correlation between income and prevalence of SARS-CoV-2 infection (19,20), but a clear inverse gradient has been found in Brazil (Hallal et al., 2020).

There are several mechanisms that could explain the slightly higher incidence observed in the most deprived areas after the early period of the epidemic. People living in deprived areas may be more likely to live in crowded housing, which act as a barrier to isolating effectively positive cases and increases the likelihood of the infection being spread to other co-habitants (Bambra et al., 2020; The Health Foundation. Wi, 2020; Patel et al., 2020a); especially in a context where family transmission is the main setting of exposure (Signorelli et al., 2020). Equally, it has been proposed that those living in the most deprived areas are less likely to be able to work remotely (Bambra et al., 2020), and that they carry out manual jobs that may increase their exposure risk compared to those living in wealthier areas (The Health Foundation. Wi, 2020), which could explain why incidence inequalities were higher particularly during the post-lockdown period. It is also possible that, as prevalence of

**Table 2**

Age-adjusted rates (AAR) of cases, hospitalisations and deaths from SARS-CoV-2 infection in Italian municipalities by level of deprivation (Q1 least deprived, Q5 most deprived). Stratified by sex and epidemic period\*.

		Pre-lockdown		Lockdown		Post-lockdown	
Incidence							
ID	Sex	Number	AAR	Number	AAR	Number	AAR
Q1	Females	844	19.9	6023	51.8	2896	14.8
Q2		1412	16.7	10,949	47.1	7357	18.2
Q3		2198	18.6	15,459	48.5	9840	17.6
Q4		2136	15.4	18,257	48.2	11,025	16.6
Q5		918	5.3	8364	17.9	14,475	16.0
Q1	Males	1258	31.6	4674	44.3	3277	16.9
Q2		2441	31.2	8858	42.7	8090	20.4
Q3		3958	37.6	12,761	45.6	10,773	19.9
Q4		3831	31.0	14,291	43.3	11,795	18.3
Q5		1539	9.9	7379	17.4	15,797	17.8
Case-hospitalisation (within 40 days of diagnosis)							
ID	Sex	Number	AAR	Number	AAR	Number	AAR
Q1	Females	409	17.3	1314	4.9	284	2.8
Q2		844	28.4	2772	6.2	643	2.5
Q3		1363	29.6	4340	7.1	921	2.6
Q4		1381	34.5	4389	6.2	1094	2.8
Q5		512	24.9	2197	6.5	1151	2.5
Q1	Males	854	39.3	1740	10.5	359	3.9
Q2		1843	57.9	3975	15.0	802	3.6
Q3		2975	57.5	6037	16.6	1129	3.8
Q4		2898	59.2	6101	14.4	1366	4.1
Q5		1009	38.8	2886	13.0	1583	3.8
Case-fatality (within 40 days of diagnosis)							
ID	Sex	Number	AAR	Number	AAR	Number	AAR
Q1	Females	163	2.3	766	1.2	37	0.3
Q2		292	2.7	1266	1.1	94	0.3
Q3		515	3.0	1854	1.2	213	0.5
Q4		487	3.1	2280	1.2	194	0.4
Q5		156	2.4	807	1.1	161	0.3
Q1	Males	342	3.9	813	2.2	59	0.6
Q2		749	4.9	1553	2.4	112	0.5
Q3		1252	4.9	2342	2.5	179	0.6
Q4		1228	5.0	2598	2.5	199	0.6
Q5		396	4.2	1101	2.4	266	0.6
ID = Index of deprivation; AAR = Age Adjusted Rate per 1,000,000 person-days for incidence and per 1000 person-days for case-hospitalisations and case-fatality							

ID = Index of deprivation; AAR = Age Adjusted Rate per 1,000,000 person-days for incidence and per 1000 person-days for case-hospitalisations and case-fatality

\* Cases were allocated to the pre-lockdown period if they had a date of sampling/diagnosis between the 20th of February and the 16th of March, to the lockdown period if their date of sampling/diagnosis was between the 17th of March and the 24th of May and to the post-lockdown period if that date was between the 25th of May and the 15th of October. Cases were classified as hospitalized or dead if they had a date of recovery/death within 40 days of sampling/diagnosis.

chronic diseases is highest in deprived areas (Gnavi et al., 2020), people living in these municipalities are more likely to suffer from symptomatic COVID-19 and therefore seek testing than those living in the least deprived ones. Some studies have found higher hospitalisation and mortality rates in the most deprived areas of the UK (Williamson et al., 2020; Patel et al., 2020b; Rose et al., 2020; Office for National Statistics, 2020b; Lone et al., 2020), as well as in the US (Nayak et al., 2020; Azar et al., 2020). These findings could reflect the known social gradient in co-morbidities and risk factors for COVID-19 severity, such as obesity, diabetes, cardiovascular disease or respiratory diseases, by which those living in the most deprived areas suffer the biggest burden. However, a study in Scotland on mortality in hospitalized patients with COVID-19 infection did not find differences in case-fatality according to deprivation in the area of residence (Khan et al., 2020a). Equally, we did not find an association between the deprivation level of the municipality of residence and the risk of hospitalisation or death. One possible explanation is that, as reported in the literature, the extent of inequalities in mortality is less pronounced in southern Mediterranean countries, like Italy or Spain, than in the US or the UK (Mackenbach et al., 2008; Regidor et al., 2015). It is also possible that inequalities in mortality for COVID-19 are mainly driven by individual socioeconomic status, or that they occur mainly in big urban areas and (Baena-Díez et al., 2020), given that we measured deprivation as a contextual variable and excluded large municipalities, that our study did not capture this pattern.

Another possible explanation may lie in the methodological differences across studies, as there is significant heterogeneity in the

indicators used to assess deprivation and the geographical areas studied. This could explain why, for example, other studies carried out in the UK and the US have not found such association (Guha et al., 2020; Apea et al., 2021; Khan et al., 2020b). Further studies would help in clarifying the individual and contextual influence of deprivation on COVID-19 outcomes.

### 5.3. Strengths and weaknesses of the study

This is the first study analysing the relation between COVID-19 and inequalities in Italy. To the best of our knowledge, it is also the first study analysing the association between deprivation and various COVID-19 outcomes through the various epidemic periods. Using individual data allowed us to classify each case according to the variables studied and to keep the maximum possible disaggregation level, as well to adjust the analysis for several individual characteristics and contextual factors other than level of deprivation of the municipality of residence.

The association between deprivation and COVID-19 is complex and is likely to be influenced by contextual and individual factors. One of the limitations of our study is that we did not have an individual measure of deprivation and, thus, we could not test the interaction between deprivation at the contextual and individual levels. In any case, we included municipalities with very different population sizes. It is likely that the deprivation index represents better the context in those with smaller population than in the larger ones, where different realities may exist

**Table 3**

Incidence Rate Ratios (IRR) and 95 % confidence interval (95 % CI) of the results of the multilevel negative binomial regression analysis for the association between COVID-19 related outcomes and deprivation in Italian municipalities. Adjusted for sex, age, population density and region of residence.

	Pre-lockdown	Lockdown	Post-lockdown
<b>Incidence IRR [95%CI]</b>			
Q1 (least deprived)	Ref	Ref	Ref
Q2	0.80 [0.71–0.91]	0.95 [0.88–1.03]	1.12 [1.03–1.21]
Q3	0.92 [0.80–1.05]	1.01 [0.93–1.09]	1.11 [1.02–1.20]
Q4	1.00 [0.87–1.16]	1.18 [1.08–1.29]	1.16 [1.06–1.27]
Q5 (most deprived)	1.17 [0.98–1.41]	1.14 [1.03–1.27]	1.47 [1.32–1.63]
<b>Case-hospitalisation (within 40 days of diagnosis) IRR [95%CI]</b>			
Q1 (least deprived)	Ref	Ref	Ref
Q2	0.88 [0.71–1.09]	1.00 [0.85–1.16]	0.92 [0.77–1.08]
Q3	0.88 [0.71–1.09]	0.93 [0.79–1.09]	0.95 [0.81–1.13]
Q4	0.81 [0.65–1.03]	0.86 [0.73–1.03]	0.99 [0.83–1.19]
Q5 (most deprived)	0.68 [0.51–0.92]	0.89 [0.72–1.11]	0.99 [0.81–1.22]
<b>Case-fatality (within 40 days of diagnosis) IRR [95%CI]</b>			
Q1 (least deprived)	Ref	Ref	Ref
Q2	0.96 [0.82–1.12]	0.94 [0.86–1.02]	0.94 [0.71–1.25]
Q3	1.01 [0.86–1.17]	0.94 [0.86–1.02]	1.26 [0.96–1.66]
Q4	1.06 [0.90–1.24]	0.94 [0.86–1.02]	1.20 [0.90–1.59]
Q5 (most deprived)	0.92 [0.75–1.13]	0.95 [0.85–1.07]	1.02 [0.73–1.41]
ICC	0.522**	0.389**	0.384**

ICC: Intra Class Correlation Coefficient.

\* Random intercepts were included in the models to account for clustering of observations at the municipality level.

\*\* Likelihood Ratio test < 0.05.

inside the same municipality. On the other hand, we cannot infer the findings at the individual level since we would incur the ecological fallacy bias, by assuming that socioeconomic status is homogeneous in all residents living in the same municipality. Another limitation is that we did not have data on the number of COVID-19 tests done by municipality, or the number of cases ascertained out of the total estimated. We know that this changed through time, thereby, the number of cases in each period should be taken with caution. During the pre-lockdown and lockdown periods, particularly, there was a limited COVID-19 testing capacity in Italy affecting to a greater extent the areas with higher incidence. It is possible that access to testing varied according to deprivation, with those living in the most deprived areas being less likely to access testing than those living in more affluent municipalities, which could confound our results towards underestimation of inequalities during these periods. Also, we measured the level of deprivation of the cases' municipality of residence. However, we do not know if cases acquired the infection in these municipalities or elsewhere. There are, also, other factors which could confound of the association between deprivation and the outcomes for which we did not have data to adjust, such as ethnicity or occupation.

Finally, deprivation is a complex concept difficult to measure. The deprivation index we used takes into account five characteristics (namely: low level of education, being unemployed, living in rent, living in crowded house, living in a single-parent family), but there may be other important components not captured by the index.

## 6. Conclusions

The COVID-19 pandemic has had a large impact on the Italian population in terms of morbidity and mortality. The impact, however, has not been homogeneous across the different population subgroups. In terms of deprivation, we found an increased incidence of COVID-19 in the most deprived municipalities during lockdown and post-lockdown. We did not find differences in case-hospitalisation rates or case-fatality rates across deprivation groups in any epidemic period.

## Funding

This study was partially funded by EU grant 874850 MOOD and is catalogued as MOOD 026. The contents of this publication are the sole responsibility of the authors and don't necessarily reflect the views of the European Commission.

## Declaration of competing interest

All authors declare no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years, no other relationships or activities that could appear to have influenced the submitted work.

## Acknowledgments:

We would like to thank the COVID-19 working group from the ISS and the Regional Authorities:

Maria C. Rota, Matteo Spuri, Daniele Petrone, Stefania Bellino, Paolo D'Ancona, Stefano Boros, Ornella Punzo, Antonietta Filia, Maria R. Castrucci, Alessandra Ciervo, Corrado Di Benedetto, Stefania Giannitelli, Paola Stefanelli, Marco Tallon, Roberta Urciuoli (Istituto Superiore di Sanità);

Regional representatives: Manuela Di Giacomo (Abruzzo); Michele La Bianca (Basilicata); Anna D. Mignuolo (Calabria); Angelo D'Argenzio (Campania); Erika Massimiliani (Emilia- Romagna); Tolinda Gallo (Friuli Venezia Giulia); Paola Scognamiglio (Lazio); Camilla Sticchi (Liguria); Danilo Cereda (Lombardia); Daniel Fiacchini (Marche); Francesco Sforza (Molise); Maria G. Zuccaro (P.A. Bolzano); Pier P. Benetollo (P.A. Trento); Daniela Tiberti (Piemonte); Maria Chironna (Puglia); Maria A. Palmas (Sardegna); Salvatore Scondotto (Sicilia); Lucia Pecori (Toscana); Anna Tosti (Umbria); Mauro Ruffier (Valle D'Aosta); Filippo Da Re (Veneto).

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.healthplace.2021.102642>.

## References

- Abedi, V., Olulana, O., Avula, V., Chaudhary, D., Khan, A., Shahjouei, S., et al., 2020 Sep 1. Racial, economic, and health inequality and COVID-19 infection in the United States [cited 2020 Dec 22]. *J Racial Ethn Heal Disparities* 1–11. <https://doi.org/10.1007/s40615-020-00833-4>.
- ISTAT. Rapporto annuale 2020-La situazione del Paese [Annual report 2020-the country's situation] [Internet]. 2020 [cited 2021 Jul 5]. <https://www.istat.it/it/archivio/244848>.
- Apea, V., Wan, Y., Dhairyawan, R., Puthucherry, Z., Pearce, R., Orkin, C., Prowle, J., et al., 2021. Ethnicity and outcomes in patients hospitalised with COVID-19 infection in East London: an observational cohort study. *BMJ Open* 11 (1). <https://doi.org/10.1136/BMJOPEN-2020-042140>.
- Azar, K.M.J., Shen, Z., Romanelli, R.J., Lockhart, S.H., Smits, K., Robinson, S., et al., 2020 Jul 1. Disparities in outcomes among COVID-19 patients in A large health care system in California [cited 2020 Sep 16]. *Health Aff* 39 (7), 1253–1262. <http://www.healthaffairs.org/doi/10.1377/hlthaff.2020.00598>.
- Baena-Díez, J.M., Barroso, M., Cordeiro-Coelho, S.I., Díaz, J.L., Grau, M., 2020 Aug 7. Impact of COVID-19 outbreak by income: hitting hardest the most deprived [cited 2020 Sep 16]. *J Public Health (Bangkok)* 42 (4). <https://doi.org/10.1093/pubmed/fdaa136>. <https://academic.oup.com/jpubhealth/advance-article/doi/10.1093/pubmed/fdaa136/5881845>.
- Bambra, C., Riordan, R., Ford, J., Matthews, F., 2020 Jun 13. The COVID-19 pandemic and health inequalities [cited 2020 Sep 16]. *JECH J Epidemiol Community Health* 2020–214401. <http://jech.bmj.com/>.
- Benchimol, E., Smeeth, L., Guttman, A., Harron, K., Moher, D., Petersen, I., et al., 2015. The REporting of studies conducted using observational routinely-collected health data (RECORD) statement. *PLoS Med* 12 (10) <https://doi.org/10.1371/JOURNAL.PMED.1001885>.
- Carrat, F., De Lamballerie, X., Rahib, D., Blanché, H., Lapidus, N., Artaud, F., et al., 2020. Title Seroprevalence of SARS-CoV-2 among adults in three regions of France following the lockdown and associated risk factors: a multicohort study. *medRxiv*, 20195693. <https://doi.org/10.1101/2020.09.16.20195693>.

- Consiglio dei Ministri, 2020. Decreto del presidente del consiglio dei ministri 26 aprile 2020: Ulteriori disposizioni attuative del decreto-legge 23 febbraio 2020. n. 6, recante misure urgenti in materia di contenimento e gestione dell'emergenza epidemiologica da COVID-19, applicabili [Internet]. Rome. <http://www.trovanorme.salute.gov.it/norme/dettaglioAtto?id=73916&completo=true>.
- Coronavirus, firmato il Dpcm 1 marzo 2020 | [Internet]. [cited 2021 Jan 10]. [www.governo.it](http://www.governo.it). <http://www.governo.it/it/articolo/coronavirus-firmato-il-dpcm-1-marzo-2020/14210>.
- Di Girolamo, C., Bartolini, L., Caranci, N., Moro, M., 2020. Socioeconomic inequalities in overall and COVID-19 mortality during the first outbreak peak in Emilia-Romagna Region (Northern Italy) [cited 2021 Jan 21] *Epidemiol Prev* 44 (Suppl. 2), 5–6. <https://pubmed.ncbi.nlm.nih.gov/33412821/>.
- [cited 2020 Dec 22] Disparities in the Risk and Outcomes of COVID-19 [Internet], 2020. [www.facebook.com/PublicHealthEngland](https://www.facebook.com/PublicHealthEngland).
- Gnavi, R., Picariello, R., Pilutti, S., Di Monaco, R., Oleandri, S., Costa, G., 2020 Sep 1. Epidemiology in support of intervention priorities: the case of diabetes in turin (piedmont region, northern Italy). *Epidemiol. Prev.* 44 (5–6), 172–178.
- Grantz, K.H., Rane, M.S., Salje, H., Glass, G.E., Schachterle, S.E., Cummings, D.A.T., 2016 Nov 29. Disparities in influenza mortality and transmission related to sociodemographic factors within Chicago in the pandemic of 1918 [cited 2020 Dec 30] *Proc Natl Acad Sci U S A* 113 (48), 13839–13844. <https://pubmed.ncbi.nlm.nih.gov/27872284/>.
- Guan, W., Ni, Z., Hu, Y., Liang, W., Ou, C., He, J., et al., 2020 Apr 30. Clinical characteristics of coronavirus disease 2019 in China [cited 2020 Dec 7] *N Engl J Med* 382 (18), 1708–1720. <https://www.nejm.org/doi/10.1056/NEJMoa2002032>.
- Guha, A., Bonnu, J., Dey, A., Addison, D., 2020 Apr 22. Community and Socioeconomic Factors Associated with COVID-19 in the United States: zip code level cross sectional analysis [cited 2020 Dec 22];2020.04.19.20071944 medRxiv Prepr Serv Heal Sci. <https://doi.org/10.1101/2020.04.19.20071944>.
- Hallal, P.C., Hartwig, F.P., Horta, B.L., Silveira, M.F., Struchiner, C.J., Vidaletti, L.P., et al., 2020 Nov 1. SARS-CoV-2 antibody prevalence in Brazil: results from two successive nationwide serological household surveys [cited 2020 Dec 22] *Lancet Glob Heal* 8 (11), e1390–e1398. [www.thelancet.com/lancetgh](https://www.thelancet.com/lancetgh).
- Ho, F.K., Celis-Morales, C.A., Gray, S.R., Katikireddi, V., Niedzwiedz, C.L., Hastie, C., et al., 2020 Dec 22. Modifiable and Non-modifiable Risk Factors for COVID-19: Results from UK Biobank. <https://doi.org/10.1101/2020.04.28.20083295>.
- Istituto Nazionale di Statistica, 2020 Sep 16. Resident Population on 1st January [Internet]. <http://dati.istat.it/Index.aspx?lang=en&SubSessionId=54e65c42-b372-4a37-a15e-2736b21f5ac8>.
- Istituto Superiore di Sanità, 2020 May 12. EpiCentro. Integrated Surveillance of COVID-19: Main National Data [Internet]. <https://www.epicentro.iss.it/coronavirus/sars-cov-2-sorveglianza-dati>.
- Khan, K.S., Torpiano, G., McLellan, M., Mahmud, S., 2020 Aug 13. The impact of socioeconomic status on 30-day mortality in hospitalized patients with COVID-19 infection [cited 2020 Dec 22];jmv.26371 *J Med Virol*. <https://onlinelibrary.wiley.com/doi/abs/10.1002/jmv.26371>.
- Khan, K.S., Torpiano, G., McLellan, M., Mahmud, S., 2020 Aug 13. The impact of socioeconomic status on 30-day mortality in hospitalized patients with COVID-19 infection [cited 2020 Oct 13];jmv.26371 *J Med Virol*. <https://onlinelibrary.wiley.com/doi/abs/10.1002/jmv.26371>.
- Liu, S.H., Liu, B., Li, Y., Norbury, A., 2020 Dec 22. Time Courses of COVID-19 Infection and Local Variation in Socioeconomic and Health Disparities in England. <https://doi.org/10.1101/2020.05.29.20116921>.
- Lone, N.I., McPeake, J., Stewart, N.I., Blayney, M.C., Seem, R.C., Donaldson, L., et al., 2020. Influence of Socioeconomic Deprivation on Interventions and Outcomes for Patients Admitted with COVID-19 to Critical Care Units in Scotland: A National Cohort study-NC-ND License. *Lancet Reg Heal - Eur* [Internet]. <https://doi.org/10.1016/j.lanepe.2020.100005> [cited 2021 Jan 10];0(0):100005. Available from: <http://creativecommons.org/licenses/by-nc-nd/4.0/>.
- Mackenbach, J.P., Stirbu, I., Roskam, A.-J.R., Schaap, M.M., Menvielle, G., Leinsalu, M., et al., 2008 Jun 5. Socioeconomic inequalities in health in 22 European countries [cited 2020 Oct 16] *N Engl J Med* [Internet] 358 (23), 2468–2481. <http://www.nejm.org/doi/abs/10.1056/NEJMsa0707519>.
- Mamo, C., Marinacci, C., Demaria, M., Mirabelli, D., Costa, G., 2005. Factors other than risks in the workplace as determinants of socioeconomic differences in health in Italy [cited 2020 Sep 16] *Int J Occup Environ Health* 11 (1), 70–76. <https://pubmed.ncbi.nlm.nih.gov/15859194/>.
- Mansiaux, Y., Salez, N., Lapidus, N., Setbon, M., Andreoletti, L., Luerue-Ville, M., et al., 2015 Mar 1. Causal analysis of H1N1pdm09 influenza infection risk in a household cohort [cited 2020 Dec 30] *J Epidemiol Community Health* 69 (3), 272–277. <https://doi.org/10.1136/jech-2014-024678>.
- Minichilli, F., Santoro, M., Bianchi, F., Caranci, N., De Santis, M., Pasetto, R., 2017. Evaluation of the use of the socioeconomic deprivation index at area level in ecological studies on environment and health. *Epidemiol. Prev.* 41 (3–4), 187–196.
- Mukherji, N., 2020 Jul 14. The social and economic factors underlying the incidence of COVID-19 cases and deaths in US counties [cited 2020 Dec 22]; 2020.05.04.20091041 medRxiv. <https://doi.org/10.1101/2020.05.04.20091041>.
- Nakagawa, S., Johnson, P.C.D., Schielzeth, H., 2017 Sep 1. The coefficient of determination R<sup>2</sup> and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded [cited 2021 Jan 10] *J R Soc Interface* 14 (134). <https://pubmed.ncbi.nlm.nih.gov/28904005/>.
- National Health Institute, 2020 Sep 16. COVID-19 integrated surveillance: key national data [Internet]. <https://www.epicentro.iss.it/en/coronavirus/sars-cov-2-integrated-surveillance-data>.
- Nayak, A., Islam, S., Mehta, A., Ko, Y.-A., Patel, S., Goyal, A., et al., 2020. Impact of social vulnerability on COVID-19 incidence and outcomes in the United States, 04.10.20060962 medRxiv Prepr Serv Heal Sci [Internet]. 2020 Apr 17 [cited 2020 Dec 22]. <https://doi.org/10.1101/2020.04.10.20060962>.
- Niedzwiedz, C.L., O'Donnell, C.A., Jani, B.D., Demou, E., Ho, F.K., Celis-Morales, C., et al., 2020 May 29. Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort study using UK Biobank [cited 2020 Dec 22] *BMC Med* 18 (1), 160. <https://bmcmecine.biomedcentral.com/articles/10.1186/s12916-020-01640-8>.
- Office for National Statistics. Coronavirus (COVID-19) Infection Survey [Internet]. 2020 [cited 2021 Jan 15]. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionnsinthecommunityinengland/characteristicsofpeopletestingpositiveforcovid19inenglandseptember2020>.
- Office for National Statistics. Deaths Involving COVID-19 by Local Area and Socioeconomic Deprivation [Internet]. [cited 2020 Jul 8]. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsinvolvecovid19bylocalareasanddeprivation/deathsoccurringbetween1marchand31may2020>.
- Pace, M., Cayotte, E., Agafitei, L., Zupanec, T., Wojtyniak, B., Gissler, M., et al., 2013. Revision of the European Standard Population : report of Eurostat's task force. 2013 edition. Publications Office.
- Patel, J.A., Nielsen, F.B.H., Badiani, A.A., Assi, S., Unadkat, V.A., Patel, B., et al., 2020 Sep 16. Poverty, Inequality and COVID-19: the Forgotten Vulnerable [Internet], vol. 183. Public Health. Elsevier B.V.; 2020, pp. 110–111. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7221360/>.
- Patel, A.P., Paranjpe, M.D., Kathiresan, N.P., Rivas, M.A., Khera, A.V., 2020 Jul 6. Race, socioeconomic deprivation, and hospitalization for COVID-19 in English participants of a national biobank [cited 2020 Dec 22] *Int J Equity Health* [Internet] 19 (1), 114. <https://equityhealth.biomedcentral.com/articles/10.1186/s12939-020-01227-y>.
- Pollán, M., Pérez-Gómez, B., Pastor-Barriuso, R., Oteo, J., Hernán, M.A., Pérez-Olmeda, M., et al., 2020 Aug 22. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study [cited 2020 Sep 16] *Lancet* [Internet] 396 (10250), 535–544. [https://doi.org/10.1016/S0140-6736\(20\)31483-5](https://doi.org/10.1016/S0140-6736(20)31483-5).
- Presidenza del Consiglio dei Ministri, 2020a. Ulteriori disposizioni attuative del decreto-legge del 23 febbraio 2020, n.6, recante misure urgenti in materia di contenimento e gestione dell'emergenza epidemiologica da COVID-19 [Internet]. Roma. <http://www.governo.it/sites/new.governo.it/files/DPCM4MARZO2020.pdf>.
- Presidenza del Consiglio dei Ministri. Comunicato stampa del Consiglio dei Ministri n. 46 [Internet]. [cited 2020 Sep 16]. Available from: [www.governo.it/it/articolo/comunicato-stampa-del-consiglio-dei-ministri-n-46/14610](http://www.governo.it/it/articolo/comunicato-stampa-del-consiglio-dei-ministri-n-46/14610).
- Regidor, E., Vallejo, F., Giraldez-García, C., Ortega, P., Santos, J.M., Astasio, P., et al., 2015 Aug 29. Low mortality in the poorest areas of Spain: adults residing in provinces with lower per capita income have the lowest mortality [cited 2020 Dec 22] *Eur J Epidemiol* [Internet] 30 (8), 637–648. <https://pubmed.ncbi.nlm.nih.gov/25773751/>.
- Riccardo, F., Ajelli, M., Andrianou, X.D., Bella, A., Del Manso, M., Fabiani, M., et al., 2020 Dec 10. Epidemiological characteristics of COVID-19 cases and estimates of the reproductive numbers 1 month into the epidemic, Italy, 28 January to 31 March 2020 [cited 2021 Jan 17] *Eurosurveillance* 25 (49), 2000790. <https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2020.25.49.2000790>.
- Rosano, A., Pacelli, B., Zengarini, N., Costa, G., Cislighi, C., Caranci, N., 2020 Mar 1. Aggiornamento e revisione dell'indice di deprivazione italiano 2011 a livello di sezione di censimento. *Epidemiol. Prev.* 44 (2–3), 162–170.
- Rose, T.C., Mason, K., Pennington, A., McHale, P., Taylor-Robinson, D.C., Barr, B., 2020 May. Inequalities in COVID19 Mortality Related to Ethnicity and Socioeconomic Deprivation [Internet]. medRxiv [cited 2020 Jul 8]. Cold Spring Harbor Laboratory Press. <https://www.medrxiv.org/content/10.1101/2020.04.25.20079491v1>.
- Rutter, P.D., Mytton, O.T., Mak, M., Donaldson, L.J., 2012. Socio-economic disparities in mortality due to pandemic influenza in England [cited 2020 Jul 7] *Int J Public Health* 57 (4), 745–750. <https://pubmed.ncbi.nlm.nih.gov/22297400/>.
- Signorelli, C., Odone, A., Stirparo, G., Cereda, D., Gramegna, M., Trivelli, M., et al., 2020 Nov 20. SARS-CoV-2 transmission in the Lombardy Region: the increase of household contagion and its implication for containment measures [cited 2020 Dec 22] *Acta Biomed* 91 (4), 2020195. [www.actabiomedica.it](https://www.actabiomedica.it).
- Socioeconomic Inequalities in the Risk of SARS-CoV-2 Infection – First Results from an Analysis of Surveillance Data from Germany [Internet]. [cited 2020 Dec 22]. Available from: <https://edoc.rki.de/handle/176904/6996>.
- Spieteri, G., Fielding, J., Diercke, M., Campese, C., Enouf, V., Gaymard, A., et al., 2020 Mar 5. First cases of coronavirus disease 2019 (COVID-19) in the WHO European region, 24 January to 21 February 2020 [cited 2020 May 12] *Eurosurveillance* 25 (9), 2000178. <https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2020.25.9.2000178>.
- Stivanello, E., Perlangeli, V., Resi, D., Marzaroli, P., Pizzi, L., Pandolfi, P., 2020 Sep 11. COVID-19 cases before and after the “i stay at home” decree, bologna local health authority [cited 2021 Jan 10] *Italy. Acta Biomed* 91 (3), 1–7. [www.actabiomedica.it](https://www.actabiomedica.it).
- Summers, J.A., Wilson, N., Baker, M.G., Shanks, D.G., 2010 Dec. Mortality risk factors for pandemic influenza on New Zealand troop ship, 1918 [cited 2020 Dec 30] *Emerg Infect Dis* 16 (12), 1931–1937. <https://pubmed.ncbi.nlm.nih.gov/21122224/>.
- The, R., 2018. Project of Statistical Computing [Internet]. [www.r-project.org](http://www.r-project.org).
- The Health Foundation, 2020 Jul 8. Will COVID-19 be a watershed moment for health inequalities? <https://www.health.org.uk/publications/long-reads/will-covid-19-be-a-watershed-moment-for-health-inequalities>.
- RStudio Team, 2015. RStudio: Integrated Development for R. [Internet]. RStudio, Boston. <http://www.rstudio.com/>.
- Wachtler, B., Michalski, N., Nowossadeck, E., Diercke, M., Warendorf, M., et al., 2020. Socioeconomic inequalities and COVID-19 – A review of the current international



- literature. *Journal of Health Monitoring* 5 (S7), 3–17. <https://doi.org/10.25646/7059>.
- Whittle, R.S., Diaz-Artiles, A., 2020 Sep 4. An ecological study of socioeconomic predictors in detection of COVID-19 cases across neighborhoods in New York City [cited 2020 Dec 22] *BMC Med* 18 (1), 271. <https://bmcmecine.biomedcentral.com/articles/10.1186/s12916-020-01731-6>.
- Williamson, E.J., Walker, A.J., Bhaskaran, K., Bacon, S., Bates, C., Morton, C.E., et al., 2020 Aug 20. Factors associated with COVID-19-related death using OpenSAFELY [cited 2020 Dec 22] *Nature* 584 (7821), 430–436. <https://pubmed.ncbi.nlm.nih.gov/32640463/>.