

PRE-PUBLICACIONES del seminario matemático


2004

“garcía de galdeano”

Supercyclic vectors and
the angle criterion

Eva A. Gallardo-Gutiérrez
Jonathan R. Partington

n. 15



seminario
matemático

garcía de galdeano

Universidad de Zaragoza

SUPERCYCLIC VECTORS AND THE ANGLE CRITERION

EVA A. GALLARDO-GUTIÉRREZ AND JONATHAN R. PARTINGTON

ABSTRACT. It is shown that the *Angle Criterion* for testing supercyclic vectors depends in an essential way on the geometrical properties of the underlying space. In particular, we exhibit non-supercyclic vectors for the backward shift acting on c_0 that still satisfy such a criterion. Nevertheless, if \mathcal{B} is a locally uniformly convex Banach space, the Angle Criterion yields an equivalent condition for a vector to be supercyclic. Furthermore, it is proved that our hypotheses on \mathcal{B} cannot be weakened to the case of a strictly convex norm on \mathcal{B} .

1. INTRODUCTION

Let \mathcal{B} be a complex Banach space. A bounded linear operator T acting on \mathcal{B} is said to be cyclic if there is a vector $x \in \mathcal{B}$ such that the linear span generated by the orbit $\{T^n x\}_{n \geq 0}$ is dense in \mathcal{B} . In such a case, the vector x is called cyclic. If the orbit itself is dense, then T is called hypercyclic and x a hypercyclic vector for T . While cyclicity is a classical concept, hypercyclic operators have engaged the attention of many operator theorists since the end of the eighties. For a source of references on the subject, updated till 1999, we refer to Grosse-Erdmann's survey [5].

When the scalar multiples of the elements in the orbit are considered, the concept of supercyclic operator arises. We say that a bounded linear operator T acting on \mathcal{B} is supercyclic if there is $x \in \mathcal{B}$, also called supercyclic, such that the *projective orbit*

$$\{\lambda T^n x : n \geq 0 \text{ and } \lambda \in \mathbb{C}\}$$

is dense in \mathcal{B} . The concept of supercyclic operator was first introduced by Hilden and Wallen [6] in the seventies. Nevertheless, the first example of a supercyclic operator in the Banach space setting can be traced back to Rolewicz's work [10]. He proved that the unilateral backward shift B acting on the sequence space c_0 or

Date: March 2004.

2000 Mathematics Subject Classification. Primary 47A16, 47B33.

Key words and phrases. Supercyclic operators, supercyclic vectors, angle criterion.

First author is partially supported by Plan Nacional I+D grant no. BFM2003-00034 and Junta de Andalucía FQM-260.

ℓ^p , $1 \leq p < \infty$, and defined on the canonical basis $\{e_n\}_{n \geq 0}$ by

$$\begin{aligned} B e_n &= e_{n-1} & (n \geq 1) \\ B e_0 &= 0 \end{aligned}$$

is supercyclic. Actually, Rolewicz provided concrete examples of supercyclic vectors for B on c_0 by constructing hypercyclic vectors for complex scalar multiples λB whenever $|\lambda| > 1$. For a comprehensive treatment of properties of supercyclic operators, we refer the reader to Montes and Salas' recent survey [9].

Testing whether a vector $x \in \mathcal{B}$ possesses any kind of cyclic behaviour for a given operator T seems to be a difficult task. Of course, it is clear that $T^n x \neq 0$ for all $n \geq 0$. If, in addition, there exists a non-zero functional $\phi \in \mathcal{B}^*$ such that the set of complex numbers $\{\phi(T^n x)\}_{n \geq 0}$ is not dense in \mathbb{C} , then x is not hypercyclic for T . This fact was used in [4] to prove that certain operators are not hypercyclic (see *The Inner Product Criterion* [4, Chapter 2]). The underlying idea is simple and follows from the fact that the image of a dense set under a non-zero linear functional must be dense. Actually, this idea had been previously used by other authors. For instance, in some applications it can be enough to prove that $\{\phi(T^n x)\}_{n \geq 0}$ is bounded, or is contained in a straight line or is the set of powers of a fixed complex number. This latter argument was used in [2, Prop. 4.1] to prove a result of Kitai [7, Cor. 2.4] that asserts that the adjoint of a hypercyclic operator has always empty point spectrum.

On the other hand, in order to check that a given non-zero vector x is not supercyclic for T on a Hilbert space \mathcal{H} , it would be enough to find a vector y such that the whole orbit $\{T^n x\}_{n \geq 0}$ lies outside a cone around y . It is clear that the scalar multiples of the orbit of x cannot approximate y and, consequently, x cannot be supercyclic. This geometric idea was first used in [8, Sec. 5] to construct a non-supercyclic vector for certain weighted shifts on ℓ^2 . Moreover, this geometric idea works in any Hilbert space for any operator and provides a Hilbert space criterion for supercyclic vectors, which was given in [3].

The Angle Criterion for Hilbert spaces. *Let T be a bounded linear operator on a separable Hilbert space \mathcal{H} . Then the vector $x \in \mathcal{H}$ is supercyclic for T if and only if*

$$\sup_n \frac{|\langle T^n x, y \rangle|}{\|T^n x\| \|y\|} = 1$$

for all non-zero vectors $y \in \mathcal{H}$.

If we restrict ourselves to the Banach space setting, it follows that a necessary condition for a vector $x \in \mathcal{B}$ to be supercyclic for T is that

$$\sup_n \frac{|\phi(T^n x)|}{\|T^n x\| \|\phi\|} = 1$$

for all non-zero functionals $\phi \in \mathcal{B}^*$. However, as we shall see by means of a counterexample in Section 2, this condition is not sufficient for testing supercyclic vectors in general Banach spaces. Actually, it will depend, essentially, on the geometrical properties of the Banach space.

To this purpose, we recall that a Banach space \mathcal{B} is said to be *strictly convex* if the conditions $\|x\| = \|y\| = 1$ and $\|x + y\| = 2$ for vectors $x, y \in \mathcal{B}$ imply that $x = y$. The space \mathcal{B} is *locally uniformly convex* if for each $x \in \mathcal{B}$ and sequence (y_n) in \mathcal{B} , with $\|x\| = 1$ and $\|y_n\| = 1$ for each n , the condition $\|x + y_n\| \rightarrow 2$ implies that $\|y_n - x\| \rightarrow 0$. Clearly local uniform convexity implies strict convexity.

The relevance of these geometric properties to supercyclicity is that, as we shall show in Section 3, the angle criterion is necessary and sufficient for supercyclicity if the underlying space is locally uniformly convex, but it is insufficient assuming the slightly weaker property of strict convexity.

These convexity conditions are described in many places, for example [1], where it is noted that every separable Banach space can be given an equivalent locally uniformly convex norm. Moreover, the L^p spaces with $1 < p < \infty$ even have uniformly convex norms, and these are certainly locally uniformly convex. Thus, in principle, the angle criterion can be used to test supercyclicity by, if necessary, first passing to an equivalent locally uniformly convex norm.

2. A COUNTEREXAMPLE IN THE BANACH SPACE SETTING

In this section, we show that the Angle Criterion is no longer sufficient for testing supercyclic vectors in the Banach space setting.

First, we state the necessary condition for a vector x to be supercyclic in a Banach space \mathcal{B} , which is the analogue to that one stated for Hilbert spaces.

The Angle Criterion for Banach spaces. *Let T be a bounded linear operator on a separable Banach space \mathcal{B} . If the vector $x \in \mathcal{B}$ is supercyclic for T then*

$$(1) \quad \sup_n \frac{|\phi(T^n x)|}{\|T^n x\| \|\phi\|} = 1$$

for all non-zero functionals $\phi \in \mathcal{B}^*$.

We include the proof for the sake of completeness.

Proof: Let $\epsilon > 0$ be a positive number. There is a vector $z \in X$ such that $\|z\| = 1$ and $|\phi(z)| \geq (1 - \epsilon)\|\phi\|$. Since x is a supercyclic vector, there is a sequence (n_k) and complex numbers (λ_k) such that $\|\lambda_k T^{n_k} x - z\| \rightarrow 0$. But then

$$\limsup_{k \rightarrow \infty} \frac{|\phi(T^{n_k} x)|}{\|T^{n_k} x\| \|\phi\|} \geq (1 - \epsilon),$$

implying (since ϵ was arbitrary) that condition (1) holds. □

Recall that c_0 denotes the Banach space of all sequences $(x(n))_{n \geq 0}$ converging to zero endowed with the supremum norm $\|(x(n))_{n \geq 0}\|_\infty = \max_n |x(n)|$. As mentioned in the introduction, Rolewicz [10] showed that if $|\lambda| > 1$ then λB is hypercyclic on c_0 and ℓ^p for $1 \leq p < \infty$. We have the following

Theorem 2.1. *Let λ be a real number with $\lambda > 1$ and let $T = \lambda B$, regarded as an operator on c_0 . Then T possesses non-supercyclic vectors satisfying that*

$$\sup_n \frac{|\phi(T^n x)|}{\|T^n x\| \|\phi\|} = 1$$

for all non-zero functionals $\phi \in \ell^1$.

Proof: We shall construct a vector x for which $\alpha T^n x$ is always far from the unit basis vector e_0 whenever any scalar $\alpha \in \mathbb{C}$ is considered. To do this we fix a real constant $R > \lambda$, and construct an element x of c_0 which, apart from its other properties, satisfies

$$(2) \quad \left| \frac{x(n+1)}{x(n)} \right| \geq \frac{1}{R} \quad \text{for each } n.$$

If condition (2) holds, then

$$\|\alpha T^n x - e_0\|_\infty \geq \max\{|\alpha \lambda^n x(n) - 1|, |\alpha \lambda^n x(n+1)|\},$$

but if the first term is less than $\frac{1}{2}$, then the second term is at least $\frac{1}{2R}$. Hence such a vector x will not be supercyclic.

To proceed with the construction of x , let us fix (ϕ_k) a dense sequence in the unit sphere of $c_0^* = \ell^1$ consisting of vectors of finite support of length L_k . Let $(v_k) \subset c_0$ be a sequence of vectors also of finite support of length L_k such that $|v_k(j)| = 1$ for $0 \leq j < L_k$ and $\phi_k(v_k) = \|\phi_k\|_1 = \|v_k\|_\infty = 1$. Choosing M_k sufficiently large such that $R^{L_k - M_k} \leq \lambda^{-M_k}$, we define the norm-one vectors u_k as follows:

$$u_k(j) = \begin{cases} v_k(j) & \text{if } 0 \leq j < L_k, \\ R^{L_k - j} & \text{if } L_k \leq j < M_k, \end{cases}$$

Observe that the number of the non-zero coordinates of u_k is actually M_k .

Let (N_k) be the sequence of positive integers given by $N_0 = 0$ and $N_{k+1} = N_k + M_k$. If S denotes the forward shift on c_0 , which satisfies $BS = \text{Id}$, the vector $x \in c_0$ we look for is defined by

$$(3) \quad x = \sum_{k=0}^{\infty} \frac{1}{\lambda^{N_k}} S^{N_k} u_k.$$

Observe that, x satisfies the non-supercyclicity condition (2) since $R^{L_k - M_k} \leq \lambda^{-M_k}$. In addition, by construction, $\|T^{N_k} x\|_\infty = 1$ as well as $|\phi_k(T^{N_k} x)| = 1$. Therefore,

$$\sup_n \frac{|\phi_k(T^n x)|}{\|T^n x\| \|\phi_k\|} = 1$$

for each k . Clearly, the statement of the theorem follows since (ϕ_k) is dense in the unit sphere of c_0^* and this latter fact suffices. \square

3. LOCALLY UNIFORMLY CONVEX BANACH SPACES

The following result shows that if the underlying Banach space \mathcal{B} is a locally uniformly convex space, then the Angle Criterion in \mathcal{B} is also a sufficient condition for supercyclicity.

Theorem 3.1. *Let \mathcal{B} be a locally uniformly convex Banach space and T a bounded linear operator on \mathcal{B} . Then the vector x is supercyclic for T if and only if*

$$\sup_n \frac{|\phi(T^n x)|}{\|T^n x\| \|\phi\|} = 1$$

for all non-zero functionals $\phi \in \mathcal{B}^*$.

Proof: It is sufficient to show that the angle criterion implies supercyclicity. Let $y \in \mathcal{B}$ a non-zero vector. We may suppose, without loss of generality, that $\|y\| = 1$. By the Hahn–Banach theorem, it is possible to find $\phi \in \mathcal{B}^*$ with $\|\phi\| = \phi(y) = 1$.

On the other hand, the angle criterion implies that there is a sequence (n_k) such that $\phi(y_k) \rightarrow 1$, where $y_k = \epsilon_k T^{n_k} x / \|T^{n_k} x\|$ and (ϵ_k) are complex constants of modulus 1. Note that $\|y_k\| = 1$ for all k and so $\|y + y_k\| \leq 2$.

Since $|\phi(y + y_k)| \rightarrow 2$ we conclude that $\|y + y_k\| \rightarrow 2$ and by the local uniform convexity we deduce that $\|y_k - y\| \rightarrow 0$. Since this holds for all $y \in \mathcal{B}$, the operator T is supercyclic. \square

Now let \mathcal{B} denote c_0 equipped with the equivalent strictly convex norm

$$\|x\| = \left(\|x\|_\infty^2 + \sum_{n=0}^{\infty} \frac{|x(n)|^2}{(n+1)^2} \right)^{1/2}.$$

Thus,

$$(4) \quad \|x\|_\infty \leq \|x\| \leq \left(1 + \frac{\pi^2}{6} \right)^{1/2} \|x\|_\infty \quad \text{for all } x \in \mathcal{B}.$$

We shall use the space \mathcal{B} to show that the Angle Criterion is not sufficient for supercyclicity even in a strictly convex Banach space.

Theorem 3.2. *Let λ be a real number with $\lambda > 1$ and let $T = \lambda B$, regarded as an operator on the strictly convex space \mathcal{B} . Then T possesses non-supercyclic vectors satisfying that*

$$\sup_n \frac{|\phi(T^n x)|}{\|T^n x\| \|\phi\|} = 1$$

for all non-zero functionals $\phi \in \mathcal{B}^*$.

Proof: We modify the proof of Theorem 2.1 as follows. Having fixed $R > \lambda$ we can no longer guarantee condition (2), but instead we shall find a vector $x \in \mathcal{B}$ such that for each n there is an $n' > n$ with

$$(5) \quad \left| \frac{x(n')}{x(n)} \right| \geq \frac{1}{R\beta},$$

where β denotes the constant $\left(1 + \frac{\pi^2}{6}\right)^{1/2}$ occurring in (4). This guarantees that $\|\alpha T^n x - e_0\| \geq \frac{1}{2R\beta}$ for any scalar $\alpha \in \mathbb{C}$. Therefore, the proof runs similarly to that of Theorem 2.1.

Again we take functionals (ϕ_k) dense in the unit sphere of \mathcal{B}^* , and we may take them to have finite support of length L_k . Let (v_k) be also vectors of finite support of length L_k such that $\phi_k(v_k) = \|\phi_k\| = \|v_k\|_\infty = 1$. We can longer take the v_k to have coordinates of modulus 1.

We define the vectors u_k by

$$u_k(j) = \begin{cases} v_k(j) & \text{if } 0 \leq j < L_k, \\ 0 & \text{if } L_k \leq j < P_k, \\ \beta^{-1} R^{P_k-j} & \text{if } P_k \leq j < M_k, \\ 0 & \text{if } j \geq M_k. \end{cases}$$

That is, $u_k = v_k + w_k$, where

$$w_k(j) = \begin{cases} 0 & \text{if } 0 \leq j < P_k, \\ \beta^{-1} R^{P_k-j} & \text{if } P_k \leq j < M_k, \\ 0 & \text{if } j \geq M_k. \end{cases}$$

Here P_k is chosen sufficiently large such that $\|u_k\| \leq 1 + 1/k$. Observe, we can do this because, first,

$$\|v_k\|_\infty \geq \beta^{-1}$$

and so, $\|v_k + w_k\|_\infty = \|v_k\|_\infty$. Secondly, we can make the series

$$\sum_{n=P_k}^{\infty} \frac{|w_k(n)|^2}{(n+1)^2}$$

arbitrarily small by choosing P_k large. Let M_k be chosen sufficiently large such that $R^{P_k-M_k} \leq \beta\lambda^{-M_k}$. As in the proof of Theorem 2.1, we consider (N_k) the sequence defined by $N_0 = 0$ and $N_{k+1} = N_k + M_k$, for each $k \geq 0$. We require that the vector x , defined by

$$x = \sum_{k=0}^{\infty} \frac{1}{\lambda^{N_k}} S^{N_k} u_k,$$

is not supercyclic for T .

To check the non-supercyclicity of x , observe that since $R^{P_k-M_k} \leq \beta\lambda^{-M_k}$, condition (5) holds for $n = N_{k+1} - 1$ and n' the index of the numerically largest coordinate of $\lambda^{-N_{k+1}} S^{N_{k+1}} u_{k+1}$, which is at least $\beta^{-1}\lambda^{-N_{k+1}}$ in absolute value (the

observation that (5) holds for other values of n is immediate from the construction of each u_k). Therefore, x is not a supercyclic vector. However,

$$\sup_n \frac{|\phi_k(T^n x)|}{\|T^n x\| \|\phi_k\|} = 1$$

since $\phi_k(T^{N_k} x) = 1$ and $\|T^{N_k} x\| \rightarrow 1$ as $k \rightarrow \infty$. Now, the density of (ϕ_k) in the unit sphere of \mathcal{B}^* yields the statement of the theorem. \square

Acknowledgments

This work was initiated while the first author was visiting University of Leeds. She is grateful to the Mathematical Department at University of Leeds for its hospitality. The authors are also grateful to the London Mathematical Society for financial support.

REFERENCES

- [1] Y. Benyamini and J. Lindenstrauss, *Geometric nonlinear functional analysis*, Vol. 1, A.M.S. Colloquium Publications, 48, American Mathematical Society, Providence, RI, 2000.
- [2] K. C. Chan and J. H. Shapiro, *The cyclic behavior of translation operators on Hilbert spaces of entire functions*, Indiana Univ. Math. J. **40** (1991), 1421–1449.
- [3] E. A. Gallardo-Gutiérrez and A. Montes-Rodríguez, *The role of the angle in the supercyclic behavior*, J. Functional Analysis, **203** (2003), 27–43.
- [4] E. A. Gallardo-Gutiérrez and A. Montes-Rodríguez, *The role of the spectrum in the cyclic behavior of composition operators*, Memoirs of A. M. S., **167**, (2004) no. 791.
- [5] K. G. Grosse-Erdmann, *Universal families and hypercyclic operators*, Bull. Amer. Math. Soc (NS) **36** (1999), 345–281.
- [6] H. M. Hilden and L. J. Wallen, *Some cyclic and non-cyclic vectors of certain operators*, Indiana Univ. Math. J. **23** (1974), 557–565.
- [7] C. Kitai, *Invariant Closed Sets for Linear Operators*, Thesis, Univ. Toronto, (1982).
- [8] A. Montes-Rodríguez and H. N. Salas, *Supercyclic subspaces: spectral theory and weighted shifts*, Adv. Math. **163**, (2001), 74–134.
- [9] A. Montes-Rodríguez and H. N. Salas, *Supercyclic subspaces*, Bull. London Math. Soc. **35** (2003), 721–737.
- [10] S. Rolewicz *On orbits of elements*, Studia Math. **32** (1969), 17–22.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE ZARAGOZA, PLAZA SAN FRANCISCO S/N,
50009 ZARAGOZA, SPAIN.

E-mail address: `eva@unizar.es`

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, U.K.

E-mail address: `J.R.Partington@leeds.ac.uk`