

1 **TITLE:**

2 Validation of coffee by-products as novel food ingredients

3

4 **AUTHORS:**

5 Amaia Iriondo-DeHond <sup>a</sup>, Natalia Aparicio García <sup>a</sup>, Beatriz Fernandez-Gomez <sup>a</sup>,

6 Eduardo Guisantes-Batan <sup>b</sup>, Francisco Velázquez Escobar <sup>c</sup>, Gracia Patricia Blanch <sup>d</sup>,

7 Manuel Ignacio San Andres <sup>e</sup>, Sebastián Sanchez-Fortun <sup>e</sup>, María Dolores del Castillo <sup>a\*</sup>

8

9 **AFFILIATIONS:**

10 <sup>a</sup> Instituto de Investigación en Ciencias de la Alimentación (CIAL) (CSIC-UAM), C/

11 Nicolás Cabrera, 9, Campus de la Universidad Autónoma de Madrid, 28049 Madrid,

12 Spain; [amaia.iriondo@csic.es](mailto:amaia.iriondo@csic.es), [n.aparicio@ictan.csic.es](mailto:n.aparicio@ictan.csic.es), [b.fernandez@csic.es](mailto:b.fernandez@csic.es),

13 [mdolores.delcastillo@csic.es](mailto:mdolores.delcastillo@csic.es)

14 <sup>b</sup> Universidad de Castilla-La Mancha, Instituto Regional de Investigación Científica

15 Aplicada (IRICA), Av. Camilo José Cela s/n, 13071 Ciudad Real, Spain;

16 [eduardo.guisantes@uclm.es](mailto:eduardo.guisantes@uclm.es)

17 <sup>c</sup> Technische Universität Berlin, 135/PC14 Max Volmer Laboratorium für

18 biophysikalische Chemie, 1023 Berlin, Germany; [velazquez.escobar@chem.tu-berlin.de](mailto:velazquez.escobar@chem.tu-berlin.de)

19 <sup>d</sup> Instituto de Ciencia y Tecnología de Alimentos y Nutrición (ICTAN) (CSIC), C/ José

20 Antonio Novais, 10, 28040 Madrid, Spain; [gblanch@ictan.csic.es](mailto:gblanch@ictan.csic.es)

21 <sup>e</sup> Facultad de Veterinaria, Universidad Complutense de Madrid, Av. Puerta de Hierro, s/n,

22 28040 Madrid, Spain; [fortun@ucm.es](mailto:fortun@ucm.es), [misanand@ucm.es](mailto:misanand@ucm.es)

23

24 \* **Corresponding author.** Dr. M. Dolores del Castillo, Tel: +34 91 0017900 Ext. 953,

25 Fax: +34 91 0017905, e-mail: [mdolores.delcastillo@csic.es](mailto:mdolores.delcastillo@csic.es)

26

27 **Abbreviations**

28 ABTS, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt; ATR,  
29 attenuated total reflection; CGA, chlorogenic acid; 5-CQA, 5-caffeoylquinic acid; CS,  
30 coffee silverskin; DCFH-DA, 2',7'-dichlorodihydrofluorescein diacetate; DDD,  
31 dichlorodiphenyldichloroethane; DDE, dichlorodiphenyldichloroethylene; DDT,  
32 dichlorodiphenyltrichloroethane; DMEM, Dulbecco's Modified Eagle's Medium;  
33 DMSO, dimethyl sulfoxide; EFSA, European Food Safety Authority; FOS,  
34 fructooligosaccharides; FT, Fourier-Transform; HCH, hexachlorocyclohexane; IDF,  
35 insoluble dietary fiber; IR, infrared; MTT, 3-(4,5-dimethylthiazole-2,5-  
36 diphenyltetrazolium bromide; OECD, Organization for Economic Co-operation and  
37 Development; OTA, ochratoxin A; ROS, reactive oxygen species; RUT, rutin; SDF,  
38 soluble dietary fiber; TAC, total antioxidant capacity; tBOOH, tert-butyl hydroperoxide;  
39 TDF, total dietary fiber; TFC, total flavonoid content; TPC, total phenolic compounds.

40

41 **Abstract**

42 This research aimed to validate coffee husk, parchment and silverskin as new health-  
43 promoting food ingredients. Characterization of the novel ingredients was carried out by  
44 Raman and infrared spectroscopy and analysis of total phenolic compounds, chlorogenic  
45 acid, caffeine and dietary fiber. Antioxidant properties of the novel ingredients were  
46 tested by ABTS and intracellular ROS formation in HepG2 cells. Pesticides, mycotoxins,  
47 acrylamide and acute toxicity experiments following OECD Test Guidelines 425 were  
48 performed to assess the food safety of extracts, solid residues and raw materials. Husk  
49 and silverskin are proposed as a source of two food ingredients: aqueous extracts enriched  
50 in phytochemicals and antioxidant dietary fiber while parchment is proposed as a natural  
51 source of antioxidant dietary fiber. No lesions were found in selected isolated vital organs  
52 from treated animals. Coffee by-products can be converted into safe food ingredients  
53 allowing a whole food waste recovery. Analyses of contaminants is essential for  
54 achieving this goal.

55

56

57 **Keywords:** Antioxidants, coffee by-products, dietary fiber, food safety, novel food  
58 ingredient, sustainability.

59

## 60        **1. Introduction**

61    Coffee is a cherry formed by various anatomic parts with distinct morphologies (Figure  
62    1) and unique chemical compositions. It comprises a plethora of nutrients and bioactive  
63    phytochemicals with diverse preventive and therapeutic effects for chronic diseases (del  
64    Castillo et al., 2017). Nowadays, coffee beans are almost exclusively used for the  
65    preparation of the beverage. While the coffee industry is very diverse, coffee processing  
66    can generally be described by ten steps: planting, harvesting and processing the cherries,  
67    drying, milling and trading, tasting, roasting, grinding and finally brewing. Even though  
68    most of the health-promoting phytochemicals and nutrients are found in the coffee cherry,  
69    90% of the cherry is discarded during processing as agricultural waste or by-product (del  
70    Castillo, Fernandez-Gomez, Martinez-Saez, Iriondo-DeHond, & Mesa, 2018).

71    According to the European Commission, a “food waste” is defined as “food (including  
72    inedible parts) lost from the food supply chain, not including food diverted to material  
73    uses such as bio-based products, animal feed, or sent for redistribution” (European  
74    Commission, 2014). The coffee industry is responsible for the generation of large  
75    amounts of residues that represent a great pollution hazard if discharged into the  
76    environment (Chanakya & De Alwis, 2004). The future of coffee depends on sustainable  
77    practices, and a wide range of initiatives can be implemented in every part of the coffee  
78    supply chain: improving the conditions at origin, recycling packaging materials, reducing  
79    emissions, developing eco-friendly facilities or designing new coffee products. The  
80    conversion of coffee waste into food ingredients is a research priority in the Nutrition and  
81    Food Chemistry field.

82    Esquivel et al. (2012) defined coffee husk as the outer skin and pulp obtained from the  
83    wet processing of coffee berries (Esquivel & Jiménez, 2012). Coffee husk encloses the  
84    coffee beans and compromises nearly 45% of the berry. It has a high content in

85 carbohydrates (35-85%), soluble fibers (30.8 %), minerals (3-11%) and proteins (5-11%).  
86 It is also rich in insoluble dietary fiber and can be a source of phytochemicals such as  
87 tannins (5-9%) and cyanidins (20%) for the food and pharmaceutical industries (M.  
88 Dolores del Castillo et al., 2018; Esquivel & Jiménez, 2012). Phytochemicals have  
89 recently received considerable interest because of their safety and potential positive  
90 physiological effects on the human body (Singh & Geetanjali, 2013). To date, coffee husk  
91 has been used as a source of dietary fiber ([www.pectcof.com](http://www.pectcof.com)) and also as a source of  
92 anthocyanins (Murthy, Manjunatha, Sulochannama, & Madhava Naidu, 2012).  
93 Considering its chemical composition and the proposed applications, coffee husk has a  
94 great potential as a food ingredient and as a natural source of nutrients and bioactive  
95 compounds.

96 Parchment is a strong fibrous endocarp that covers both hemispheres of the coffee seed  
97 and separates them from each other (Belitz, Grosch, & Schieberle, 2009). It represents  
98 5.8% of berry dry weight and is formed by ( $\alpha$ -) cellulose (40–49%), hemicellulose (25–  
99 32%), lignin (33–35%) and ash (0.5–1%) (del Castillo et al., 2018). In wet coffee  
100 processing, the parchment is removed after drying and hulling, which allows it to be  
101 collected and used separately from other by-products (Esquivel & Jiménez, 2012). To  
102 date, no applications have been found for coffee parchment as a food ingredient.

103 Coffee silverskin (CS) represents 4.2 % (w/w) of the coffee cherry and constitutes a thin  
104 tegument of the outer layer of the two beans forming the green coffee seed. It is obtained  
105 as a by-product of the roasting process. CS has a high dietary fiber content (68-80%),  
106 polysaccharides are also abundant components (60-70 %) and total sugar content varies  
107 greatly (1.6–12%) (Pourfarzad, Mahdavian-Mehr, & Sedaghat, 2013). CS contains  
108 protein, fat, and ash, at 16.2–19.0%, 1.56–3.28%, and 5-7%, respectively (del Castillo et  
109 al., 2018). Borrelli and colleagues found that CS, obtained from several Italian roasting

110 plants, had a high amount of soluble dietary fiber (about 14% of the total fiber) and a very  
111 high antioxidant activity (Borrelli, Visconti, Mennella, Anese, & Fogliano, 2002).  
112 Previous studies have suggested the use of CS as a source of dietary fiber and prebiotics  
113 (Ballesteros, Teixeira, & Mussatto, 2014; Borrelli, Esposito, Napolitano, Ritieni, &  
114 Fogliano, 2004).

115 In 2016, the European Food Safety Authority (EFSA) reported that only 1.5% of 83,000  
116 food samples analyzed exceeded the legal limit of pesticides, and the national competent  
117 authorities had to take appropriate enforcement actions (European Food Safety Authority  
118 (EFSA), 2016). To date, no studies have analyzed pesticide content in coffee by-products.  
119 Moreover, coffee is susceptible to contamination by mycotoxins, toxic compounds that  
120 result from fungal secondary metabolism under certain conditions, which cause different  
121 toxicological effects in humans (Afsah-Hejri, Jinap, Hajeb, Radu, & Shakibazadeh, 2013;  
122 Gamboa-Gaitán, 2012). The most studied mycotoxin in coffee is ochratoxin A (OTA),  
123 the only mycotoxin subjected to European legislation (European Commission (EC)  
124 1881/2006). Other contaminants may also be formed during food processing, e.g.  
125 Maillard reaction products such as acrylamide. This low molecular weight compound is  
126 highly soluble in water, and has a well-documented carcinogenic potential (European  
127 Commission, 2017). It is formed in food processing as a degradation product of  
128 asparagine and sugars, typically at 120 °C and low moisture (European Commission,  
129 2017). Therefore, it could be found in roasted coffee beans and CS. Previous studies have  
130 confirmed the presence of acrylamide in CS (333 µg/kg) (Garcia-Serna, Martinez-Saez,  
131 Mesias, Morales, & Castillo, 2014) in the same amount as that reported in roasted coffee  
132 and approximately 10 times lower than that reported in instant coffee (European  
133 Commission, 2017).

134 The aim of this research was to perform a comparative study of the three coffee by-  
135 products obtained before (husk and parchment) and after the roasting process (silverskin);  
136 validate their potential and safety as sustainable food ingredients; explore novel  
137 applications for them through environment-friendly procedures and implement novel  
138 analytical tools such as vibrational spectroscopy.

## 139 **2. Materials and methods**

### 140 *2.1. Reagents*

141 Chlorogenic acid, 2,21-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) diammonium  
142 salt (ABTS), caffeine, Folin-Ciocalteu reagent, tert-butyl hydroperoxide (tBOOH),  
143 dimethyl sulfoxide (DMSO), 3-(4,5-dimethylthiazole-y)-2,5-diphenyltetrazolium  
144 bromide (MTT) and 21,71-dichlorodihydro-fluorescein diacetate (DCFH-DA) were  
145 purchased from Sigma Chemical (Sigma-Aldrich, St Louis, MO, USA). Dulbecco's  
146 Modified Eagle's Medium (DMEM) was purchased from Lonza (Basel, Switzerland).

### 147 *2.2. Raw materials*

148 Coffee husk and parchment from Arabica species were kindly provided by Delikia from  
149 Finca Morenitas in Nicaragua. CS from Arabica and Robusta species from Colombia was  
150 provided by Fortaleza S.A. (Spain). All coffee by-products were derived from wet-  
151 processing.

### 152 *2.3. Raman and infrared spectroscopy*

153 Fourier-Transform (FT) Raman spectra were recorded with an RFS-100/S spectrometer  
154 (Bruker, Billerica, USA). A single 4x4 mm<sup>2</sup> endocarp shell was positioned between a thin  
155 quartz window (1 mm) and a silver-mirror. Spectra were recorded according to previous

156 measuring protocols for green coffee (Keidel, von Stetten, Rodrigues, Máguas, &  
157 Hildebrandt, 2010).

158 Complementary endocarp FT-infrared (IR) spectra were recorded in a Tensor27 FT-  
159 spectrometer (Bruker, Billerica, USA), equipped with a diamond attenuated total  
160 reflection (ATR) cell (Durascope Resultec). The single endocarp shell was positioned and  
161 pressed towards the ATR-diamond surface. Absorbance spectrum was calculated from  
162 the recorded sample spectrum and a previously measured reference spectrum (empty  
163 ATR-cell). Further details regarding spectrometers and data processing are given  
164 elsewhere (Ulijasz et al., 2009).

165 For the band assignment of the endocarp IR and the Raman spectra, we refer to previous  
166 experimental works in cellulose, lignin and further related compounds (Adebajo, Frost,  
167 Klopogge, & Kokot, 2006; Gierlinger, Keplinger, Harrington, & Schwanninger, 2013;  
168 Nabais et al., 2008).

#### 169 *2.4. Characterization of coffee by-product extracts*

##### 170 *2.4.1. Preparation of aqueous extracts from coffee by-products*

171 Coffee by-product extracts were produced as described in the patent WO 2013/004873  
172 (del Castillo et al., 2013). Briefly, 50 mg of coffee husk, parchment or silverskin were  
173 added per milliliter of boiling water. This mixture was stirred at 250 rpm for 10 min;  
174 filtered by a 250 µm filter and then through Whatman paper No. 4; and the filtrate and  
175 the insoluble residue were freeze-dried. Extracts were prepared three times.

##### 176 *2.4.2. Caffeine and 5-CQA*

177 Caffeine and 5-CQA content was determined by ultra-performance liquid  
178 chromatography - tandem mass spectrometer (UPLC-MS/MS) (Thermo-Scientific, San

179 José, CA, USA) as described by Fernandez-Gomez, Lezama, et al., (2016). For  
180 quantification, samples were diluted in milli-Q water and added with 50 µg/ml of  
181 phluoroglucinol as internal standard for the most abundant chlorogenic acid (CGA)  
182 described for coffee (5-caffeoylquinic acid, 5-CQA) and 50 µg/ml of salicylic acid as  
183 internal standard for caffeine. To verify the reproducibility and possible variations, all  
184 experiments were performed in triplicate. Compound concentrations were expressed in  
185 mg caffeine or 5-CQA/ g of sample.

#### 186 *2.4.3. Overall antioxidant capacity*

187 The trapping capacity of cationic free radicals was evaluated using the method of radical  
188 ABTS<sup>+</sup> bleaching described by Re et al., (1999) and modified by Oki, Nagai, Yoshinaga,  
189 Nishiba, & Suda (2006) for its use in microplate. Aqueous solutions of CGA (0.15–2.0  
190 mmol/l) were used for calibration. Absorbance was measured in microplate using a UV-  
191 Visible Spectrophotometer (BioTek Instruments, Winooski, VT, USA). All  
192 measurements were performed in triplicate, and results were expressed as mg CGA eq.  
193 per gram of sample.

#### 194 *2.4.4. Total phenolic compounds*

195 Folin–Ciocalteu adapted to a micromethod format was used for the analysis of total  
196 phenolic compounds in samples (Contini, Baccelloni, Massantini, & Anelli, 2008). The  
197 reaction was initiated by mixing 10 µl of sample with 150 µl of Folin–Ciocalteu solution.  
198 After 3 minutes at room temperature, 50 µl of sodium bicarbonate solution were added.  
199 The kinetics of the reaction at 37 °C was followed for 120 minutes by measuring the  
200 absorbance at 735 nm once every minute using a UV-Visible Spectrophotometer (BioTek  
201 Instruments, Winooski, VT, USA). Sample blank and reagent blank were also analyzed  
202 in each set of samples. A CGA calibration curve was used for quantification (0.1–0.8

203 mg/ml). Results were expressed as  $\mu\text{mol CGA eq./g}$ . All measurements were performed  
204 in triplicate.

#### 205 *2.4.5. Total flavonoid content*

206 Total flavonoid content was determined using a previously described colorimetric method  
207 (Xu & Chang, 2007). Briefly, 100  $\mu\text{L}$  of samples or rutin (RUT) standard solution were  
208 mixed with 30  $\mu\text{L}$  of a 5 %  $\text{NaNO}_2$  solution. After 5 minutes of incubation, 30  $\mu\text{L}$  of a  
209 10%  $\text{AlCl}_3$  solution were added and allowed to stand for another 6 minutes before adding  
210 100  $\mu\text{L}$  of 2M  $\text{NaOH}$ . Absorbance was measured immediately at 510 nm using a UV-  
211 Visible Spectrophotometer (BioTek Instruments, Winooski, VT, USA). Results were  
212 calculated and expressed as milligrams of RUT equivalents per 100 mg of extract. The  
213 determination was carried out in triplicate.

#### 214 *2.4.6. Cell culture and treatments*

215 HepG2 human hepatocyte cell line was kindly provided by Dr. Paloma Morales (Facultad  
216 de Veterinaria, Universidad Complutense de Madrid, Spain). Cells were cultured in  
217 Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% fetal bovine  
218 serum (FBS), 1% L-glutamine and 1% penicillin/streptomycin in standard conditions (37  
219  $^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , in a humidified incubator, BINDER CB series 2010, Tuttlingen, Germany).

#### 220 *2.4.7. Cell viability assays*

221 The effect of different concentrations of husk, parchment, and CS extracts on cell viability  
222 was measured using the MTT assay (Bakondi et al., 2003). Powdered extracts were  
223 prepared in PBS and sterile filtered when used in cell culture. Cells were cultured at a  
224 density of  $1.0 \times 10^5$  cells per well of a 96-well plate. After 24 hours, cells were treated with  
225 husk, parchment, and CS extracts diluted in DMEM culture medium (0.1, 1, 10 and 20  
226 mg/ml) for 24 hours. DMSO (50%) was used as death control. Subsequently, cells were

227 incubated in MTT solution (0.5 mg/ml) for 1 h at 37 °C. The supernatant was then  
228 removed, 100 µL of DMSO were added, and the optical density at 570 nm was measured  
229 using a microplate reader (BioTek Synergy HT Multi-Mode Microplate Reader,  
230 Winooski, VT, USA). Experiments were carried out in triplicate.

#### 231 *2.4.8. Intracellular ROS scavenging assay*

232 The analysis was performed by measuring the fluorescence intensity of the DCFH-DA  
233 probe, which was proportional to the amount of ROS formed (Gomes, Fernandes, &  
234 Lima, 2005). A 10 mM solution of DCFH-DA was prepared (5 mg in 1 ml DMSO), and  
235 a 50 µL aliquot was separated. Then, 800 µL of DMSO were added to the 50 µL solution.  
236 After 24 hours of extract incubation, cells were pre-loaded with 2.5 µL/well of this last  
237 solution for 30 minutes at 37 °C. After incubation, DCFH becomes dichlorofluorescein  
238 (DCF) due to intracellular oxidants and will emit fluorescence. Next, the culture medium  
239 was removed; cells were washed with PBS; and the three coffee by-product extracts (1  
240 mg/ml) were added for 1 hour. tBOOH 1 mM was used as a positive control. Then,  
241 fluorescence was measured at 485 nm/528 nm (BioTek Synergy HT Multi-Mode  
242 Microplate Reader). Experiments were carried out in triplicate.

#### 243 *2.5. Characterization of the insoluble fraction obtained from coffee by-products during* 244 *the extraction process*

##### 245 *2.5.1. Dietary fiber*

246 Insoluble (IDF), soluble (SDF) and total (TDF) dietary fiber content was determined  
247 using the Total Dietary Fiber Assay Kit (Megazyme International Ireland, Ireland)  
248 following the manufacturer's instructions, and based on the enzymatic–gravimetric  
249 method. Results are expressed as percentage (%).

##### 250 *2.5.2. Total protein*

251 Protein content was determined by Kjeldahl mineralization followed by a colorimetric  
252 analysis of nitrogen for quantification (AOAC-32.1.22, 920.87). NH<sub>4</sub>Cl was used for a  
253 standard calibration curve. A conversion factor (6.25) was used to calculate protein  
254 content. Results were expressed as % dry matter (d.m.).

### 255 *2.5.3. Total lipid content*

256 Using a modified protocol described by Folch, Lees, & Sloane Stanley (1987), lipid  
257 content was determined by a solid-liquid extraction with n-hexane. Fifty milliliters of n-  
258 hexane were added to the insoluble fractions (1 g) and the mixture was homogenized  
259 using an Ultra-Turrax for 3 minutes and subjected to an ultrasonic bath for 20 minutes to  
260 improve lipid extraction. Then, samples were centrifuged at 1620 g for 20 minutes.  
261 Supernatant's solvent was then gently removed under controlled vacuum. Total lipid  
262 content was obtained by weighing difference. The assay was performed in duplicate and  
263 the results expressed as % lipid content.

### 264 *2.5.4. Overall antioxidant capacity*

265 Direct ABTS<sup>+</sup> assay was carried out according to Açar, Gökmen, Pellegrini, & Fogliano  
266 (2009). Ten mg of sample were mixed with 90 mg of cellulose and stirred, 10 mg of the  
267 mixture were mixed together with 1.7 ml of ABTS<sup>+</sup> solution in a thermomixer (25 °C, 2  
268 minutes, 600 rpm). After centrifugation, absorbance of the supernatant was measured in  
269 a microplate. A CGA calibration curve (0–250 µg/ml) was used. Measurements were  
270 performed in triplicate and results expressed as mg CGA eq. per gram of sample.

## 271 *2.6. Safety assays*

### 272 *2.6.1. Pesticides*

273 The gas chromatography system (HPGC-7890A) was equipped with split/splitless  
274 injector and a MS detector (model MS-5975-VL-MSD with Triple –Axis Detector). The  
275 automatic split/splitless injector operated at 250°C and the split (1:10) was used to detect  
276 the analytes with helium as the carrier gas at an approximate flowrate of 1 ml/min. Gas  
277 chromatographic analysis of samples (husk extract and insoluble fraction, raw parchment  
278 and Arabica CS extract and insoluble fraction) was performed using a 30 m × 0.25 mm  
279 ID capillary fused silica column coated with a 0.25 µm film of TRB-5 (Teknokroma,  
280 Spain). Oven temperature-ramp followed a 12 °C/min increase-rate from 60°C (2 min) to  
281 140°C and 6°C/min up to 320°C. Desired temperature was maintained for 5 min.

282 The ion source temperature of the MS was 230°C, mass range 40–500 amu, scan speed  
283 1666 amu/s, interface temperature 230°C. First, the SCAN mode and then the selected ion  
284 monitoring (SIM) mode were used for the analyses. An advanced pressure control  
285 supplied helium to the interface at constant pressure (95 kPa). In all cases, analyses were  
286 made in triplicate. Pesticide identification was based on the retention time and the relative  
287 abundance of the specific ions (used in the SIM mode), compared from the sample extract  
288 (Supplementary Material, Table S1). The number of monitored ions, preferably and if  
289 possible a minimum of three, was dependent on the pesticide. Analyzed pesticides were  
290 HCH- $\alpha$ , HCH- $\beta$ , HCH- $\gamma$ , HCH- $\delta$ , heptachlor, Aldrin, hexachlorbenzene, endosulfan- $\alpha$ ,  
291 endosulfan- $\beta$ , chlorpyrifos methyl, parathion methyl, parathion ethyl, dieldin, endrin, 4,4-  
292 DDE, 2,4-DDD, 44DDT, 24DDE and mirex.

### 293 2.6.2. *Mycotoxins*

294 Samples (husk extract and insoluble fraction, raw parchment and Arabica CS extract and  
295 insoluble fraction) were ground with an A-10 Basic laboratory mill (IKA, Staufen,  
296 Germany) and stored at -20 °C until use. Ground samples (1 g) were mixed with  
297 acetonitrile:water (80:20, v/v, 2.5 ml) in a 15 ml centrifuge tube, vortexed (5 min),

298 sonicated (3 min), vortexed again (1 min) and then samples were centrifuged (3000 g, 10  
299 min). Subsequently, the supernatant was separated. The residue was extracted again  
300 following the same procedure and the supernatants were combined. One ml of the  
301 supernatant was transferred to a new 15 ml centrifuge tube and diluted with water (7 ml)  
302 to a total volume of 8 ml and centrifuged (3000 g, 10 min).

303 Mycotoxins were extracted using ISOLUTE® Myco 60 mg/3 ml (Biotage, Sweden) SPE  
304 columns following the procedure recommended by the supplier. Briefly, SPE column was  
305 conditioned with 2 ml acetonitrile, equilibrated with 2 ml of ammonium acetate (10 mM)  
306 and 3 ml of the diluted sample extract were loaded. Then, the column was washed with  
307 10 mM ammonium acetate (3 ml) and with 10 mM ammonium acetate:acetonitrile (90:10,  
308 v/v, 3 ml) and dried for 30 seconds at maximum vacuum. Finally, the sample was eluted  
309 with 0.1% formic acid in acetonitrile (2 ml) and then with 0.1% formic acid in methanol  
310 (2 ml). The combined eluates were dried under vacuum (5 mbar at 35 °C) and  
311 reconstituted in 250 µL of 0.1 % acetic acid in 20 % acetonitrile:methanol (1:1, v/v).

312 HPLC-QToF analysis: The analytical system used consisted of a 1260 Infinity HPLC  
313 system coupled to a 6545 quadrupole-time of flight (Q-ToF) mass spectrometer detector  
314 (Agilent, Waldbronn, Germany). Control software was Mass Hunter Workstation  
315 (version B.06.11). The Q-ToF used a Dual Jet Stream Electrospray Ionization (Dual AJS-  
316 ESI) source operated in the positive ionization mode, and the following parameters were  
317 set: capillary voltage, 4000 V; fragmentor, 120 V; nozzle voltage 500 V; gas temperature,  
318 130 °C; drying gas, 13 L/min; nebulizer, 30 psig; sheath gas temperature, 300 °C; sheath  
319 gas flow, 11 L/min; acquisition range, 80-1000 m/z. Samples were analyzed after  
320 injection (30 µl) on a Zorbax Eclipse Plus C18 Rapid Resolution HD column (2.1 × 50  
321 mm, 1.8 µm, Agilent, Santa Clara, CA) protected with a 5 mm guard column of the same  
322 material thermostated at 30 °C. The solvents system were 5 mM ammonium formate +

323 0.1% formic acid (solvent A) and 5 mM ammonium formate + 0.1% formic acid in  
324 methanol (solvent B). The elution gradient was (time, % of solvent A): 0 min, 90%; 0.5  
325 min, 90%; 10 min, 30%; 15 min, 2%; 18 min, 2%; 20 min, 90% and a post time of 5 min.  
326 Compounds were identified and quantified using the algorithm “Find by Formula” that  
327 evaluated the mass accuracy together with the isotopic relative abundance and isotopic  
328 separation.

### 329 *2.6.3. Acrylamide*

330 The quantification of acrylamide in Arabica CS extract and insoluble fraction was carried  
331 out in a laboratory that participates in appropriate proficiency testing schemes (which  
332 comply with the ‘International Harmonised Protocol for the Proficiency Testing of  
333 (Chemical) Analytical Laboratories’) developed under the auspices of  
334 IUPAC/ISO/AOAC and using approved analytical methods for detection and  
335 quantification (HPLC-MS/MS, ISO 18862:2016) according to Commission Regulation  
336 (EU) 2017/2158. This analysis was performed by Coffee Consulting S.L.

### 337 *2.6.4. Acute Toxicity Assay*

338 In accordance to OECD (Organization for Economic Co-operation and Development)  
339 Test Guidelines 425 (Up and Down Procedure), healthy young adult nulliparous and  
340 nonpregnant female rats, weighing  $180,52 \pm 6,42$  g (8 weeks old) at the start of the  
341 experiment, were procured from Charles River (Sant Cugat del Vallés, Spain) . The  
342 present study was approved by the Institutional Animal Ethics Committee (Reg. No.  
343 PROEX 011/17) of Community of Madrid, Spain.

344 Rats were housed in cages with free access to standard food (A04 Safe Diets, Augy,  
345 France) and water *ad libidum*. According to OECD Test Guidelines 425 (OECD, 2008),  
346 limit test of raw parchment (section 2.2) and husk and Arabica CS extracts (section 2.4.1)

347 was performed at 2000 mg/kg b.w. by gavage as a single dose to one rat. Husk and CS  
348 extracts were dissolved in water and parchment was dissolved in corn oil. As a control,  
349 one rat was dosed with water and another rat with corn oil to establish a comparative  
350 negative control group. Rats were closely observed for the first 30 min, then for 4 h. After  
351 survival of the treated rat, 4 additional female rats per group were administered with the  
352 same dose under the same conditions.

353 Body weight changes, signs of toxicity, behavior and mortality were observed for 24  
354 hours after administration and once daily for 14 days. Then, rats were sacrificed by  
355 exposure to excess carbon dioxide in a gas chamber for necropsy examination. Internal  
356 organs (heart, lungs, liver, kidneys, spleen, adrenals, sex organs and brain) were collected  
357 for histopathological examination.

358 Tissue samples were routinely processed for histology and fixed in buffered formalin 10%  
359 (Panreac©, Barcelona, Spain, stabilized with methanol at pH 7) for 24 hours at room  
360 temperature. Samples were then embedded in synthetic paraffin (Casa Alvarez, Madrid,  
361 Spain) with a melting point of 56 °C, using an automatic tissue processor (ASP300,  
362 Leica®, Wetzlar, Germany), with a program of automatic transmissions alcohols  
363 increased histological grading. Blocks were performed in a block forming unit (console  
364 Leica© EG1140H and cold plate Leica© EG1130) and 4 micron thick sections were  
365 obtained from rotation microtome Leica© brand, model RM 2155. Sections were  
366 deparaffinized in xylene and hydrated in alcohol and water. Conventional staining  
367 method, hematoxylin & eosin, was used by means of Leica© auto stainer SP4040. Then,  
368 dehydrated first ascending alcohol series and xylene were used, and finally, mounted with  
369 DPX (Nustain©, Nottingham, UK).

370 *2.7. Statistical analysis*

371 Data were expressed as the mean  $\pm$  SD. Prior to statistical analysis, all data were tested  
372 for variance homogeneity using the Levene's test. Logarithmic transformation of the  
373 response variable was used to achieve homocedasticity when needed. One-way analysis  
374 of variance (ANOVA) was performed. Statistical comparisons of the different treatments  
375 were performed using Tukey's test. Values of  $p < 0.05$  were considered statistically  
376 significant. All statistical analyses were performed using SPSS Statistics 24.

### 377 **3. Results and discussion**

#### 378 *3.1. Characterization of coffee husk and its potential applications*

379 Coffee husk extract (Figure 1a) showed the highest extraction yield (28 %), followed by  
380 CS extract (14 %) and parchment extract (2.3 %). Data on the characterization of coffee  
381 husk extract are summarized in Table 1. 5-CQA and caffeine were detected in all three  
382 extracts. Thus, caffeine concentration was higher than that corresponding to the phenolic  
383 compound whose concentration increased towards the fruit's core. Therefore, extract  
384 from coffee husk, the outer part of the cherry, showed the lowest values of the two  
385 bioactive compounds (1.7 mg 5-CQA/g and 13.9 mg caffeine /g). Husk extract also  
386 presented the lowest ( $p < 0.05$ ) amount of total phenolic compounds (TPC) and total  
387 flavonoid content (TFC) under the tested conditions (15.6 mg CGA eq./g and 0.9 g RUT  
388 eq./g, respectively). The lowest total antioxidant capacity (TAC) corresponded to husk  
389 extract, in accordance with the obtained data of TPC, TFC and 5-CQA content. Some  
390 authors have previously reported the amount of caffeine and CGA in raw coffee husk, 13  
391 and 25 mg/g, respectively (Murthy & Naidu, 2012; Pandey et al., 2000). Husk extract  
392 obtained under the conditions described in this work (WO 2013/004873) presented the  
393 same levels of caffeine as in the starting material.

394 Table 2 shows the viability of HepG2 cells after 24 hours of incubation with husk extract  
395 at different concentrations (0.1 to 20 mg/ml). At concentrations of 0.1 and 1 mg/ml, cell  
396 viability was not significantly ( $p > 0.05$ ) altered, while 20 mg/ml resulted cytotoxic. To  
397 date, there are no published studies about the effect of water soluble bioactive compounds  
398 from coffee husk on cell viability. Table 2 also shows the effect of husk extract on the  
399 prevention of intracellular ROS formation. The addition of tBOOH 1 mM to the culture  
400 medium significantly increased ( $p < 0.05$ ) intracellular ROS production ( $\approx 44\%$ )  
401 compared to non-treated cells. Cells pre-treated for 24 hours with husk extract (1 mg/ml)  
402 showed a significant reduction ( $p < 0.05$ ) in ROS levels compared to cells treated with  
403 tBOOH. ROS values were similar for non-treated cells (control) and those treated with  
404 the extract in absence of tBOOH, suggesting no effect in the physiological production of  
405 radicals. This is the first time that the effect of coffee husk on intracellular ROS formation  
406 has been described. Understanding the cellular defense action mechanism requires further  
407 research beyond the scope of the current study. Nevertheless, coffee husk extract should  
408 be considered a potential agent for the prevention of cellular damage induced by oxidative  
409 stress.

410 Soluble dietary fiber (SDF) was found in the insoluble fraction generated during the  
411 extraction process of coffee husk in significantly higher concentrations ( $p < 0.05$ ) than in  
412 the rest of the studied samples (12.3 %, Table 3). Total dietary fiber (TDF) values of husk  
413 insoluble residue are in the range described by Gouvea, Torres, Franca, Oliveira, &  
414 Oliveira (2009), but in lower amounts compared to the results obtained by Navya &  
415 Pushpa (2013). Coffee husk has been characterized as a source of soluble dietary fiber  
416 aimed for use in food and pharmaceutical applications. Due to its properties as an  
417 emulsifier and stabilizer, this extracted dietary fiber is a promising new ingredient for the  
418 food and beverage industry ([www.pectcof.com](http://www.pectcof.com)).

419 In addition, husk residue presented a protein content of 8.4 % (Table 3), which is within  
420 the range reported by other authors (Brand et al., 2001; Gouvea et al., 2009). Lipid content  
421 was also in the same range reported by Gouvea et al. (2009). Regarding the total  
422 antioxidant capacity of this (water-insoluble) fraction, coffee husk showed the highest  
423 values, with 48.6 mg CGA/g ( $p < 0.05$ ). This is the first time the antioxidant character  
424 of the dietary fiber from coffee husk is described. In addition to its antioxidant properties,  
425 consumption of SDF has nutritional relevance due to its positive impact on health by  
426 decreasing serum cholesterol, postprandial blood glucose and insulin contents in the  
427 human body (Bruntha Devi, Vijayabharathi, Sathyabama, Gurusiddappa Malleshi, &  
428 Brindha Priyadarisini, 2011). The ability of SDF to retard the absorption of glucose in the  
429 small intestine is a desirable characteristic in the development of foods for diabetic  
430 populations (Onyango, Noetzold, Bley, & Henle, 2004).

431 With regard to contaminants, no pesticides were detected either in husk extract or husk  
432 insoluble fraction. OTA was detected in husk insoluble fraction (4.3  $\mu\text{g}/\text{kg}$ ) in amounts  
433 below the maximum levels of 5  $\mu\text{g}/\text{kg}$  established by the European Commission  
434 (European Commission, 2005) (Table 4). In contrast, OTA levels in husk extract were  
435 below the quantification limit ( $< 0.3 \mu\text{g}/\text{kg}$ ). Furthermore, neither aflatoxin B1 nor  
436 enniantin B were detected in husk extract or its solid residue. García-Moraleja *et al.*  
437 (2015) reported OTA, Aflatoxin B1 and Enniantin B as the most frequent mycotoxins in  
438 coffee beverages (Garcia-Moraleja, Font, Manes, & Ferrer, 2015).

439 The chemical composition and antioxidant capacity of coffee husk underline its potential  
440 as a food ingredient with promising health-promoting properties. For instance, coffee  
441 pulp has been proposed as a potential source of anthocyanins (pigments widely distributed  
442 in colored fruits and flowers). Murthy et al. (2012) reported cyanadin-3-rutinoside as the  
443 major anthocyanin in coffee pulp. Interest in anthocyanins has emerged due to their

444 potential health benefits as antioxidant, anticarcinogenic, anti-inflammatory and  
445 hypoglycemic agents and as insulin sensitivity promoters (Lila, 2004). Coffee husk has  
446 also been used in the production of fructooligosaccharides (FOSs), low caloric, non-  
447 carcinogenicity fructose oligomers which promote decreasing levels of phospholipids,  
448 triglycerides, and cholesterol. This helps gut absorption of calcium and magnesium  
449 thereby stimulating probiotic growth in the human colon (Mussatto, Aguilar, Rodrigues,  
450 & Teixeira, 2009).

451 Considering the extraction yield, the chemical composition of both fractions, the *in vitro*  
452 antioxidant capacity and the effect on HepG2 intracellular ROS production, we propose  
453 the fractioning of coffee husk into two novel ingredients. First, an enriched extract with  
454 potential implementation as a food preservative, natural colorant or health-promoting  
455 food ingredient, and another fraction used as a source of antioxidant dietary fiber.

### 456 *3.2. Characterization of coffee parchment and its potential applications*

457 As an extended approach to standard analytical characterization, we investigated  
458 parchment using vibrational spectroscopy as a non-invasive technique. Vibrational  
459 spectroscopy is a reliable tool to characterize structural composition in coffee (Keidel et  
460 al., 2010) and a rapid analytical tool for extracts and future food and health applications.  
461 In the Raman spectrum (Figure 2A), a strong, broad envelope containing C-H stretching  
462 modes was observed, possibly of CH<sub>2</sub>-functional groups in the highest frequency region  
463 (2700-3600 cm<sup>-1</sup>). Main peaks were at 2937 and 2896 cm<sup>-1</sup>. At lower frequencies a peak  
464 at 2712 cm<sup>-1</sup> was identified, which is characteristic of the H-stretching modes of a O-  
465 CH<sub>2</sub>-O group or an CH<sub>3</sub> umbrella overtone. Within the broad band envelope, C-H  
466 stretching modes rising also aromatic systems have the strong contribution, since they are  
467 highly Raman active. Conversely, fewer IR-bands (Figure 2B 2700-3600cm<sup>-1</sup> region)  
468 were found. At even higher frequencies, namely at 3322 cm<sup>-1</sup>, we found a weak Raman-

469 active yet strong and broad IR-active peak. This was assigned to alcohol O-H stretching.  
470 Only two IR-active peaks were identified at lower frequencies at 2854 and 2924 cm<sup>-1</sup>,  
471 both related to CH<sub>2</sub>-stretching coordinates (Supplementary Material, Table S2).

472 At 1602 and 1658 cm<sup>-1</sup>, two C=C stretching modes with strong Raman activity and  
473 negligible IR intensity were identified. In fact, this region in the IR spectrum was rather  
474 broad. As the broad peak at 1632 cm<sup>-1</sup> might contain small contributions of amid-I  
475 modes, a low, yet detectable protein concentration in the parchment (*vide infra*) is  
476 assumed. The band at 1602 cm<sup>-1</sup> was assigned to an aromatic ring def., whereas the peak  
477 at higher frequencies contained contributions rising from a C=C double bond, possibly in  
478 the vicinity of a benzyl (or phenyl) group. At 1732 cm<sup>-1</sup> a strong IR-active and a weak  
479 Raman band associated with carbonyl C=O stretching coordinates were detected.

480 The further assignment mainly refers to the analysis of the parchment Raman spectrum.  
481 At lower frequencies, namely between 1000 and 1500 cm<sup>-1</sup>, (in-plane) bending and  
482 deformation modes gave rise to the main contribution in this region. These allowed a  
483 further structural assignment of endocarp composition. The band at 1461 cm<sup>-1</sup> was  
484 assigned to the CH<sub>3</sub> (or CH<sub>2</sub>) flowering mode of methyl (methylene) coordinates,  
485 whereas the peak at 1378 cm<sup>-1</sup> contained further coordinates of the CH<sub>2</sub>, HCC, HCO and  
486 COH bending modes. Unlike double bond stretching modes, the potential energy  
487 distribution of these vibrations is less localized and includes relatively large contributions  
488 of distinct coordinates. Deformation of the O-H groups in aliphatic and aryl alcohols  
489 contribute to the bands at 1333 and 1278 cm<sup>-1</sup>, respectively. A band containing glycoside  
490 symmetric ring breathing was located at 1095 cm<sup>-1</sup>, whereas the peak at 1038 cm<sup>-1</sup>  
491 mainly contained contributions of C-C and C-O stretching coordinates. Finally, the peak  
492 at 899 cm<sup>-1</sup> was assigned to HCC and HCO bending modes. The adjacent peak at 926  
493 cm<sup>-1</sup> probably contained further angle bending modes (CHO, CCH or similar), however

494 an unambiguous assignment was not possible. While the Raman spectrum contained very  
495 strong bending modes, only a broad and prominent envelope at 1011 cm<sup>-1</sup> with band  
496 shoulders at 1048 and 1100 cm<sup>-1</sup> was found. Here the highest peak and further low-  
497 frequency peaks could be assigned to C-O stretching modes of aryl alcohols, whereas the  
498 high frequency shoulders contained C-O stretching contributions of aliphatic alcohols,  
499 esters or ethers.

500 In agreement with physicochemical studies on coffee parchment composition, the bands  
501 at 1658 and 1602 cm<sup>-1</sup> have been assigned to lignin. Conversely, cellulose did not display  
502 any bands in this region. The corresponding marker bands were observed at 899, 1095  
503 and 1121 cm<sup>-1</sup>. In the high frequency region (2700-3500 cm<sup>-1</sup>), the strongest peak at  
504 2937 cm<sup>-1</sup> corresponds to lignin, whereas the adjacent peaks at 2896 and 2712 cm<sup>-1</sup>  
505 correspond to the C-H stretching modes of cellulose. The O-H stretching peak has  
506 contributions from both molecules. The strong C=O stretching mode observed at 1732  
507 cm<sup>-1</sup> could not be assigned to either of the previously mentioned compounds. It could  
508 possibly reflect a third compound group, such as hemicellulose. The observed IR and  
509 Raman activity of the C=O group (*vide supra*), can only be understood if the C=O group  
510 in discussion is conjugated, as for example in ferulated xylan. The presence of ashes, coal  
511 and dark materials give rise to strong thermal emission in the Raman spectrum. If present,  
512 the contribution is low or negligible.

513 Two components could be unambiguously confirmed in coffee parchment, namely lignin  
514 and cellulose. Bekalo and Reinhardt (2010) also reported the presence of these  
515 compounds in coffee parchment. Strong indications for a third species were presented.  
516 This combined spectroscopic approach provides complementary information. However,  
517 further studies of different parchment varieties, vibrational spectra of isolated compounds  
518 as well as extended chemometric analysis are required. It is interesting to note that the

519 obtained results provide the first vibrational spectroscopy (IR and Raman)  
520 characterization of coffee by-products.

521 Parchment showed the lowest extraction yield (2.3 %). This extract (Figure 1b) presented  
522 low levels of 5-CQA (6.1 mg/g); however, it had the highest caffeine content (58.2 mg/g)  
523 ( $p < 0.05$ ). No bibliographical references have been found regarding caffeine and CGA  
524 content in aqueous extracts of coffee parchment. As shown in Table 1, parchment extract  
525 had the greatest antioxidant capacity ( $p < 0.05$ ) of all the samples analyzed, followed by  
526 CS extracts. Parchment and Robusta CS extract showed the highest TPC and TFC ( $p <$   
527  $0.05$ ). Table 2 includes data on the effect of parchment extract on cell viability and  
528 intracellular ROS formation. At concentrations of 0.1 and 1 mg/ml cell viability was not  
529 significantly ( $p > 0.05$ ) altered, while 10 and 20 mg/ml resulted cytotoxic. Pre-treatment  
530 of cells with parchment extract for 24 hours (1 mg/ml) significantly reduced ( $p < 0.05$ )  
531 ROS levels compared to non-treated cells and cells treated with tBOOH. Parchment  
532 extract had a greater antioxidant effect on HepG2 cells compared to husk extract. Results  
533 are in line with those corresponding to TAC and TPC. Parchment extract presented the  
534 highest values for these parameters among the studied samples. To the best of our  
535 knowledge, no studies have been published on the effect of coffee parchment on cell  
536 viability and intracellular ROS formation.

537 The insoluble fraction of coffee parchment obtained after extraction presented the highest  
538 amount of TDF ( $p < 0.05$ ) (92.6 %) mainly composed of IDF (92.1 %) (Table 3). Protein  
539 content in the parchment insoluble fraction was 3.1%, which was the lowest value  
540 compared to the rest of the samples. Our results showed that parchment insoluble fraction  
541 also displayed the lowest ( $p < 0.05$ ) lipid content (0.3 %). To date, no other data on protein  
542 and lipid content of coffee parchment have been published. Furthermore, it also had the  
543 lowest antioxidant capacity (4.1 mg CGA/g).

544 Considering the food safety of raw parchment, none of the studied pesticides were  
545 detected in this sample. Aflatoxin B1 and enniantin B were not present in parchment  
546 (Table 4). However, OTA was detected at 2.7 µg/kg, which is under the limit established  
547 by the European Commission (5 µg/kg). Therefore, thermal stabilization of the raw  
548 material should be thermally stabilized to avoid microbial contamination and the  
549 production of mycotoxins during its storage, to increase the shelf-life of the sample and  
550 to ensure its safe use as a food ingredient for human beings.

551 Until now, applications for coffee parchment as a food ingredient remain largely  
552 unexplored. Despite the promising chemical composition of aqueous parchment extract  
553 and its high *in vitro* antioxidant capacity, we suggest that coffee parchment should be  
554 used as it is after a simple thermal stabilization, since extraction yields are rather poor.  
555 New applications of coffee parchment in the food industry as a natural source of  
556 antioxidant dietary fiber should be considered to revalue this by-product.

### 557 *3.3. Characterization of coffee silverskin and its potential applications*

558 The extraction yield of water-soluble compounds from CS (Figure 1c) was similar for  
559 Arabica and Robusta species. As CS is in direct contact with the coffee seeds, CS extract  
560 presented the highest content of 5-CQA (9.4 mg/g and 21.3 mg/g, for Arabica and  
561 Robusta species, respectively). The concentration ratio of caffeine/5-CQA was 2.5 in both  
562 the Arabica and Robusta CS extract. However, values of both compounds were lower in  
563 the Arabica CS extract (Table 1). The obtained caffeine content was higher than that  
564 reported by other authors (Bresciani, Calani, Bruni, Brighenti, & Del Rio, 2014;  
565 Napolitano, Fogliano, Tafuri, & Ritieni, 2007). In agreement with our results, Mesías et  
566 al. (2014) found that 5-CQA and caffeine content were significantly higher in Robusta  
567 than in Arabica CS extract.

568 EFSA reported a safety level for daily caffeine consumption of 400 mg for the general  
569 population and 200 mg for lactating women. There is currently not enough information  
570 available to determine a safe level of caffeine intake for children (European Food Safety  
571 Authority (EFSA), 2015). Caffeine also presents beneficial health effects due to its  
572 stimulating properties (Nehlig, 2016). Arabica and Robusta CS extracts could be a natural  
573 source of sustainable caffeine. So far, published results suggest that caffeine content  
574 should not be considered as a safety concern in the application of coffee by-products as  
575 food ingredients (Garcia-Serna et al., 2014; Martinez-Saez et al., 2014). However, more  
576 conclusive studies are required.

577 Robusta CS extract presented significantly higher amounts of TPC and TFC than Arabica  
578 CS extract ( $p < 0.05$ ), and TPC and TFC concentration was significantly higher in these  
579 two species than in husk extract ( $p < 0.05$ ) (Table 1). In this case, the TPC value obtained  
580 for CS extracts was in the same order of magnitude as the values obtained by Mesías et  
581 al. (2014). The antioxidant capacity values of the different CS extracts did not differ  
582 significantly ( $p > 0.05$ ). There is usually a greater proportion of antioxidants in Robusta  
583 CS than in Arabica CS (Mesías et al., 2014; Napolitano et al., 2007). Other authors have  
584 also confirmed the antioxidant character and high phenolic and flavonoid content in this  
585 sample (Borrelli et al., 2004; Rodrigues, Palmeira-de-Oliveira, et al., 2015a). Although  
586 TAC was significantly greater in parchment extract ( $p < 0.05$ ), parchment extraction yield  
587 is low. Therefore, CS could be a more suitable source of bioactive compounds with a  
588 higher antioxidant capacity. The application of CS as a natural source of antioxidants has  
589 been previously described (del Castillo et al., 2013; Garcia-Serna, Martinez-Saez, Mesias,  
590 Morales, & Castillo, 2014; Iriondo-DeHond et al., 2016; Mesías et al., 2014; Rodrigues,  
591 Palmeira-de-Oliveira, et al., 2015).

592 Since Arabica coffee represents around 70 % of global production (Bunn, Läderach,  
593 Ovalle Rivera, & Kirschke, 2015), this species was used in the *in vitro* studies. Arabica  
594 CS extract was not cytotoxic in HepG2 cells at none of the tested concentrations. For  
595 doses of 10 and 20 mg/ml of Arabica CS extract, cell viability increased significantly ( $p$   
596  $< 0.05$ ) (Table 2). The non-cytotoxic character of CS has also been studied in other cell  
597 lines (Iriondo-DeHond et al., 2016; Iriondo-DeHond, Haza, Ávalos, del Castillo, &  
598 Morales, 2017; Rodrigues, Gaspar, et al., 2015). Arabica CS extract was more effective  
599 in the reduction of induced ROS than husk extract, and as effective as parchment extract.  
600 This is in agreement with their corresponding antioxidant capacity studied *in vitro*. The  
601 protective effect of CS in the reduction of induced ROS formation has been reported for  
602 other cell lines (Fernandez-Gomez, Ramos, et al., 2016; Iriondo-DeHond et al., 2016).

603 The TDF content in the remaining CS solid insoluble fraction after the aqueous extraction  
604 process was similar for Arabica and Robusta varieties (Table 3). Values obtained for CS  
605 dietary fiber are similar to those previously described (Ballesteros et al., 2014; Borrelli et  
606 al., 2004; Garcia-Serna et al., 2014). Most of the dietary fiber found in the sample was  
607 insoluble, and values of SDF were lower than those recently described (Esquivel &  
608 Jiménez, 2012; Napolitano et al., 2007). In agreement with previous results (Ballesteros  
609 et al., 2014; Borrelli et al., 2004; Pourfarzad et al., 2013), the highest protein content  
610 corresponded to Robusta CS (19 %). Lipid content of the insoluble residues of CS was  
611 2.6 % and 3.6 % for Arabica and Robusta, respectively, in the same range as that  
612 previously described by Ballesteros et al. (2014) and Borrelli et al. (2004).

613 Food safety of Arabica CS was validated by the absence of pesticides, aflatoxin B1 and  
614 enniantin B (Table 4). OTA was present in the insoluble fraction of this sample (2.9  
615  $\mu\text{g}/\text{kg}$ ) in a concentration below the limit established by the European Commission (5  
616  $\mu\text{g}/\text{kg}$ ) and was not detected in CS extract. Other authors have described the presence of

617 OTA in CS (Toschi, Cardenia, Bonaga, Mandrioli, & Rodriguez-estrada, 2014). The  
618 analysis of this contaminant is mandatory to ensure the safety of CS extract as a food  
619 ingredient. Since CS is the only by-product generated during the roasting process of  
620 coffee beans, acrylamide content was also studied in this sample. Levels of 489  $\mu\text{g}/\text{kg}$  of  
621 this water-soluble compound were found in CS extract. This value is between the values  
622 established by the European Commission for roasted coffee (400  $\mu\text{g}/\text{kg}$ ) and instant  
623 coffee (850  $\mu\text{g}/\text{kg}$ ) (European Commission, 2017). In contrast, CS insoluble fraction  
624 presented levels  $< 50 \mu\text{g}/\text{kg}$ . Mitigation of acrylamide in CS can be achieved following  
625 the instructions recommended for roasted and instant coffees by the European  
626 Commission.

627 Arabica CS, CS extract and the insoluble fraction recovered from the water extraction  
628 process have been implemented to improve the formulation of biscuits. In this study, CS  
629 was used as a natural colorant and as a source of fiber (Garcia-Serna et al., 2014). CS has  
630 also been used to improve the quality, shelf-life, sensory qualities and appearance of  
631 Barbari bread (Pourfarzad et al., 2013). Other authors proposed the use of CS as a source  
632 of prebiotics (Borrelli et al., 2004; Jiménez-Zamora, Pastoriza, & Rufián-Henares, 2015;  
633 Mussatto et al., 2013). In addition, CS has been used in combination with roasted coffee  
634 and cocoa powder to obtain innovative coffee blends that were rich in bioactive  
635 compounds, such as CGAs, trigonelline, theobromine, and caffeine (Ribeiro, Leitão,  
636 Ramalho, & Lidon, 2014). Other novel antioxidant beverages based on raw CS and CS  
637 extract from Arabica and Robusta species have been developed to study their inhibitory  
638 effect on fat accumulation using *Caenorhabditis elegans* as an *in vivo* model (Martinez-  
639 Saez et al., 2014). According to our results, improved fractioning of CS will yield a  
640 superior aqueous extract rich in bioactive compounds with the potential to reduce diabetes

641 and obesity, among other conditions. Similarly, the potential of coffee husk extract to  
642 reduce the risk of chronic diseases should also be studied.

643 Considering the extraction yield and chemical characterization of each by-product  
644 fraction, we suggest single-fraction separation and development of two new  
645 multifunctional ingredients. As proposed for coffee husk, CS is a suitable source for an  
646 enriched extract which could be implemented as a food ingredient with antioxidant and  
647 antidiabetic properties (del Castillo et al., 2016). Furthermore, the insoluble fraction can  
648 be used as a source of antioxidant dietary fiber.

#### 649 *3.4. Food toxicity of novel ingredients*

650 Single oral administration of the three validated coffee by-products (raw parchment and  
651 husk and Arabica CS extracts) at a dose of 2000 mg/kg b.w. showed no visible signs of  
652 toxicity, abnormal behavior or mortality. Relative organ weights are shown in Table S3  
653 (Supplementary Material). No significant differences ( $p > 0.05$ ) were found among  
654 groups. Intake of an acute dose of ingredients (2000 mg/kg b.w) did not cause significant  
655 changes in histological parameters of vital organs. Livers of the control and treated rats  
656 showing normal hepatic lobules, polyhedral hepatocytes with central vesicular nuclei and  
657 eosinophilic granular cytoplasm are shown in Figure 3a and 3b, respectively. Similarly,  
658 the control and treated kidney sections showed a normal histological picture composed  
659 of renal corpuscles, which appear as rounded structures, and glomeruli are surrounded by  
660 narrow Bowman's spaces and cortical tubules with a small number of distal convoluted  
661 tubules and collecting tubules (Figure 3c and 3d).

662 To the best of our knowledge, very few studies regarding the presence of contaminants  
663 in coffee by-products have been previously published. No lethal effects were observed in  
664 rats treated with 2000 mg/kg b.w. by oral administration. Results are supported by data

665 on contaminants. As previously discussed, pesticides, aflatoxin B1 and enniatin B were  
666 not found in the coffee by-products; OTA was detected in raw parchment and in the  
667 insoluble fraction of husk and CS in concentrations lower than 5 µg/kg; acrylamide was  
668 only detected in CS and its derivatives in lower or similar concentrations to those found  
669 in roasted coffee (400 µg/kg) (European Commission, 2017). Caffeine concentrations can  
670 be considered safe and were lower than concentrations reported for coffee beverages  
671 (European Food Safety Authority (EFSA), 2015).

#### 672 **4. Conclusions**

673 This study provided data for the validation of CS and coffee husk as two novel safe food  
674 ingredients: an enriched extract with multifunctional properties and an insoluble fraction  
675 composed of dietary fiber of different nature (soluble and insoluble). The fractioning  
676 procedure is a sustainable, low-cost process which could be easily performed by the  
677 coffee industry without the production of new waste. We also propose that coffee  
678 parchment be used as a single food ingredient composed of antioxidant insoluble dietary  
679 fiber after a thermal stabilization process. Thus, coffee by-products can be converted into  
680 promising safe health-promoting food ingredients, providing sustainable economic and  
681 environmental benefits. Analysis of contaminants is essential to achieve this goal.

#### 682 **Acknowledgements**

683 The SUSCOFFEE Project (AGL2014-57239-R): “Sustainable coffee production and  
684 consumption: Valorization of coffee waste into food ingredients” funded this work. A.  
685 Iriondo-DeHond is a fellow of the FPI predoctoral program of MINECO (BES-2015-  
686 072191). The authors thank Dr. Miguel Herrero for his help with the UPLC-MS/MS  
687 analysis.

#### 688 **References**

- 689 Açar, Ö. C., Gökmen, V., Pellegrini, N., & Fogliano, V. (2009). Direct evaluation of the  
690 total antioxidant capacity of raw and roasted pulses, nuts and seeds. *European*  
691 *Food Research and Technology*, 229(6), 961–969. [https://doi.org/10.1007/s00217-](https://doi.org/10.1007/s00217-009-1131-z)  
692 009-1131-z
- 693 Adebajo, M. O., Frost, R. L., Kloprogge, J. T., & Kokot, S. (2006). Raman  
694 spectroscopic investigation of acetylation of raw cotton. *Spectrochimica Acta Part*  
695 *A: Molecular and Biomolecular Spectroscopy*, 64(2), 448–453.  
696 <https://doi.org/10.1016/J.SAA.2005.07.045>
- 697 Afsah-Hejri, L., Jinap, S., Hajeb, P., Radu, S., & Shakibazadeh, S. (2013). A Review on  
698 Mycotoxins in Food and Feed: Malaysia Case Study. *Comprehensive Reviews in*  
699 *Food Science and Food Safety*, 12(6), 629–651. [https://doi.org/10.1111/1541-](https://doi.org/10.1111/1541-4337.12029)  
700 4337.12029
- 701 Bakondi, E., Gönczi, M., Szabó, É., Bai, P., Pacher, P., Gergely, P., ... Virág, L. (2003).  
702 Role of intracellular calcium mobilization and cell-density-dependent signaling in  
703 oxidative-stress-induced cytotoxicity in HaCaT keratinocytes. *Journal of*  
704 *Investigative Dermatology*, 121(1), 88–95. [https://doi.org/10.1046/j.1523-](https://doi.org/10.1046/j.1523-1747.2003.12329.x)  
705 1747.2003.12329.x
- 706 Ballesteros, L. F., Teixeira, J. a., & Mussatto, S. I. (2014). Chemical, Functional, and  
707 Structural Properties of Spent Coffee Grounds and Coffee Silverskin. *Food and*  
708 *Bioprocess Technology*, 7(12), 3493–3503. [https://doi.org/10.1007/s11947-014-](https://doi.org/10.1007/s11947-014-1349-z)  
709 1349-z
- 710 Bekalo, S. A., & Reinhardt, H.-W. W. (2010). Fibers of coffee husk and hulls for the  
711 production of particleboard. *Materials and Structures*, 43(8), 1049–1060.  
712 <https://doi.org/DOI 10.1617/s11527-009-9565-0>

713 Belitz, H.-D., Grosch, W., & Schieberle, P. (2009). Coffee, tea, cocoa. In *In: Food*  
714 *Chemistry. Eds. Belitz, H.-D., Grosch, W. and Schieberle, P., 4th edition, Springer,*  
715 *Leipzig* (pp. 938–951).

716 Borrelli, R. C., Esposito, F., Napolitano, A., Ritieni, A., & Fogliano, V. (2004).  
717 Characterization of a new potential functional ingredient: coffee silverskin. *Journal*  
718 *of Agricultural and Food Chemistry*, 52(5), 1338–1343.  
719 <https://doi.org/10.1021/jf034974x>

720 Borrelli, R. C., Visconti, A., Mennella, C., Anese, M., & Fogliano, V. (2002). Chemical  
721 characterization and antioxidant properties of coffee melanoidins. *Journal of*  
722 *Agricultural and Food Chemistry*, 50(22), 6527–33.

723 Brand, D., Pandey, a., Rodriguez-Leon, J. a., Roussos, S., Brand, I., & Soccol, C. R.  
724 (2001). Packed bed column fermenter and kinetic modeling for upgrading the  
725 nutritional quality of coffee husk in solid-state fermentation. *Biotechnology*  
726 *Progress*, 17(6), 1065–1070. <https://doi.org/10.1021/bp010112+>

727 Bresciani, L., Calani, L., Bruni, R., Brighenti, F., & Del Rio, D. (2014). Phenolic  
728 composition, caffeine content and antioxidant capacity of coffee silverskin. *Food*  
729 *Research International*, 61, 196–201.  
730 <https://doi.org/10.1016/j.foodres.2013.10.047>

731 Bruntha Devi, P., Vijayabharathi, R., Sathyabama, S., Gurusiddappa Malleshi, N., &  
732 Brindha Priyadarisini, V. (2011). Health benefits of finger millet (Eleusine  
733 coracana L.) polyphenols and dietary fiber: a review.  
734 <https://doi.org/10.1007/s13197-011-0584-9>

735 Bunn, C., Läderach, P., Ovalle Rivera, O., & Kirschke, D. (2015). A bitter cup: climate  
736 change profile of global production of Arabica and Robusta coffee. *Climatic*

737 *Change*, 129(1–2), 89–101. <https://doi.org/10.1007/s10584-014-1306-x>

738 Chanakya, H. N., & De Alwis, A. A. P. (2004). Environmental Issues and Management  
739 in Primary Coffee Processing. *Process Safety and Environmental Protection*,  
740 82(4), 291–300. <https://doi.org/10.1205/095758204323162319>

741 Contini, M., Baccelloni, S., Massantini, R., & Anelli, G. (2008). Extraction of natural  
742 antioxidants from hazelnut (*Corylus avellana* L.) shell and skin wastes by long  
743 maceration at room temperature. *Food Chemistry*, 110(3), 659–669.  
744 <https://doi.org/10.1016/j.foodchem.2008.02.060>

745 del Castillo, M. D., Fernandez-Gomez, B., Martinez-Saez, N., Iriondo-DeHond, A.,  
746 Martirosyan, D. M., & Mesa, M. D. (2016). Coffee silverskin extract for aging and  
747 chronic diseases. In D. M. Martirosyan (Ed.), *Functional Foods In Health And*  
748 *Disease*.

749 del Castillo, M. D., Fernandez-Gomez, B., Martinez-Saez, N., Iriondo-DeHond, A., &  
750 Mesa, M. D. (2018). Coffee By-Products. In A. Farah (Ed.), *Coffee: Chemistry,*  
751 *Quality and Health Implications*. Oxfordshire: Royal Society of Chemistry.

752 del Castillo, M. D., Ibañez, M. E., Amigo, M., Herrero, M., Plaza del Moral, M., &  
753 Ullate, M. (2013, January). Application of products of coffee silverskin in anti-  
754 ageing cosmetics and functional food. WO 2013/004873.

755 del Castillo, M. D., Iriondo-DeHond, A., Fernandez-Gomez, B., Martinez-Saez, N.,  
756 Iriondo-DeHond, M., & Zhou, J.-R. (2017). Applications of recovered compounds  
757 in food products. In C. Galanakis (Ed.), *Handbook of Coffee Processing By-*  
758 *Products: Sustainable Applications* (First Edit). Academic Press-Elsevier.

759 Esquivel, P., & Jiménez, V. M. (2012). Functional properties of coffee and coffee by-

760 products. *Food Research International*, 46(2), 488–495.  
761 <https://doi.org/10.1016/j.foodres.2011.05.028>

762 European Commission. (2005). COMMISSION REGULATION (EC) No 123/2005 of  
763 26 January 2005 amending Regulation (EC) No 466/2001 as regards ochratoxin A.  
764 *Official Journal of the European Union*, 2005(123), 2004–2006. Retrieved from  
765 [http://eur-lex.europa.eu/legal-](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32005R0123&qid=1478028660364&from=ES)  
766 [content/EN/TXT/PDF/?uri=CELEX:32005R0123&qid=1478028660364&from=E](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32005R0123&qid=1478028660364&from=ES)  
767 [S](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32005R0123&qid=1478028660364&from=ES)

768 European Commission. (2014). Proposal for a directive of the European parliament and  
769 of the council amending Directives 2008/98/EC on waste, 94/62/EC on packaging  
770 and packaging waste, 1999/31/EC on the landfill of waste, 2000/53/EC on end-of-  
771 life vehicles, 2006/66/EC on batteries and. *Official Journal of the European*  
772 *Union.*, 0201, 1–23. <https://doi.org/10.1017/CBO9781107415324.004>

773 European Commission. (2017). COMMISSION REGULATION (EU) 2017/2158 of 20  
774 November 2017 establishing mitigation measures and benchmark levels for the  
775 reduction of the presence of acrylamide in food. *Official Journal of the European*  
776 *Union*, 204, 24–44. [https://doi.org/http://eur-](https://doi.org/http://eur-lex.europa.eu/pri/en/oj/dat/2003/l_285/l_28520031101en00330037.pdf)  
777 [lex.europa.eu/pri/en/oj/dat/2003/l\\_285/l\\_28520031101en00330037.pdf](https://doi.org/http://eur-lex.europa.eu/pri/en/oj/dat/2003/l_285/l_28520031101en00330037.pdf)

778 European Food Safety Authority (EFSA). (2015). Scientific Opinion on the safety of  
779 caffeine. *EFSA Journal*, 13(5), 1–21. <https://doi.org/10.2903/j.efsa.2015.4102>

780 European Food Safety Authority (EFSA). (2016). *Chemicals in food 2016*. Parma.  
781 <https://doi.org/10.2805/359743>

782 Fernandez-Gomez, B., Lezama, A., Amigo-Benavent, M., Ullate, M., Herrero, M.,  
783 Martín, M. Á., ... del Castillo, M. D. (2016). Insights on the health benefits of the

784 bioactive compounds of coffee silverskin extract. *Journal of Functional Foods*, 25,  
785 197–207. <https://doi.org/10.1016/j.jff.2016.06.001>

786 Fernandez-Gomez, B., Ramos, S., Goya, L., Mesa, M. D., del Castillo, M. D., & Martín,  
787 M. Á. (2016). Coffee silverskin extract improves glucose-stimulated insulin  
788 secretion and protects against streptozotocin-induced damage in pancreatic INS-1E  
789 beta cells. *Food Research International*.  
790 <https://doi.org/10.1016/j.foodres.2016.03.006>

791 Folch, J., Lees, M., & Sloane Stanley, G. . (1987). a Simple, 55(5), 999–1033.  
792 <https://doi.org/10.1016/j.ultrasmedbio.2011.03.005>

793 Gamboa-Gaitán, M. Á. (2012). Presencia de aspergillus y otros simbioses fúngicos en  
794 granos de café procedentes de Colombia, 17(1), 39–50. Retrieved from  
795 [https://www.scopus.com/record/display.uri?eid=2-s2.0-](https://www.scopus.com/record/display.uri?eid=2-s2.0-84861470773&origin=inward&txGid=6c7e709f19507c1f886634e01289d794)  
796 [84861470773&origin=inward&txGid=6c7e709f19507c1f886634e01289d794](https://www.scopus.com/record/display.uri?eid=2-s2.0-84861470773&origin=inward&txGid=6c7e709f19507c1f886634e01289d794)

797 Garcia-Moraleja, A., Font, G., Manes, J., & Ferrer, E. (2015). Analysis of mycotoxins  
798 in coffee and risk assessment in Spanish adolescents and adults. *Food and*  
799 *Chemical Toxicology*, 86, 225–233. <https://doi.org/10.1016/j.fct.2015.10.014>

800 Garcia-Serna, E., Martinez-Saez, N., Mesias, M., Morales, F. J., & Castillo, M. D. Del.  
801 (2014). Use of Coffee Silverskin and Stevia to Improve the Formulation of  
802 Biscuits. *Polish Journal of Food and Nutrition Sciences*, 64(4), 243–251.  
803 <https://doi.org/10.2478/pjfn-2013-0024>

804 Gierlinger, N., Keplinger, T., & Harrington, M. (2012). Imaging of plant cell walls by  
805 confocal Raman microscopy. *Nature Protocols*, 7(9), 1694–1708.  
806 <https://doi.org/10.1038/nprot.2012.092>

807 Gierlinger, N., Keplinger, T., Harrington, M., & Schwanninger, M. (2013). Raman  
808 Imaging of Lignocellulosic Feedstock. In *Cellulose - Biomass Conversion*. InTech.  
809 <https://doi.org/10.5772/50878>

810 Gomes, A., Fernandes, E., & Lima, J. L. F. C. (2005). Fluorescence probes used for  
811 detection of reactive oxygen species. *Journal of Biochemical and Biophysical*  
812 *Methods*, 65(2–3), 45–80. <https://doi.org/10.1016/j.jbbm.2005.10.003>

813 Gouvea, B. M., Torres, C., Franca, A. S., Oliveira, L. S., & Oliveira, E. S. (2009).  
814 Feasibility of ethanol production from coffee husks. *Biotechnology Letters*, 31(9),  
815 1315–1319. <https://doi.org/10.1007/s10529-009-0023-4>

816 Iriondo-DeHond, A., Haza, A. I., Ávalos, A., del Castillo, M. D., & Morales, P. (2017).  
817 Validation of coffee silverskin extract as a food ingredient by the analysis of  
818 cytotoxicity and genotoxicity. *Food Research International*, 100, 791–797.  
819 <https://doi.org/10.1016/j.foodres.2017.08.012>

820 Iriondo-DeHond, A., Martorell, P., Genovés, S., Ramón, D., Stamatakis, K., Fresno, M.,  
821 ... Del Castillo, M. D. (2016). Coffee silverskin extract protects against accelerated  
822 aging caused by oxidative agents. *Molecules*, 21(6), 1–14.  
823 <https://doi.org/10.3390/molecules21060721>

824 Jiménez-Zamora, A., Pastoriza, S., & Rufián-Henares, J. a. (2015). Revalorization of  
825 coffee by-products. Prebiotic, antimicrobial and antioxidant properties. *LWT -*  
826 *Food Science and Technology*, 61(1), 12–18.  
827 <https://doi.org/10.1016/j.lwt.2014.11.031>

828 Keidel, A., von Stetten, D., Rodrigues, C., Máguas, C., & Hildebrandt, P. (2010).  
829 Discrimination of Green Arabica and Robusta Coffee Beans by Raman  
830 Spectroscopy. *Journal of Agricultural and Food Chemistry*, 58(21), 11187–11192.

831 <https://doi.org/10.1021/jf101999c>

832 Lila, M. A. (2004). Anthocyanins and Human Health: An In Vitro Investigative  
833 Approach. *Journal of Biomedicine & Biotechnology*, 2004(5), 306–313.  
834 <https://doi.org/10.1155/S111072430440401X>

835 Martinez-Saez, N., Ullate, M., Martin-Cabrejas, M. a., Martorell, P., Genovés, S.,  
836 Ramon, D., & Del Castillo, M. D. (2014). A novel antioxidant beverage for body  
837 weight control based on coffee silverskin. *Food Chemistry*, 150, 227–234.  
838 <https://doi.org/10.1016/j.foodchem.2013.10.100>

839 Mesías, M., Navarro, M., Martínez-Saez, N., Ullate, M., del Castillo, M. D., & Morales,  
840 F. J. (2014). Antiglycative and carbonyl trapping properties of the water soluble  
841 fraction of coffee silverskin. *Food Research International*, 62, 1120–1126.  
842 <https://doi.org/10.1016/j.foodres.2014.05.058>

843 Murthy, P. S., Manjunatha, M., Sulochannama, G., & Madhava Naidu, M. (2012).  
844 Extraction, Characterization and Bioactivity of Coffee Anthocyanins. *European*  
845 *Journal of Biological Sciences*, 4(1), 13–19.  
846 <https://doi.org/10.5829/idosi.ejbs.2012.4.1.6149>

847 Murthy, P. S., & Naidu, M. M. (2012). Recovery of Phenolic Antioxidants and  
848 Functional Compounds from Coffee Industry By-Products. *Food and Bioprocess*  
849 *Technology*, 5(3), 897–903. <https://doi.org/10.1007/s11947-010-0363-z>

850 Mussatto, S. I., Aguilar, C. N., Rodrigues, L. R., & Teixeira, J. A. (2009).  
851 Fructooligosaccharides and  $\beta$ -fructofuranosidase production by *Aspergillus*  
852 *japonicus* immobilized on lignocellulosic materials. *Journal of Molecular*  
853 *Catalysis B: Enzymatic*, 59(1–3), 76–81.  
854 <https://doi.org/10.1016/j.molcatb.2009.01.005>

- 855 Mussatto, S. I., Ballesteros, L. F., Martins, S., Maltos, D. A. F., Aguilar, C. N., &  
856 Teixeira, J. A. (2013). Maximization of Fructooligosaccharides and  $\beta$ -  
857 Fructofuranosidase Production by *Aspergillus japonicus* under Solid-State  
858 Fermentation Conditions. *Food and Bioprocess Technology*, 6(8), 2128–2134.  
859 <https://doi.org/10.1007/s11947-012-0873-y>
- 860 Nabais, J. M. V., Nunes, P., Carrott, P. J. M., Ribeiro Carrott, M. M. L., García, a. M.,  
861 & Díaz-Díez, M. a. (2008). Production of activated carbons from coffee endocarp  
862 by CO<sub>2</sub> and steam activation. *Fuel Processing Technology*, 89(3), 262–268.  
863 <https://doi.org/10.1016/j.fuproc.2007.11.030>
- 864 Napolitano, A., Fogliano, V., Tafuri, A., & Ritieni, A. (2007). Natural occurrence of  
865 ochratoxin A and antioxidant activities of green and roasted coffees and  
866 corresponding byproducts. *Journal of Agricultural and Food Chemistry*, 55(25),  
867 10499–10504. <https://doi.org/10.1021/jf071959+>
- 868 Navya, P. N., & Pushpa, S. M. (2013). Production, statistical optimization and  
869 application of endoglucanase from *Rhizopus stolonifer* utilizing coffee husk.  
870 *Bioprocess and Biosystems Engineering*, 36(8), 1115–1123.  
871 <https://doi.org/10.1007/s00449-012-0865-3>
- 872 Nehlig, A. (2016). Effects of coffee/caffeine on brain health and disease: What should I  
873 tell my patients? *Practical Neurology*, 16(2), 89–95.  
874 <https://doi.org/10.1136/practneurol-2015-001162>
- 875 Oki, T., Nagai, S., Yoshinaga, M., Nishiba, Y., & Suda, I. (2006). Contribution of b-  
876 Carotene to Radical Scavenging Capacity Varies among Orange-fleshed Sweet  
877 Potato Cultivars. *Food Sci. Technol. Res*, 12, 156–160.
- 878 Onyango, C., Noetzold, H., Bley, T., & Henle, T. (2004). Proximate composition and

879 digestibility of fermented and extruded uji from maize–finger millet blend. *LWT -*  
880 *Food Science and Technology*, 37(8), 827–832.  
881 <https://doi.org/10.1016/J.LWT.2004.03.008>

882 Pandey, A., Soccol, C. R., Nigam, P., Brand, D., Mohan, R., & Roussos, S. (2000).  
883 Biotechnological potential of coffee pulp and coffee husk for bioprocesses.  
884 *Biochemical Engineering Journal*, 6(2), 153–162. [https://doi.org/10.1016/S1369-](https://doi.org/10.1016/S1369-703X(00)00084-X)  
885 [703X\(00\)00084-X](https://doi.org/10.1016/S1369-703X(00)00084-X)

886 Pourfarzad, A., Mahdavian-Mehr, H., & Sedaghat, N. (2013). Coffee silverskin as a  
887 source of dietary fiber in bread-making: Optimization of chemical treatment using  
888 response surface methodology. *LWT - Food Science and Technology*, 50(2), 599–  
889 606. <https://doi.org/10.1016/j.lwt.2012.08.001>

890 Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., & Rice-Evans, C. (1999).  
891 Antioxidant activity applying an improved ABTS radical cation decolorization  
892 assay. *Free Radical Biology and Medicine*, 26(9–10), 1231–1237.  
893 [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3)

894 Ribeiro, V. S., Leitão, A. E., Ramalho, J. C., & Lidon, F. C. (2014). Chemical  
895 characterization and antioxidant properties of a new coffee blend with cocoa,  
896 coffee silverskin and green coffee minimally processed. *Food Research*  
897 *International*, 61, 39–47. <https://doi.org/10.1016/j.foodres.2014.05.003>

898 Rodrigues, F., Gaspar, C., Palmeira-de-Oliveira, A., Sarmiento, B., Helena Amaral, M.,  
899 & P P Oliveira, M. B. (2015). Application of Coffee Silverskin in cosmetic  
900 formulations: physical/antioxidant stability studies and cytotoxicity effects. *Drug*  
901 *Development and Industrial Pharmacy*, 00(00), 1–8.  
902 <https://doi.org/10.3109/03639045.2015.1035279>

903 Rodrigues, F., Palmeira-de-Oliveira, A., das Neves, J., Sarmento, B., Amaral, M. H., &  
904 Oliveira, M. B. P. P. (2015a). Coffee silverskin: a possible valuable cosmetic  
905 ingredient. *Pharmaceutical Biology*, 53(3), 386–94.  
906 <https://doi.org/10.3109/13880209.2014.922589>

907 Rodrigues, F., Palmeira-de-Oliveira, A., das Neves, J., Sarmento, B., Amaral, M. H., &  
908 Oliveira, M. B. P. P. (2015b). Coffee silverskin: a possible valuable cosmetic  
909 ingredient. *Pharmaceutical Biology*, 53(3), 386–94.  
910 <https://doi.org/10.3109/13880209.2014.922589>

911 Schenzel, K., & Fischer, S. (2001). NIR FT Raman Spectroscopy—a Rapid Analytical  
912 Tool for Detecting the Transformation of Cellulose Polymorphs. *Cellulose*, 8(1),  
913 49–57. <https://doi.org/10.1023/A:1016616920539>

914 Schenzel, K., Fischer, S., & Brendler, E. (2005). New Method for Determining the  
915 Degree of Cellulose I Crystallinity by Means of FT Raman Spectroscopy.  
916 *Cellulose*, 12(3), 223–231. <https://doi.org/10.1007/s10570-004-3885-6>

917 Singh, R., & Geetanjali. (2013). Nutraceuticals: Promising Health Product.  
918 *International Research Journal of Medical Sciences*, 1(1), 14–17. Retrieved from  
919 <http://www.isca.in>

920 Toschi, T. G., Cardenia, V., Bonaga, G., Mandrioli, M., & Rodriguez-estrada, M. T.  
921 (2014). Coffee Silverskin : Characterization , Possible Uses , and Safety Aspects.  
922 *Journal of Agricultural and Food Chemistry*, 62, 10836–10844.

923 Ulijasz, A. T., Cornilescu, G., von Stetten, D., Cornilescu, C., Velazquez Escobar, F.,  
924 Zhang, J., ... Vierstra, R. D. (2009). Cyanochromes Are Blue/Green Light  
925 Photoreversible Photoreceptors Defined by a Stable Double Cysteine Linkage to a  
926 Phycoviolobin-type Chromophore. *Journal of Biological Chemistry*, 284(43),

927 29757–29772. <https://doi.org/10.1074/jbc.M109.038513>

928 Xu, B. J., & Chang, S. K. C. (2007). A comparative study on phenolic profiles and  
929 antioxidant activities of legumes as affected by extraction solvents. *Journal of*  
930 *Food Science*, 72(2). <https://doi.org/10.1111/j.1750-3841.2006.00260.x>

931

932