

An ecological perspective for analysing rural depopulation and abandonment

Francisco Lloret^{1,2}  | Adrián Escudero³  | Joan Lloret^{4,5}  | Fernando Valladares⁶ 

¹Ecology Unit, Department of Biología Animal, Biología Vegetal i Ecologia, Edifici C, Universitat Autònoma Barcelona (UAB), Barcelona, Spain; ²CREAF, Edifici C, Universitat Autònoma Barcelona (UAB), Barcelona, Spain; ³Biodiversity and Conservation Area, Universidad Rey Juan Carlos, Móstoles, Spain; ⁴Instituto de Geociencias, CSIC-UCM, Madrid, Spain; ⁵Departamento de Geodinámica, Estratigrafía y Paleontología, Facultad de Ciencias Geológicas, Universidad Complutense, Madrid, Spain and ⁶Departamento Biogeografía y Cambio Global, Museo Nacional de Ciencias Naturales, (MNCN-CSIC), Madrid, Spain

Correspondence

Francisco Lloret

Email: francisco.lloret@uab.cat

Funding information

Agència de Gestió d'Ajuts Universitaris i de Recerca, Grant/Award Number: 2021 SGR 00849; Ministerio de Ciencia e Innovación, Grant/Award Number: PID2020-115264RB-I00 and PID2021-126927NB-I00; Fundación Biodiversidad

Handling Editor: Robert Fish

Abstract

1. Population loss in rural areas is rapidly increasing in high-income countries, raising concerns and debate, given its socio-economic consequences. Despite the evident environmental dimension of the phenomenon, ecological knowledge has been neglected in the analysis of actions aiming to reverse rural depopulation. Particularly, cultural landscapes reflect memories of ecological processes that have configured current patterns of biodiversity and ecosystem services.
2. Based on ecological principles, we present a conceptual procedure to assess the behaviour of social–ecological systems subjected to depopulation, projecting their expected trajectories through time within a framework defined jointly by demographic and environmental–ecological dimensions (in our case, biodiversity, carbon storage, pollution control, water resources and soil conservation).
3. We applied this procedure to various alternative interventions designed to confront depopulation in Spain: (1) non-intervention, (2) maintenance of the historical landscape configuration, (3) active conservation, (4) extensive, sustainable land use and (5) intensified land use.
4. We conclude that extensive, sustainable land use better optimizes criteria of demographic consolidation, environmental impact, resilience and implementation of actions confronting depopulation. We highlight the need to incorporate ecological knowledge into the assessment and application of actions confronting rural depopulation.

KEYWORDS

biodiversity, depopulation, ecosystem services, fireworks graphs, Spanish rural areas

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Authors. *People and Nature* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

1 | INTRODUCTION

The rural exodus and abandonment registered in many countries, with special intensity in high-income countries (Hospers & Reverda, 2015), has given rise to social concerns and strong tensions, due to the important economic, political, cultural, demographic and environmental consequences (Hospers & Reverda, 2015; Laborda Soriano et al., 2021). Reversing the depopulation of rural territories is a critical issue on political agendas, and it is a policy priority for the EU (ESPON, 2017). However, the economic, logistical and social incentives mobilized to stimulate the repopulation of these territories have not been successful so far (Navarro-Valverde et al., 2021), and, moreover, they may even exacerbate socio-economic and environmental problems (Hermoso et al., 2022). Significantly, these initiatives have been mainly aimed at promoting economic and social development, while neglecting the ecological dimension (Bruno et al., 2021; MacDonald et al., 2000; Martínez-Abraín et al., 2020; Regos et al., 2016), despite the current dramatic context of climatic crisis and biodiversity loss (IPBES, 2019; IPCC, 2021). In fact, ecological and social dimensions should not be separated when assessing rural depopulation (Hospers & Reverda, 2015). For instance, both social and ecological systems operate at spatial and temporal scales that go beyond the territory experiencing population decline. Actions to overcome the urban–rural dichotomy will benefit from considering depopulation and rural abandonment from an ecological perspective, reinforcing the conception of a common system in which rural and urban subsystems mutually nourish each other (Davoudi & Stead, 2002; Tacoli, 1998).

Ecological memory is essential to understand the ongoing ecosystem transformations in areas experiencing massive depopulation. This memory corresponds to the degree to which ecological processes are shaped by its history (Peterson, 2002) and, in our case, they are determined to a great extent by previous human activities (Balaguer et al., 2014) that have configured current landscapes, which, therefore, can be considered as cultural landscapes (Farina, 2000) (Figure 1). In these cultural landscapes subject to intervention, crops and human-driven open habitats prevail to the detriment of extended forests (Sabatini et al., 2017)—whose remnants have also largely been simplified in terms of their structure




FIGURE 1 Biodiversity decay associated with human activity occurring during the last millennia in cultural landscapes of Europe that have originated novel historical ecosystems and species assemblages in territories currently experiencing depopulation.

and biological composition (Lloret, Solé, et al., 2009)—and other habitats such as wetlands and natural grasslands. The historical legacy of human societies is particularly evident in European landscapes, especially in the Mediterranean region (Blondel et al., 2010). However, a 'primeval' situation of naturalness is frequently used as a reference value for range management and restoration (Aronson et al., 1993). But, the primeval ecosystems that were set in these landscapes cannot be recognized exactly in current landscapes, and any faithful restitution of non-modified ecosystems is elusive (Balaguer et al., 2014). Nevertheless, ecological processes that maintain biodiversity, primary productivity and carbon fixation, soil maintenance and the regulation of water and nutrient flows keep operating in these cultural landscapes, which, in fact, provide many of these essential services at an appreciable extent (Bernues et al., 2014; Gómez-Baggethun et al., 2011).

The history of cultural landscapes has determined the complex relationship between humans and biodiversity. A complex balance appears between, on the one hand, extinctions, impoverishment of food webs and disruption of some ecosystem functions (Root-Bernstein & Ladle, 2018). On the other hand, there is a maintenance of diverse and rich semi-natural habitats (Blondel & Aronson, 1999; Carrión et al., 2010), some of them with difficult access, but also in non-remote lands where low intensive agricultural and pasture uses, such as in 'dehesas'—that is, cultural grazed savanna-like ecosystems, common in the western Iberian Peninsula—(Ramírez-Hernández et al., 2014) are established. In these habitats, selective and evolutionary processes have operated in many lineages, thus permitting the persistence of many species and intricate biological interactions (Talle et al., 2016; Figure 1). These agroecosystems not subjected to intensification perform a set processes functionally mimicking natural ecosystems (Power, 2010), although a replacement of natural complex food webs controlled by large vertebrates or water and nutrient cycles cannot be achieved. For example, pressure from livestock has replaced at some extent wild herbivory selective forces, while extensive cultivation has substituted natural habitats affected by recurrent wildfires and herbivore foraging (Navarro et al., 2015). In consequence, a relevant part of the diversity of these cultural landscapes is confined to semi-natural habitats that require active human presence and active management (i.e., grazing, extensive agriculture; Estrada-Carmona et al., 2022; Maurer et al., 2022). When crops, grasslands and forests cease to be exploited, well-known ecological processes operate, largely resembling secondary succession (Escribano et al., 2014). This transformation of vegetation leads to new thickets, woodlands and forests, while open habitats such as grasslands and old fields diminish. A balance is established between the advent of conditions favouring biodiversity that is unable to life in human-driven habitats and the loss of conditions for biodiversity directly associated with these habitats. So, some species may benefit from the derived increase in semi-natural shrublands and forests following human depopulation (Escribano et al., 2014; Estrada-Carmona et al., 2022). This is the case of some components of biodiversity that have been

largely decimated (Martínez-Abrain et al., 2020) or can hardly survive in regularly altered landscapes, but may find an opportunity to recover when human intervention is limited (Hirt et al., 2021). In contrast, rural abandonment may bring plants and animals typical from agriculture and steppe-like habitats such as some specialized bird species (Traba & Morales, 2019), to the brink of extinction (Moreno Saiz et al., 2019). Moreover, legacy effects of past anthropic use often remain through time in depopulated areas; for instance, invasive species introduced by humans (Guillermé et al., 2020) and the absence of now-extinct key megaherbivores may impede the recovery of biodiversity assemblages, which would need top-down trophic regulations (Svenning et al., 2019).

Here, our main aim is to address rural depopulation from an ecological viewpoint, assessing the effect of different management alternatives that potentially induce 'depopulation reversal', while simultaneously considering their impacts on biodiversity and various ecosystem services. To do this, after establishing different alternatives of intervention in rural depopulated areas, we present a simple conceptual procedure to assess, on the one hand, changes in human demography and, on the other hand, loss of biodiversity and ecosystem services. This procedure aims to provide a roadmap for the assessment of demographic-environmental systems. We qualitatively exemplify our assessment in Spanish territories suffering drastic rural depopulation, mainly in mountain areas (ca. 700–2000m AMSL; <https://atlasau.mitma.gob.es/#c=indicador&i=pobvar.pobvar010&view=map4>, accessed 8 June 2023, Atlas Digital de las Zonas Urbanas, Ministerio de Transportes, Movilidad y Agenda Urbana), where a lot of studies are addressing the topic (e.g. Caceres-Feria et al., 2021; Larráz & San-Martin, 2021; Regos et al., 2016; Rodríguez-Rodríguez et al., 2021) and the phenomenon is gaining space on the political agenda. Specifically, our objectives are to (i) build an operational framework for assessing massive rural depopulation in a given territory incorporating an ecological perspective; (ii) establish a methodology to evaluate different intervention alternatives; (iii) analyse the consequences of some of these alternatives on the interface defined by human population (demographic) and different environmental dimensions; and (iv) provide recommendations for intervention and planning in such depopulated territories.

2 | METHODS

2.1 | Assessing alternative interventions to confront rural depopulation

Our procedure is based on the analysis of temporal trajectories of a social-ecological system under different management alternatives in a two-dimensional space defined by (i) a human demographic dimension and (ii) an environmental dimension characterizing ecosystems in such depopulated territory. Here, the demographic dimension considers two variables: mean population density and the inequality Gini index of the number of people in inhabited nuclei (farmhouses, hamlets, villages and towns), which informs on the spatial

patterns of human occupancy in the territory. Thus, a trend towards high inequality in the size of populations in the inhabited nuclei of the depopulated territory under consideration would indicate an increase in the people living in cities or mid-size towns, coupled with a decrease in non-urban land. In turn, the environmental dimension is adequately addressed by ecosystem services, when considering social-ecological systems.

The procedural steps are as follows:

1. An evaluation of historical changes in the selected demographic parameters in the depopulated territory, in our case for the ca. 1900–2000 period. Next, the evolution of these demographic parameters is projected from the present to the future (until ca. 2100 in our case, a period which approximately encompasses three human generations) in response to different alternative interventions to confront depopulation (see below section '2.3. Alternative interventions to confronting rural depopulation' for the analysed Spanish case; Figure 2).
2. An evaluation of the trends of critical environmental parameters under the above-mentioned alternatives, considering the same time frame used to estimate the evolution of demographic parameters (see above).
3. A projection of the temporal trajectory of the social-ecological system into the future for each of the alternatives. This trajectory




FIGURE 2 Demographic trends (density and inequality estimated by the Gini index of inhabited nuclei) in Spanish mountainous areas subject to rural depopulation. The Gini index is a measure of the inequality of the population distribution in the territory, with values ranging from 0 to 1, with 0 indicating perfect equality—all nuclei have the same population—and 1 maximal inequality, with a single nucleus houses the entire population. Different vectors represent expected future trends under different alternatives for confronting rural depopulation, from now (dot) to the end of the 21st century, following changes in the recent past (black line) (see main text for a description of the alternatives). Source for historical records: Goerlich et al., 2015.

corresponds to a vector whose components are the variation in the demographic parameters (e.g. population density, Gini index of population in inhabited nuclei) and the change in the environmental dimension (e.g. biodiversity, carbon storage, pollution, water resources and soil erosion) over a given time. In the case of the demographic parameter, we consider the difference between a future reference date (e.g. the year 2100 in our case) and the present. The component of the trajectory vector corresponding to the environmental dimension is an estimate of the effect that each alternative produces on the respective environmental dimension for the same period. This estimate should ideally be obtained from quantitative models, but, as in our case, it would often correspond to semi-quantitative trends supported by empirical studies undertaken in the region, or in areas with similar environmental and socio-economic contexts. To build the trajectories, it is essential to establish the territorial range in which the impacts of the proposed alternatives and the depopulation process occur. The result is a set of vectors that represent the foreseeable simultaneous change in demographic and environmental parameters. Here, we propose to name the resulting representation as 'fireworks graph', and it allows us to compare the vectors that describe the effects of the different alternatives (Figure 3).

2.2 | Implementation of the demographic–environmental space: The Spanish case

We illustrate the application of the proposed procedure for the assessment of rural depopulation in Spain, based in the published literature; so, the study does not directly involve human data or participants. We built the fireworks graphs as detailed above after considering published results addressing the relationship between depopulation and ecosystem functioning (related to ecosystem services) and biodiversity in inland and mountainous rural areas of Spain. For this purpose, we searched the literature in the Web of Science database using the keywords string 'depopulation AND (biodiversity OR carbon OR pollution OR water OR soil OR erosion OR ecosystem service) AND Spain' for title, abstract and keywords, and we incorporated additional publications from the author's records (see Data sources).

We considered two demographic parameters: mean population density and the inequality Gini index of the number of people in inhabited nuclei. In the Spanish case, the population density was low in the second half of the 19th century and early 20th century in many rural areas, with a slight-to-moderate tendency to increase, in contrast with a steeper increase in urban localities. Later on, rural areas experienced a slow population decline (e.g. Infante-Amate et al., 2016), which accelerated in the 1950s and, more particularly, the 1970–1980s (source: 'Instituto Nacional de Estadística' of Spain), a trend maintained to the present (Franch Auladell et al., 2013). Then, we qualitatively projected the trends of the trajectories of the two demographic parameters considered for the five proposed alternatives (see below section '2.3.

Alternative interventions to confronting rural depopulation') until 2100.

We considered the following environmental surrogates for our assessment: biodiversity, above-ground carbon storage, pollution, water resources and soil conservation. They are linked to well-recognized supportive (biodiversity, soil conservation and absence of pollution), regulatory (carbon storage and water resources) and provisioning ecosystem services (water resources).

The selected environmental surrogates are deliberately wide. For instance, we consider biodiversity in a broad sense, without distinguishing its organizational levels—genetic, species, habitats and ecosystems—and also including biotic interactions and species functionality. This broad estimation of these environmental dimensions allowed us an illustration of the procedure without quantitative analyses, which would require modelling of the future trajectories of demographic and environmental parameters for a given territory, in accordance with the different alternative to face rural depopulation. Although a relationship between some of these dimensions is to be expected—for example, biodiversity is likely affected by the availability of water resources, soil conservation and pollution—we tried to avoid any highly covarying environmental dimension, such as invasions by exotic species—which are closely linked to biodiversity—or fire management—which is closely related to carbon storage.


2.3 | Alternative interventions to confronting rural depopulation

We consider five broad management approaches to facing depopulation and evaluate their impact on demographic and ecological dimensions. These approaches are assessed in virtual territories broadly corresponding to a county (ca. 500–1000 km²) in mountainous areas of Spain (ca. 700–2000 m AMSL). They are not applied to specific locations because our aim is to provide a general picture of the phenomenon by integrating the available information for the region. These five alternatives do not cover the entire range of management options, but they represent contrasting and realistic strategies of intervention for depopulated territories. Note also that they are not exclusive and can be applied to different areas in the same territory. Furthermore, specific actions may correspond to more than one alternative.

2.3.1 | Alternative 1: Crossed-arms

This alternative corresponds to non-intervention, that is, it maintains the current actions oriented towards population, economic activity and biodiversity conservation and environmental quality. The capacity of the social–ecological system to organize itself in the absence of planned intervention is implicitly recognized. This alternative implies the absence of a sound strategy for the promotion of human populations, which are expected to maintain the current trend of abandonment (Figure 2). So, secondary succession associated with

FIGURE 3 Expected temporal (21st century) trends in taxonomic biodiversity (a, b), carbon storage (c, d), pollution (e, f), water resources (g, h), and soil conservation (i, j), as well as two demographic parameters—population density (a, c, e, g, i) and inequality of population size in inhabited nuclei (b, d, e, h, j)—under different alternatives for confronting rural depopulation (AC, active conservation; CA, crossed-arms; EU, extensive-use; FL, frozen landscapes; IN, intensified-use). Dashed line represents human population demographic trajectories in the recent past in the corresponding biplot.



the abandonment of crops, pastures and forests constitutes the dominant ecological process. This implies a predominance of encroachment and afforestation, which lead to the shrinking of open

and semi-natural habitats such as pastures and *dehesas* and their associated biodiversity (Ramírez-Hernández et al., 2014), but, on the other hand, the appearance of habitat for some animals (f.e. birds,

Regos et al., 2016), and an increase in C sequestration from the atmosphere (Velazquez et al., 2022). This alternative does not require any great additional financial or infrastructure investments, and economic activity is also expected to maintain current declines.

2.3.2 | Alternative 2: Frozen landscape

This alternative seeks the maintenance of the historical landscape configuration, with well-conserved remnants of natural habitats and mosaics of croplands and pastures. In Europe, this alternative is exemplified by the actions supported by the European Habitats Directive (1992), which provides a framework for biodiversity conservation by recognizing the value of historical habitats, many of them semi-natural and without trees. This alternative adopts a static view of ecological systems, in which vegetation units become references overlooking the role of secondary succession and other constituents of biodiversity and ecosystem dynamics. This static perspective has been questioned, since it does not consider population dynamics and community assembly processes, particularly responding to climate change (Normand et al., 2007). Actions that would potentially revert the trend towards depopulation are subordinated to inputs into the territory aimed at maintaining its spatial configuration, since historical landscapes depend on human activities and management (Figure 2). Promotion of economic activity derives from management plans for habitats, from agro-environmental measures implemented in cultivated areas to reconcile production, sustainability and the conservation of such habitats (Halada et al., 2011), and from ecotourism initiatives, which take advantage of traditional uses and the conservation of valuable semi-natural habitats.

2.3.3 | Alternative 3: Active conservation

This alternative is based on the premise that rural abandonment is an opportunity for biodiversity conservation and the recovery of healthy ecosystems. Therefore, actions promoting biodiversity and ecosystem health—for example, measures for the maintenance, restoration and recreation of habitats; species and habitat monitoring; reintroduction, translocation and conservation management of species populations; control of invasive species (Kati et al., 2015)—are enhanced, particularly in protected areas, such as the Natura 2000 network in Europe. Actions aiming to revert depopulation are accessory, given the secondary role of the human population for biodiversity maintenance, as it does not need to remain in the territory and may even be encouraged to abandon it. Actions derived from active conservation could favour the maintenance of even an increase of local human population (Schou et al., 2021), and they are expected to produce benefits to human societies as well, for instance by preventing diseases; but some evidence suggests negative effects on local populations in strongly protected areas (Rodríguez-Rodríguez et al., 2021; Schou et al., 2021; Figure 2). In this alternative, economic

activity is minimized and related to biodiversity-focused revenues, including ecotourism.

2.3.4 | Alternative 4: Extensive use of the territory

This alternative foments non-intensive land use. Biodiversity and environmental quality therefore benefit on a secondary level, although the maintenance of anthropogenic activity in large parts of a territory implies a certain environmental impact. This alternative likely helps stop or even reverse rural depopulation (Caceres-Feria et al., 2021; Figure 2), but it requires precise planning and a significant investment in public services and infrastructures for communication, while still valuing sustainability. The technological improvements achieved in telecommunications, together with the rise in teleworking, can make it easier to retain and attract people. The portfolio of economic activities for this alternative is significantly represented by ecological and regenerative agriculture—for example, pastoralism and sustainable agriculture based on the recovery of local plants' genetic resources and the maintenance of a heterogeneous agriculture-forest mosaic (Marull et al., 2014)—and other productive, sustainable activities with likely low environmental impacts, such as ecotourism, in combination with outdoor activities and others associated with the conservation of biodiversity and the improvement of ecosystem services (Caceres-Feria et al., 2021).

2.3.5 | Alternative 5: Intensified use of the territory

Rural depopulation provides an opportunity for activities that are not easily compatible with high population densities. This intensification alternative includes the installation of large power generation complexes, polluting industries, urban disposal, and industrial and mineral waste facilities, as well as the construction of transport infrastructure and reservoirs. Intensified land use may also be derived from large recreational resorts (Martin et al., 2018), large mono-specific tree plantations, and the intensification of agriculture and farming (Gaglio et al., 2020), via, for example, large-scale irrigation projects. One of the main arguments in favour of this alternative is that economic activities with a high financial return eventually have a positive economic impact on local populations, while other arguments refer to global needs such as the mitigation of climate change. However, local environmental priorities are obscured, the effects on the ecosystems tend to be neglected, and the objectives and actions for biodiversity conservation are disregarded. Depopulation reversal dependent on economic activity would occur without specific actions. So, this alternative can help curb depopulation to a certain extent (Figure 2), but this effect is often only local and short-term (Collantes & Pinilla, 2004)—for example, while infrastructures are being built. The effect may disappear in subsequent phases of exploitation, when the demand for local workers declines and the remaining economic activity may not be sufficient to compensate for

TABLE 1 Qualitative assessment of alternatives for confronting rural depopulation according to several criteria: (i) effects on biodiversity and environmental quality, depopulation reversal and economic revenues, (ii) feasibility of implementation, in terms of economic investment, governance and set-up time, and (iii) capacity to promote the resilience of the social-ecological system (see text for further details). Colours indicate favourable (green), intermediate (yellow), or unfavourable (red) appraisal.

Alternatives to confront rural depopulation	Effect		Implementation			Resilience	
	Biodiversity environmental quality	Depopulation reversal	Economic revenue	Economic investment	Governance	Time to set-up	
Crossed arms (CA)	Subordinated to conserved landscape	Business as usual	Business as usual	Low	Easy	Fast	Low
Frozen landscape (FL)	Static perspective	Subordinated to conserved landscape	Environmentally friendly	Medium	Medium	Medium	Medium
Active conservation (AC)	Strong prioritization	Accessory to conservation	Minimized, conservation focused	High	Difficult	Slow	Medium
Extensive use (EU)	Subsidiary of land use	Promoted to maintain land use	Sustainability focused	Medium	Medium	Medium	High
Intensified use (IN)	Disregarded	Dependent on economic activity	Strong prioritization	High	Difficult	Medium	Low

the depopulation associated with the loss of agricultural land (Larraz & San-Martin, 2021), while economic revenues from other activities, such as tourism, could increase (Silvestre & Clar, 2010). In this alternative, economic revenues become the priority, favouring a broad range of potential activities, as described above. Overall, intensification tends to consume large amounts of resources and land, and often pollutes watersheds, air and soils (Ladrera et al., 2019).

2.4 | Assessment of the alternatives for confronting rural depopulation

We have assessed the proposed five alternatives according to these criteria: (i) feasibility of the implementation of the alternative, which can, in turn, be assessed according to the economic investment needed (e.g. cost of health service in territories with low density), the estimated economic returns (e.g. revenues from irrigation plans or energy infrastructures), the governance required for implementation (e.g. approval of specific regulations) and the expected time needed to set all these up, (ii) effectiveness in reversing depopulation (i.e. demographic consolidation), (iii) environmental impact on biodiversity and ecosystem services, (iv) resilience, considered as the system's ability to absorb the impact of disturbances and stressors while maintaining its functionality (Folke et al., 2010; Walker et al., 2004). The assessments of the reversal of depopulation and the environmental impact are derived from the patterns observed in our fireworks graphs, as discussed above, while economic revenue, implementation feasibility, and resilience are estimated from the literature (see Data sources) as shown in the Results for each biodiversity and environmental dimension (Table 1).

3 | RESULTS

3.1 | Biodiversity

The alternative with the most positive effect on biodiversity in the Iberian Peninsula is expected to be active conservation (Figure 3a), given the long history of overgrazing, expansion of crops and forest thinning that have had a negative effect on some key constituents of biodiversity, such as large mammals (Martínez-Abraín et al., 2020). Overall, a general trend of biodiversity loss is likely to be found with increasing human population density (Underwood et al., 2009). When combining biodiversity and human demographic trends, this alternative is likely to lead to a certain demographic decline since it does not involve active social policies aimed at reversing depopulation. This demographic decline would, in fact, provide opportunities for the conservation of certain species (Hinojosa et al., 2018) and habitats (Regos et al., 2016) that are vulnerable to human pressures and disturbances (Henle et al., 2008). The extensive-use alternative, which shares elements of traditional land use, would also favour biodiversity to some extent by maintaining and promoting activities compatible with the persistence of multiple habitats and

critical resources for some species (Beckmann et al., 2018) in certain landscape configurations (Franz et al., 2011; Varga, 2020; but see Lopez & Pardo, 2018 for the negative impacts on biodiversity of ecotourism and outdoor activities). At the same time, this alternative would promote economic activity favouring the maintenance of the human population, and even its increase (Caceres-Feria et al., 2021; Hospers & Reverda, 2015).

The frozen landscape alternative would have a less beneficial effect on biodiversity. Although human pressure has diminished in depopulated areas during the late 20th century, these biodiversity constituents seem unable to recover the same levels exhibited in distant historical times (Blondel et al., 2010; Navarro et al., 2015). It is worth noting that, although this alternative appears to ensure the status of certain habitats and species, it lacks any integrated vision of their dynamics at the landscape level. Furthermore, this frozen landscape would not be expected to substantially reverse any prevailing depopulation, given the limited impact on the human population of actions devoted to maintaining the landscape.

The crossed-arms alternative, in its turn, would maintain the current situation of population collapse, together with a certain deterioration in biodiversity, due to the loss of semi-natural open habitats by encroachment and afforestation (Halada et al., 2011). However, an incorporation of species inhabiting mature successional stages and vertebrates sensitive to anthropogenic pressure is also expected (Hirt et al., 2021; Regos et al., 2016). Finally, the intensified-use alternative produces the greatest negative impact on biodiversity (Beckmann et al., 2018; Carmona et al., 2020; Franz et al., 2011) by affecting habitats and altering the quantity and quality of water resources and soils, while it can counteract demographic decline, at least temporarily.

The component of the trajectories associated with the impact on biodiversity conservation would not vary substantially when comprehensively considering the inequality of the size of inhabited nuclei size (see fireworks graphs in Figure 3). However, the trajectories of the alternatives of extensive-use and active conservation would differ, since the former would promote a greater number of inhabited nuclei of intermediate size, and therefore, a greater inequality in the distribution of the population in the territory, than the latter, in which small nuclei are likely to disappear (Figure 3b). Similarly, the crossed-arms and frozen landscape alternatives would maintain the current population dilution and concentration in larger nuclei sometimes out of the territory, while the intensified-use alternative could retain some population in nuclei of intermediate size, through work in infrastructures, industries and new facilities. These demographic trends remain constant throughout the analysis of other environmental dimensions, and so their results will not be repeated further.

The trajectories of the different alternatives could be modified when the loss of population in the territory implies a greater deterioration of biodiversity, due, for example, to the loss of certain habitats. In that case, the extensive-use alternative would not be sufficient to reverse the loss of diversity, although it could alleviate it. This effect would be counterbalanced by biodiversity improvements driven by depopulation.

3.2 | Carbon storage

The carbon storage of Mediterranean-type ecosystems mainly relies on above- and underground woody vegetation (Vayreda et al., 2012), although the role of organic matter in grasslands is far from negligible (Doblas-Miranda et al., 2013). Therefore, forest cover and age constitute a proxy for this service. Overall, an increase in carbon storage is expected with the crossed-arms alternative (Figure 3c,d) since it reproduces recent trends of afforestation and carbon sequestration following land abandonment in Europe (Fuchs et al., 2016). However, carbon storage can be challenged by raising wildfires due to the increase of fuel and climatic risk (Duane et al., 2021). An increase in carbon storage is also to be expected with the frozen landscape and active conservation alternatives, as they favour the regeneration and growth of forests. This increase would be lower than in the crossed-arms alternative, since the maintenance of forests would be counterbalanced by the conservation of open habitats, which in the case of active conservation would be a crucial objective as they host some specialized species (Varga, 2020). Similarly, carbon storage is not expected to increase so substantially under the extensive-use alternative, due to the enhancement of open habitats for extensive livestock grazing; but no clear drop can be expected either, thanks to the conservation of forest remnants in the landscape. Finally, the destruction of forested and mountain grasslands by specific interventions associated with the intensified-use alternative would diminish carbon stocks (Doblas-Miranda et al., 2013).

3.3 | Pollution

Pollution in depopulated rural areas is expected to diminish due to the current reduction in the population and its associated economic activity (Bruno et al., 2021) and to the ability of ecosystems, particularly aquatic ones, to recover from both concentrated and diffuse sources of pollution (Donohue et al., 2010). This environmental improvement would reach a peak with active conservation and, to a lesser extent, with the frozen landscape alternative, due to its higher remaining activity, compared with the rest (Figure 3e,f). In contrast, intensification is expected to increase pollution in soils and water due to intensive farming (Ladrera et al., 2019), mining (García-Carmona et al., 2019), recreational resorts and installations for energy production (Heal et al., 2020). In fact, depopulated territories are often sought as sites for the disposal of hazardous products, such as industrial or radioactive wastes (e.g. Llurdes et al., 2003), representing intensive use of the territory. Human activity derived from extensive use is also expected to produce some levels of pollution, although much less than in this intensified-use alternative.

3.4 | Water resources

Local water input into the social-ecological system is directly determined by climate, aquifers and vegetation conditions. Soil quality

and vegetation cover also play essential roles in regulating water supply. Water demand, in its turn, is determined by human population and economic activity. Water requirement is not constant over time, so water supply must be ensured in periods of peak demand or low availability—that is, drought periods. Therefore, depopulation and reduced economic activity will decrease the demand for water, while also enhancing water regulation by vegetation and soils (Pisabarro et al., 2019). Accordingly, those alternatives (crossed-arms, frozen landscapes, active conservation) involving low water demand derived from low levels of population density and reduced industry and agricultural intensification would, together with some degree of vegetation and soil preservation, increase both water resources and their regulation (Bruno et al., 2021). This increase would be proportional to the expected recovery of vegetation. This trend can be counterbalanced to some extent by the higher water requirements of forested lands, although this effect is limited in comparison with the impact of climate change (Otero et al., 2011). Overall, active conservation would represent the lowest water demand, while maintaining higher water stocks and groundwater recharge, as supported by the role of protected areas in providing regulating services (Castro et al., 2015). The crossed-arms and frozen landscapes alternatives would also maintain a positive balance of water resources to some extent (Figure 3g,h). In contrast, the intensified-use alternative would reduce water resources, due to high demand (e.g. for irrigation, Vidal-Legaz et al., 2013) and a likely reduction in vegetation and soil preservation. This reduction in water resources is expected to be less marked in the extensive-use alternative, due to lower demand for water and better preservation of vegetation and soils at landscape level (Roper et al., 2019).

3.5 | Soil conservation

Soil erosion has been exacerbated by past deforestation, overgrazing and recurring fires associated with pastoralism, particularly in Mediterranean mountain areas (Sanjuan et al., 2016), despite the retention of soil provided by traditional terraces (García-Ruiz, 2010). So, depopulation tends to improve soil retention and development (Beguiría et al., 2003; Bruno et al., 2021), which, in turn, produce cascade effects by enhancing revegetation processes (Navas et al., 2008) and reducing off-site costs (e.g. quality of surface and ground water; Colombo et al., 2005). However, the abandonment or destruction of terraces on croplands would endanger soil, reducing any such improvement in the crossed-arms alternative (García-Ruiz, 2010; MacDonald et al., 2000), particularly with an increased frequency and intensity of wildfires (Cerdà & Lasanta, 2005). The risk of soil loss would be less pronounced in the frozen landscape and, to a lesser extent, extensive-use alternatives, as vegetation cover would be enhanced, and terraces could be somewhat better preserved. The most effective alternative for soil conservation will be active conservation, as it would increase the vegetated area; it would be even more effective if soils were specifically included as a target for conservation and

restoration, particularly in drylands (Bautista et al., 2010). In contrast, soil conservation is put at risk in the intensified-use alternative, particularly when mining or agricultural intensification are involved (Bruno et al., 2021; Cerdà et al., 2012). In contrast, soil conservation could benefit in the extensive-use alternative if some of the soils affected by intensive land use revert to a non-intensive use. This shift would allow some soils to return to semi-natural conditions (Figure 3i).

3.6 | Assessment of the alternatives for confronting rural depopulation

Overall, our qualitative assessment of the alternatives reveals that the most favourable alternative is extensive-use, according to their capacity to reverse depopulation and maintain biodiversity and environmental quality, together with the feasibility of their implementation and their ability to promote resilience of the social-ecological system (Table 1). In this alternative, none of the criteria show low levels of achievement and the best demographic consolidation and resilience are found here. In contrast, the alternatives that obtain the worst rates are intensified-use and active conservation, with five criteria rated negatively (high environmental impact, low depopulation reversal, high economic investment, difficult governance, and low resilience in the intensified-use alternative; and weak demographic consolidation, low economic revenue after substantial economic investment, difficult governance, and long set-up time in the active conservation alternative). This assessment is supported by studies comparing the outputs of different land uses related to our alternatives in Spain (Bruno et al., 2021; Castro et al., 2015; Franz et al., 2011; Varga, 2020; Vidal-Legaz et al., 2013), and the assessments of biodiversity and ecosystem services shown above.

4 | DISCUSSION

4.1 | Integrating ecological and social dimensions into depopulation assessment

In the context of rural depopulation, social-ecological dynamics are historically determined by both natural and social processes, as reflected in cultural landscapes. This rationale implies that: (i) there are ecological legacies that must be interpreted and considered when planning and applying decisions affecting depopulation; (ii) previous historical situations do not provide exact references suitable for replication (Balaguer et al., 2014) but they provide valuable information on key ecological processes; and (iii) in a given historical context, management should aim to favour the resilience of biodiversity and ecosystem services (Connell & Ghedini, 2015).

One important issue when integrating social and ecological processes is the difference in the time and spatial scales at which they operate. For instance, Gosselin and Callois (2021) have found that

current patterns of biodiversity in terms of proportion of extinct species show, at the European national level, a time lag of approximately one century compared to human population density—extinction debts. Moreover, the abandonment of agricultural practices results in a secondary succession of vegetation, which takes decades to produce an increase in the forest surface (Duane et al., 2021), and even longer to culminate in highly structured stands (Rozas et al., 2009) and balance carbon budgets with a significant contribution of soil organic matter (Fuchs et al., 2016). Also, regulation of water quality shows important lags between production and consumption, with substantial spatial mismatching (Fremier et al., 2013). Therefore, according with these studies, a general recommendation is that short-term economic benefits should be contrasted with environmental gains, which often only emerge in the medium and long term.

Consequently, contextual dependencies need some kind of spatial–temporal up-scaling. Actions and demands in a given environment have an impact on other parts of a territory, and on other components of the social–ecological system (Bruno et al., 2021; Caceres-Feria et al., 2021; Costa et al., 2017; Davoudi & Stead, 2002). This applies to environmental impacts such as pollution, as well as the demands of urban populations for agricultural, livestock and forest resources, and recreational use, as particularly reflected in the intensified-use alternative and, to a lesser extent, the extensive-use, frozen landscape and crossed-arms ones. Furthermore, rural populations also present a demand for technological resources created in urban areas. Therefore, any solutions for facing depopulation and integrating ecological conditions should optimize the cost–benefit balance at different local, regional, and global scales. It is thus important to apply the principle that rural and urban systems are part of a functional and ecological continuum (Tacoli, 1998). They are mutually dependent in terms of flows of energy, matter, goods, information, and people, as well as in terms of social and ecosystem services. Although this principle has been recognized by ecologists (Villarreal Walker & Beck, 2012), social perception and political practitioners often confront territorial and environmental challenges separately, thereby obstructing any overall assessment or application of effective solutions.

Dealing with these conflicts implies that we must (i) prioritize those actions with the greatest synergies and evaluate existing trade-offs, and (ii) consider different spatial and temporal scales in order to optimize the distribution of actions in space and time. Note that some actions, such as ‘rewilding’—which aims to re-establish integrated ecological processes by improving homeostasis in ecosystem functioning (Svenning et al., 2019), by re-establishing key elements such as locally extinct herbivores or carnivores—shares objectives and procedures that we can recognize in more than one alternative (in our case, extensive-use and active conservation), albeit with different degrees of involvement. We have knowledge and tools (cost–benefit analysis, multi-criteria, modelling) that provide us with a well-founded and rational approach to these evaluations (e.g. Martínez-Sastre et al., 2017). The participation of the population in decision-making is essential here, and it must be sustained by the dissemination and understanding of scientific knowledge. In Spain,

strong social conflicts are expected to appear with the intensified-use and active conservation alternatives, resulting in challenges to governance. While intensification will trigger environmentalist reactions (Costa et al., 2017), active conservation will face opposition from local populations seeking to maintain their control over land resources (Hinojosa et al., 2018). These conflicts become ambivalent when the proposed infrastructures in the intensified-use alternative promote local economic activity, particularly when the latter is apparently consistent with the historical economic activity or provides environmental benefits on a broader scale (e.g. renewable energy). In turn, any implementation of active conservation should be slow, in keeping with the pace of not only ecological processes but also administrative obstacles (Perez-Barberia et al., 2023), and social acceptance, which can vary from refusal to acceptance, such as in programmes involving large carnivores (Delibes-Mateos et al., 2022; Recio et al., 2020, respectively).

The integrated maintenance over time of ecological and social processes is well-exemplified by resilience (Folke et al., 2010; Walker et al., 2004). Here, this is expected to be prominent in the extensive-use alternative, thanks to a diffuse spatial distribution of human activities that permits the functioning of essential self-regulatory ecological mechanisms (i.e. those involving biodiversity and soil and watershed integrity). This implies, for instance, that decision makers confronting depopulation should favour diverse landscapes (i.e. hosting grasslands, woodland and forests) with high species diversity as a means to buffer the impacts of pests, extreme weather events and wildfires (Fischer et al., 2006; Varga, 2020). In contrast, alternatives such as active conservation and frozen landscape, which tend to dissociate human activities from cultural landscapes, can threaten some components of the ecosystems which have historically been modelled by human activity (i.e. diverse mountain pastures, or domesticated lineages of animals and plants). For instance, an effective fire suppression policy aimed at preserving forests can also contribute to an increase in fuel load and may favour large, highly intense wildfires that, in turn, reduce the resilience of burned forests (Lloret, Piñol, et al., 2009). The maintenance—or even promotion—of ecological self-regulating processes, such as herbivory by large mammals, could counterbalance any loss of the regulatory capacity associated with human activity. However, when this capacity is completely lost, as tends to occur in the crossed-arms and intensified-use alternatives, resilience is expected to be dramatically depleted. Current trends of rural abandonment, corresponding to the crossed-arms alternative, are also increasing the fuel load, which—combined with increasing climatic risk—is triggering wildfires with higher intensities and frequency (Duane et al., 2021), jeopardizing important ecosystem services, such as soil conservation. In the intensified-use alternative, the destruction of habitats and disruption of ecological processes are particularly liable to reduce resilience (Laliberté et al., 2010). For instance, soil losses due to such intensification dramatically deplete the capacity of vegetation to establish itself (Colombo et al., 2005). This effect is even more obvious when exploitative activities constitute harmful disturbances in themselves or amplify a pre-existing natural disturbance regime (e.g. García-Carmona et al., 2019).

4.2 | Fire-works procedure for an integrative depopulation assessment

Given its multidimensional conception, the proposed procedure based on the evaluation of land management alternatives and their impact on human demographic and ecological trends is useful for an integrative semi-quantitative assessment of rural depopulation, despite reservations about a lack of precise, specific information. Interestingly, the procedure can be applied to situations other than the Spanish case, corresponding to different territorial, socio-economic or historical situations addressing the environmental consequences of depopulation. In fact, it can become more comprehensive if it incorporates additional operations, by (i) combining different environmental dimensions, for example, by weighting and multiplying their respective vector components; (ii) incorporating alternatives other than those exposed here for the Spanish case, once their effects on demography and ecological dimensions have been evaluated; (iii) combining the consequences of different alternatives operating in distinct parts of a given territory, for example, by weighting the coordinates of the vectors that describe the respective trajectories for the proportion of the territory affected by each alternative; (iv) analysing trends at different future time lags, that is, these intervals may represent different periods marked by changes in the relative weights of different alternatives; and (v) estimating lagged effects and other non-linear responses of demographic parameters and environmental dimensions.

Our procedure can also evaluate other properties of interest for the social-ecological system. For instance, resilience would correspond to the distance of the final coordinate of the vector on the axis of the environmental dimension, with respect to a historical reference without rural depopulation. The overall magnitude of the transformation of the social-ecological system under the different scenarios (response intensity or responsiveness) corresponds to the angle or slope of the vector, with respect to the axes that describe the demographic and environmental dimensions. Finally, the directionality of the change corresponds to the maintenance of the direction of the vectors in a given alternative over successive periods of time.

5 | CONCLUDING REMARKS

Planning and actions carried on to confronting rural depopulation must integrate the ecological perspective. In the current context of biodiversity loss, climate change and environmental degradation, rural depopulation has clearly acquired a dimension that goes beyond a merely socio-economic and cultural analysis. Land uses in depopulated territories range from promoting biodiversity conservation, which provides opportunities to confront the global environmental crisis and to subsequently implement mitigating measures, to intensification of land use. In addition to the specifics of the physical environment and the current social situation, the historical legacies

determining the spatial-temporal contexts of each locality are crucial to any understanding of the patterns of change in ecosystem services and biodiversity in all their complexity. Furthermore, the current crisis of globalization and climate change, and the value being placed on the use of local natural resources and environmental health, are leading to a reassessment of the opportunities provided by these territories.

We provide a procedure for an integrated analysis of demographic trends and ecological properties and processes of the social-ecological systems, particularly those resulting in biodiversity conservation and ecosystem services. Our evaluation of these alternatives in Spain indicates that management based on the extensive sustainable use of natural resources—including activities related to recreation and biodiversity conservation, as well as the provision of ecosystem services—would be the option that best fulfils the criteria for demographic consolidation, lower environmental impact, increasing resilience, and feasible implementation. More detailed analyses should be carried out, based on local characteristics and a specific assessment of the performance of demographic and environmental parameters in terms of realistic alternative approaches. Crucially, any depopulation analysis should be based on a set of basic ecological principles that must be included in the implementation of any mitigating actions.

AUTHOR CONTRIBUTIONS

Francisco Lloret, Fernando Valladares, Joan Lloret, and Adrian Escudero conceived the ideas and designed the methodology; Francisco Lloret, Fernando Valladares, Joan Lloret, and Adrian Escudero implemented the proposed methodology in the Spanish study case from published literature and discussed the interpretation; Francisco Lloret led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

ACKNOWLEDGEMENTS

This study has been supported by the Spanish Ministry of Science and Innovation (grants PID2020-115264RB-I00, PID2021-126927NB-I00), AGAUR (Government of Catalonia) (grant 2021 SGR 00849), and Fundación Biodiversidad (Proyecto BioFoRest).

CONFLICT OF INTEREST STATEMENT

The authors have no potential sources of conflict of interest to disclose, as established by the policy of the journal 'People and Nature'.

DATA AVAILABILITY STATEMENT

The article is bibliographic, based on the published literature and the study has not generated data eligible to be stored in a public archive.

ORCID

Francisco Lloret  <https://orcid.org/0000-0002-9836-4069>

Adrián Escudero  <https://orcid.org/0000-0002-1427-5465>

Joan Lloret  <https://orcid.org/0000-0002-9070-4639>

Fernando Valladares  <https://orcid.org/0000-0002-5374-4682>

REFERENCES

- Aronson, J., Floret, C., LeFloc'h, E., Ovalle, C., & Pontanier, R. (1993). Restoration and rehabilitation of degraded ecosystems in arid and semi-arid lands. I a view from the south. *Restoration Ecology*, 1, 8–17.
- Balaguer, L., Escudero, A., Martín-Duque, J., Mola, I., & Aronson, J. (2014). The historical reference in restoration ecology: Re-defining a cornerstone concept. *Biological Conservation*, 176, 12–20. <https://doi.org/10.1016/j.biocon.2014.05.007>
- Bautista, S., Orr, B. J., Alloza, J. A., & Vallejo, R. V. (2010). Evaluating the restoration of dryland ecosystems in the northern Mediterranean. In G. Schneier-Madanes & M. F. Courel (Eds.), *Water and sustainability in arid regions*. Springer. https://doi.org/10.1007/978-90-481-2776-4_18
- Beckmann, M., Gerstner, K., Akin-Fajiyi, M., Ceausu, S., Kambach, S., Kinlock, N. L., Phillips, H. R. P., Verhagen, W., Gurevitch, J., Klotz, S., Newbold, T., Verburg, P. H., Winter, M., & Seppelt, R. (2018). Conventional land-use intensification reduces species richness and increases production: A global meta-analysis. *Global Change Biology*, 25, 1941–1956. <https://doi.org/10.1111/gcb.14606>
- Beguieria, S., López-Moreno, J. I., Lorente, A., Seeger, M., & García-Ruiz, J. M. (2003). Assessing the effects of climate oscillations and land-use changes on streamflow in the Central Spanish Pyrenees. *Ambio*, 32, 283–286. <https://doi.org/10.1579/0044-7447-32.4.283>
- Bernues, A., Rodríguez-Ortega, T., Ripoll-Bosch, R., & Alfnes, F. (2014). Socio-cultural and economic valuation of ecosystem services provided by Mediterranean mountain agroecosystems. *PLoS One*, 9, e102479. <https://doi.org/10.1371/journal.pone.0102479>
- Blondel, J., & Aronson, J. (1999). *Ecology and wildlife of the Mediterranean Region*. Oxford University Press.
- Blondel, J., Aronson, J., Bodiou, J.-Y., & Boeuf, G. (2010). *The Mediterranean region: Biological diversity in space and time* (2nd ed.). Oxford University Press.
- Bruno, D., Sorando, R., Álvarez-Farizo, B., Castellano, C., Céspedes, V., Gallardo, B., Jiménez, J. J., López, M. V., López-Flores, R., Moret-Fernández, D., Navarro, E., Picazo, F., Sevilla-Callejo, M., Tormo, J., Vidal-Macua, J. J., Nicolau, J. M., & Comín, F. A. (2021). Depopulation impacts on ecosystem services in Mediterranean rural areas. *Ecosystem Services*, 52, 101039. <https://doi.org/10.1016/j.ecoser.2021.101369>
- Caceres-Feria, R., Hernandez-Ramirez, M., & Ruiz-Ballesteros, E. (2021). Depopulation, community-based tourism, and community resilience in southwest Spain. *Journal of Rural Studies*, 88, 108–116. <https://doi.org/10.1016/j.rurstud.2021.10.008>
- Carmona, C. P., Guerrero, I., Peco, B., Morales, M. B., Oñate, J. J., Pärt, T., Tschardtke, T., Liira, J., Aavik, T., Emmerson, M., Berendse, F., Ceryngier, P., Bretagnolle, V., Weisser, W. W., & Bengtsson, J. (2020). Agriculture intensification reduces plant taxonomic and functional diversity across European arable systems. *Functional Ecology*, 34, 1448–1460.
- Carrion, J. S., Fernández, S., González-Sampériz, P., Gil-Romera, G., Badal, E. C.-M., López-Merino, Y., López-Sáez, L., Fierro, E., & Burjachs, F. (2010). Expected trends and surprises in the Lateglacial and Holocene vegetation history of the Iberian Peninsula and Balearic Islands. *Review of Palaeobotany and Palynology*, 162, 458–475.
- Castro, A. J., Martín-Lopez, B., Lopez, E., Plieninger, T., Alcaraz-Segura, D., Vaughn, C. C., & Cabello, J. (2015). Do protected areas networks ensure the supply of ecosystem services? Spatial patterns of two nature reserve systems in semi-arid Spain. *Applied Geography*, 60, 1–9.
- Cerdà, A., Gimenez Morera, A., Burguet, M., Arcenegui, V., Gonzalez Peñaloza, F. A., García-Orenes, F., & Pereira, P. (2012). El impacto del cultivo, el abandono y a intensificación de la agricultura en la pérdida de agua y suelo. El ejemplo de la vertiente norte de la Serra Grossa en el este peninsular. *Cuadernos de Investigación Geográfica*, 38, 75–94. (In Spanish).
- Cerdà, A., & Lasanta, T. (2005). Long-term erosional responses after fire in the Central Spanish Pyrenees 1. Water and sediment yield. *Catena*, 60, 59–80.
- Collantes, F., & Pinilla, D. (2004). Extreme depopulation in the Spanish rural mountain areas: A case study of Aragon in the nineteenth and twentieth centuries. *Rural History*, 15, 149–166. <https://doi.org/10.1017/S0956793304001219>
- Colombo, S., Hanley, N., & Calatrava-Requena, J. (2005). Designing policy for reducing the off-farm effects of soil erosion using choice experiments. *Journal of Agricultural Economics*, 56, 81–95.
- Connell, S. D., & Ghedini, G. (2015). Resisting regime-shifts: The stabilising effect of compensatory processes. *Trends in Ecology & Evolution*, 30, 513–515. <https://doi.org/10.1016/j.tree.2015.06.014>
- Costa, D., Pereira, V., Góis, J., Danko, A., & Fiúza, A. (2017). Understanding public perception of hydraulic fracturing: A case study in Spain. *Journal of Environmental Management*, 205, 551–562. <https://doi.org/10.1016/j.jenvman.2017.09.001>
- Davoudi, S., & Stead, D. (2002). Urban-rural relationships: An introduction and a brief history. *Built Environment*, 28, 269–277.
- Delibes-Mateos, M., Glikman, J. A., Lafuente, R., Villafuerte, R., & Garrido, F. E. (2022). Support to Iberian lynx reintroduction and perceived impacts: Assessments before and after reintroduction. *Conservation Science and Practice*, 4, 3605.
- Doblas-Miranda, E., Rovira, P., Brotons, L., Martínez-Vilalta, J., Retana, J., & Vayreda, J. (2013). Soil carbon stocks and their variability across the forests, shrublands and grasslands of peninsular Spain. *Biogeosciences*, 10, 8353–8361. <https://doi.org/10.5194/bg-10-8353-2013>
- Donohue, I., Leira, M., Hobbs, W., León-Vintró, L., O'Reilly, J., & Irvine, K. (2010). Rapid ecosystem recovery from diffuse pollution after the Great Irish Famine. *Ecological Applications*, 20, 1733–1743.
- Duane, A., Castellnou, M., & Brotons, L. (2021). Towards a comprehensive look at global drivers of novel extreme wildfire events. *Climatic Change*, 165, 43. <https://doi.org/10.1007/s10584-021-03066-4>
- Escribano, G., Calviño, M., Pias, B., Virgós, E., Valladares, F., & Escudero, A. (2014). Diverse guilds provide complementary dispersal services in a woodland expansion process after land abandonment. *Journal of Applied Ecology*, 51, 1701–1711.
- ESPON. (2017). Shrinking rural regions in Europe. In I. Raugze, G. Daly, & M. van Herwijnen (Eds.), *Policy brief*. European Spatial Planning Observation Network.
- Estrada-Carmona, N., Sánchez, A. C., Remans, R., & Jones, S. K. (2022). Complex agricultural landscapes host more biodiversity than simple ones: A global meta-analysis. *Proceedings of the National Academy of Sciences of the United States of America*, 119, e2203385119. <https://doi.org/10.1073/pnas.2203385119>
- Farina, A. (2000). The cultural landscape as a model for the integration of ecology and economics. *BioScience*, 50, 313–320.
- Fischer, J., Lindenmayer, D. B., & Manning, A. D. (2006). Biodiversity, ecosystem function, and resilience: Ten guiding principles for commodity production landscapes. *Frontiers in Ecology and the Environment*, 4, 80–86. [https://doi.org/10.1890/1540-9295\(2006\)004\[0080:BEFART\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2006)004[0080:BEFART]2.0.CO;2)
- Folke, C., Carpenter, S. R., Walker, B., Scheffer, M., Chapin, T., & Rockström, J. (2010). Resilience thinking: Integrating resilience, adaptability and transformability. *Ecology and Society*, 15, 20.
- Franch Auladell, X., Marti-Henneberg, J., & Puig-Farré, J. (2013). Un análisis espacial de las pautas de crecimiento y concentración de la población a partir de series homogéneas: España (1877-2001). *Investigaciones Regionales*, 25, 43–65. (In Spanish).
- Franz, K. W., Romanowski, J., & Saavedra, D. (2011). Effects of prospective landscape changes on species viability in Segre River valley, NE Spain. *Landscape and Urban Planning*, 100, 242–250.
- Fremier, A. K., DeClerck, F. A. J., Bosque-Pérez, N. A., Carmona, N. E., Hill, R., Joyal, T., Keesecker, L., Klos, P. Z., Martínez-Salinas, A., Niemeyer, R., Sanfiorenzo, A., Welsh, K., & Wulffhorst, J. D. (2013).

- Understanding spatiotemporal lags in ecosystem services to improve incentives. *BioScience*, 63, 472–482. <https://doi.org/10.1525/bio.2013.63.6.9>
- Fuchs, R., Schulp, C. J. E., Hengeveld, G. M., Verbug, P. H., Clevers, J. G. P. W., Schelhaas, M.-J., & Herold, M. (2016). Assessing the influence of historic net and gross land changes on the carbon fluxes of Europe. *Global Change Biology*, 22, 2526–2539. <https://doi.org/10.1111/gcb.13191>
- Gaglio, M., Aschoniis, V., Castaldelli, G., & Fano, E. A. (2020). Land use intensification rather than land cover change affects regulating services in the mountainous Adige river basin (Italy). *Ecosystem Services*, 45, 01158. <https://doi.org/10.1016/j.ecoser.2020.101158>
- García-Carmona, M., García-Robles, H., Turpín Torrano, C., Fernández Ondoño, E., Lorite Moreno, J., Sierra Aragón, M., & Martín Peinado, F. J. (2019). Residual pollution and vegetation distribution in amended soils 20 years after a pyrite mine tailings spill (Aznalcóllar, Spain). *The Science of the Total Environment*, 650, 933–940. <https://doi.org/10.1016/j.scitotenv.2018.09.092>
- García-Ruiz, L. (2010). The effects of land uses on soil erosion in Spain: A review. *Catena*, 81, 1–10. <https://doi.org/10.1016/j.catena.2010.01.001>
- Goerlich, F. J., Ruiz, F., Chorén, P., & Albert, C. (2015). *Cambios en la estructura y localización de la población. Una visión de largo plazo (1842-2011)*. Fundación BBVA. 356 pp.
- Gómez-Baggethun, E., Martín-López, B., Lomas, P. L., Zorrilla, P., & Montes, C. (2011). Evolution of ecosystem services in a Mediterranean cultural landscape: Doñana case study, Spain (1956–2006). In A. Sofo (Ed.), *Biodiversity* (pp. 27–46). IntechOpen. <https://doi.org/10.5772/24633>
- Gosselin, F., & Callois, J. M. (2021). On the time lag between human activity and biodiversity in Europe at the national scale. *Anthropocene*, 35, 100303. <https://doi.org/10.1016/j.ancene.2021.100303>
- Guillaume, S., Barcet, H., de Munnik, N., Maire, E., & Marais-Sicre, C. (2020). Evolution of traditional agroforestry landscapes and development of invasive species: Lessons from the Pyrenees (France). *Sustainability Science*, 15, 1285–1299. <https://doi.org/10.1007/s11625-020-00847-1>
- Halada, L., Evans, D., Romao, C., & Petersen, J.-E. (2011). Which habitats of European importance depend on agricultural practices? *Biodiversity and Conservation*, 20, 2365–2378. <https://doi.org/10.1007/s10531-011-9989-z>
- Heal, K., Phin, A., Waldron, S., Flowers, H., Brunau, P., Coupar, A., & Cundill, A. (2020). Wind farm development on peatlands increases fluvial macronutrient loading. *Ambio*, 49, 442–459. <https://doi.org/10.1007/s13280-019-01200-2>
- Henle, K., Alard, D., Clitherow, J., Cobb, P., Firbank, L., Kull, T., McCracken, D., Moritz, R. F. A., Niemelä, J., Rebane, M., Wascher, D., Watt, A., & Young, J. (2008). Identifying and managing the conflicts between agriculture and biodiversity conservation in Europe—A review. *Agriculture Ecosystems & Environment*, 124, 60–71. <https://doi.org/10.1016/j.agee.2007.09.005>
- Hermoso, V., Carvalho, S. B., Giakoumi, S., Goldsborough, D., Katsanevakis, S., Leontiou, S., Markantonatou, V., Rumes, R., Vogiatzakis, I. N., & Yates, K. L. (2022). The EU biodiversity strategy for 2030: Opportunities and challenges on the path towards biodiversity recovery. *Environmental Science and Policy*, 127, 263–271. <https://doi.org/10.1016/j.envsci.2021.10.028>
- Hinojosa, L., Lambin, E. F., Mzoughi, N., & Napoléone, C. (2018). Constraints to farming in the Mediterranean Alps: Reconciling environmental and agricultural policies. *Land Use Policy*, 75, 726–733. <https://doi.org/10.1016/j.landusepol.2017.11.047>
- Hirt, M. R., Barnes, A. D., Gentile, A., Pollock, L. J., Rosenbaum, B., Thuiller, W., Tucker, M. A., & Brose, U. (2021). Environmental and anthropogenic constraints on animal space use drive extinction risk worldwide. *Ecology Letters*, 24, 2576–2585. <https://doi.org/10.1111/ele.13872>
- Hospers, G. A., & Reverda, N. (2015). *Managing population decline in Europe's urban and rural areas*. SpringerBriefs in Population Studies. <https://doi.org/10.1007/978-3-319-12412-4>
- Infante-Amate, J., Villa, I., Jiménez, F., Martínez Martín, M., Martínez López, D., Cunfer, G., & González de Molina, M. (2016). The rise and fall of the cortijo system: Scattered rural settlements and the colonization of land in Spain's Mediterranean mountains since 1581. *Journal of Historical Geography*, 54, 63–75. <https://doi.org/10.1016/j.jhg.2016.09.001>
- IPBES. (2019). *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. (E. S. Brondizio, J. Settele, S. Díaz, & H. T. Ngo (Eds.)). IPBES Secretariat. <https://doi.org/10.5281/zenodo.3831673>
- IPCC. (2021). Summary for policymakers. In V. Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, J. B. R. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, & B. Zho (Eds.), *Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 3–32). Cambridge University Press. <https://doi.org/10.1017/9781009157896.001>
- Kati, V., Hovardas, T., Dieterich, M., Ibsch, P. L., Mihok, B., & Selva, N. (2015). The challenge of implementing the European network of protected areas Natura 2000. *Conservation Biology*, 29, 260–270. <https://doi.org/10.1111/cobi.12366>
- Laborda Soriano, A. A., Cambra Aliaga, A., & Vidal-Sanchez, M. I. (2021). Depopulation in Spain and violation of occupational rights. *Journal of Occupational Science*, 28, 71–80. <https://doi.org/10.1080/14427591.2021.1896331>
- Ladrera, R., Belmar, O., Tomás, R., Prat, N., & Cañedo-Argüelles, M. (2019). Agricultural impacts on streams near nitrate vulnerable zones: A case study in the Ebro basin, Northern Spain. *PLoS One*, 14, e0218582. <https://doi.org/10.1371/journal.pone.0218582>
- Laliberté, E., Wells, J. A., DeClerck, F., Metcalfe, D. J., Catterall, C. P., Quieroz, I., Aubin, C., Bonser, S. P., Ding, Y., Fraterrigo, J. M., McNamara, S., Morgan, J. W., Sánchez Merlos, D., Vesk, P. A., & Mayfield, M. M. (2010). Land-use intensification reduces functional redundancy and response diversity in plant communities. *Ecology Letters*, 13, 76–86. <https://doi.org/10.1111/j.1461-0248.2009.01403.x>
- Larraz, B., & San-Martin, E. (2021). A tale of two dams: The impact of reservoir management on rural depopulation in central Spain. *Water Resources Management*, 35, 4769–4787. <https://doi.org/10.1007/s11269-021-02938-9>
- Lloret, F., Piñol, J., & Castellnou, M. (2009). Wildfires. In J. Woodward (Ed.), *The physical geography of the mediterranean* (pp. 541–558). Oxford University Press.
- Lloret, F., Solé, A., Pino, J., Vayreda, J., Font, X., & Terradas, J. (2009). Patterns of species impoverishment in managed forests of Catalonia (NE Spain). *Journal of Vegetation Science*, 20, 675–685. <https://doi.org/10.1111/j.1654-1103.2009.01059.x>
- Llurdes, J.-C., Sauri, D., & Cerdan, R. (2003). Ten years wasted: The failure of siting waste facilities in central Catalonia, Spain. *Land Use Policy*, 20, 335–342.
- Lopez, I., & Pardo, M. (2018). Tourism versus nature conservation: Reconciliation of common interests and objectives—An analysis through Picos de Europa National Park. *Journal of Mountain Science*, 15, 2505–2516. <https://doi.org/10.1007/s11629-018-4943-0>
- MacDonald, D., Crabtree, J. R., Wiesinger, G., Dax, T., Stamou, N., Fleury, P., Gutierrez Lazpita, J., & Gibon, A. (2000). Agricultural abandonment in mountain areas of Europe: Environmental consequences and policy response. *Journal of Environmental Management*, 59, 47–69.
- Martin, A., Coolsaet, B., Corbera, E., Dawson, J., Fisher, J., Franks, P., Mertz, O., Pascual, U., Rasmussen, L.V., & Ryan, C. (2018). Land

- use intensification: The promise of sustainability and the reality of trade-offs. In K. Schreckenberg, G. Mace, & M. Poudyal (Eds.), *Ecosystem services and poverty alleviation—Trade-offs and governance*. Routledge.
- Martínez-Abraín, A., Jiménez, J., Jiménez, I., Ferrer, X., Llana, L., Ferrer, M., Palomero, G., Ballesteros, F., Galán, P., & Oro, D. (2020). Ecological consequences of human depopulation of rural areas on wildlife: A unifying perspective. *Biological Conservation*, 252, 108060. <https://doi.org/10.1016/j.biocon.2020.108860>
- Martínez-Sastre, R., Ravera, F., González, J. A., López Santiago, C., Bidegain, I., & Munda, G. (2017). Mediterranean landscapes under change: Combining social multicriteria evaluation and the ecosystem services framework for land use planning. *Land Use Policy*, 67, 472–486. <https://doi.org/10.1016/j.landusepol.2017.06.001>
- Marull, J., Tello, E., Wilcox, P. T., Coll, F., Pons, M., Warde, P., Valldeperes, N., & Ollés, A. (2014). Recovering the landscape history behind a Mediterranean edge environment (The Congost Valley, Catalonia, 1854e2005): The importance of agroforestry systems in biological conservation. *Applied Geography*, 54, 1–17. <https://doi.org/10.1016/j.apgeog.2014.06.030>
- Maurer, C., Lutter, L., Martínez-Núñez, C., Pellissier, L., & Abrecht, M. (2022). Different types of semi-natural habitat are required to sustain diverse wild bee communities across agricultural landscapes. *Journal of Applied Ecology*, 59, 2604–2615. <https://doi.org/10.1111/1365-2664.14260>
- Moreno Saiz, J. C., Iriondo Alegría, J. M., Martínez García, F., Martínez Rodríguez, J., & Salazar Mendías, C. (2019). *Atlas y Libro Rojo de la Flora Vasculosa Amenazada de España*. Ministerio para la Transición Ecológica-Sociedad Española de Biología de la Conservación de Plantas (In Spanish).
- Navarro, L. M., Proença, V., Kaplan, J. O., & Pereira, H. M. (2015). Maintaining disturbance-dependent habitats. In H. Pereira & L. Navarro (Eds.), *Rewilding European landscapes* (pp. 143–147). Springer. https://doi.org/10.1007/978-3-319-12039-3_8
- Navarro-Valverde, F., Cejudo-García, E., & Cañete Pérez, J. A. (2021). The lack of attention given by neo-endogenous rural development practice to areas highly affected by depopulation. The case of Andalusia (Spain) in 2015–2020 period. *European Countryside*, 13, 353–367. <https://doi.org/10.2478/euco-2021-0022>
- Navas, A., Machin, J., Beguería, S., Lopez-Vicente, M., & Gaspar, L. (2008). Soil properties and physiographic factors controlling the natural vegetation regrowth in a disturbed catchment of the Central Spanish Pyrenees. *Agroforestry Systems*, 72, 173–185.
- Normand, S., Svenning, J.-C., & Skov, F. (2007). National and European perspectives on climate change sensitivity of the Habitats directive characteristic plant species. *Journal for Nature Conservation*, 15, 41–53.
- Otero, I., Boada, M., Badía, A., Pla, E., Vayreda, J., Sabaté, S., Gracia, C. A., & Peñuelas, J. (2011). Loss of water availability and stream biodiversity under land abandonment and climate change in a Mediterranean catchment (Olzinelles, NE Spain). *Land Use Policy*, 28, 207–218. <https://doi.org/10.1016/j.landusepol.2010.06.002>
- Perez-Barbería, F. J., Gomez, J. A., & Gordon, I. J. (2023). Legislative hurdles to using traditional domestic livestock in rewilding programmes in Europe. *Ambio*, 52, 585–597.
- Peterson, G. D. (2002). Contagious disturbance, ecological memory, and the emergence of landscape pattern. *Ecosystems*, 5, 329–338.
- Pisabarro, A., Pellitero, R., Serrano, E., & Lopez-Moreno, J. I. (2019). Impacts of land abandonment and climate variability on runoff generation and sediment transport in the Pisuerga headwaters (Cantabrian Mountains, Spain). *Geografiska Annaler: Series A, Physical Geography*, 101, 211–224.
- Power, A. G. (2010). Ecosystem services and agriculture: Tradeoffs and synergies. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 365, 2959–2971. <https://doi.org/10.1098/rstb.2010.0143>
- Ramírez-Hernández, A., Micó, E., Marcos-García, M. A., Brustel, H., & Galante, E. (2014). The “dehesa”, a key ecosystem in maintaining the diversity of Mediterranean saproxylic insects (Coleoptera and Diptera: Syrphidae). *Biodiversity and Conservation*, 23, 2069–2086. <https://doi.org/10.1007/s10531-014-0705-7>
- Recio, M. R., Sand, H., & Virgos, E. (2020). Promoting grazing or rewilding initiatives against rural exodus? The return of the wolf and other large carnivores must be considered. *Environmental Conservation*, 47, 269–276.
- Regos, A., Domínguez, J., Gil-Tena, A., Brotons, L., Ninyerola, M., & Pons, X. (2016). Rural abandoned landscapes and bird assemblages: Winners and losers in the rewilding of a marginal mountain area (NW Spain). *Regional Environmental Change*, 16, 199–211. <https://doi.org/10.1007/s10113-014-0740-7>
- Rodríguez-Rodríguez, D., Larrubia, R., & Sinoga, J. W. (2021). Are protected areas good for the human species? Effects of protected areas on rural depopulation in Spain. *Science of the Total Environment*, 763, 144399. <https://doi.org/10.1016/j.scitotenv.2020.144399>
- Root-Bernstein, M., & Ladle, R. (2018). Ecology of a widespread large omnivore, *Homo sapiens*, and its impacts on ecosystem processes. *Ecology and Evolution*, 9, 10874–10894. <https://doi.org/10.1002/ece3.5049>
- Ropero, R., Runi, R., & Aguilera, P. A. (2019). Modelling relationships between socioeconomy, landscape and water flows in Mediterranean agroecosystems: A case study in Adra catchment (Spain) using Bayesian networks. *Environmental and Ecological Statistics*, 26, 47–86.
- Rozas, V., Zas, R., & Solla, A. (2009). Spatial structure of deciduous forest stands with contrasting human influence in northwest Spain. *European Journal of Forest Research*, 209(128), 273–285. <https://doi.org/10.1007/s10342-009-0263-9>
- Sabatini, F. M., Burrascano, S., Keeton, W. S., Levers, C., Lindner, M., Pötzschner, F., Veker, P. J., Bauhus, J., Buchwald, E., Chaskovsky, O., Debaive, N., Horváth, N., Garbarino, M., Grigoriadis, N., Lombardi, F., Duarte, I. M., Meyer, I. M., Midteng, R., Mikac, S., ... Kuemmerle, T. (2017). Where are Europe's last primary forests? *Diversity and Distributions*, 24, 1426–1439. <https://doi.org/10.1111/ddi.12778>
- Sanjuan, Y., Gómez-Villa, A., Nadal-Romero, E., Álvarez-Martínez, J., Arnáez, J., Serrano-Muela, M. P., Rubiales, J. M., González-Sampériz, P., & García-Ruiz, J. M. (2016). Linking land cover changes in the sub-alpine and montane belts to changes in a torrential river. *Land Degradation and Development*, 27, 179–189. <https://doi.org/10.1002/ldr.2294>
- Schou, J. S., Bladt, J., Ejrnaes, R., Thomsen, M. N., Vedel, S. E., & Flojgaard, C. (2021). Economic assessment of rewilding versus agri-environmental nature management. *Ambio*, 50, 1047–1057. <https://doi.org/10.1007/s13280-020-01423-8>
- Silvestre, J., & Clar, E. (2010). The demographic impact of irrigation projects: A comparison of two case studies of the Ebro basin, Spain, 1900–2001. *Journal of Historical Geography*, 36, 31–326.
- Svenning, J.-C., Munk, M., & Schweiger, A. (2019). Trophic rewilding: Ecological restoration of top-down trophic interactions to promote self-regulating biodiverse ecosystems. In N. Pettorelli, S. Durant, & J. Du Toit (Eds.), *Rewilding* (pp. 73–98). Cambridge University Press. <https://doi.org/10.1017/9781108560962.005>
- Tacoli, C. (1998). Rural-urban interactions: A guide to the literature. *Environment and Urbanization*, 10, 147–166.
- Talle, M., Deak, B., Poschod, P., Valko, O., Westerbeg, L., & Milberg, P. (2016). Grazing vs. mowing: A meta-analysis of biodiversity benefits for grassland management. *Agriculture, Ecosystems & Environment*, 222, 200–212. <https://doi.org/10.1016/j.agee.2016.02.008>
- Traba, J., & Morales, M. B. (2019). The decline of farmland birds in Spain is strongly associated to the loss of fallowland. *Scientific Reports*, 9, 9473. <https://doi.org/10.1038/s41598-019-45854-0>

- Underwood, E. C., Viers, J. H., Klausmeyer, K. R., Cox, R. L., & Shaw, M. R. (2009). Threats and biodiversity in the Mediterranean biome. *Diversity and Distributions*, 15, 188–197. <https://doi.org/10.1111/j.1472-4642.2008.00518.x>
- Varga, D. (2020). Are agrarian areas in Mediterranean mountain regions becoming extinct? A methodological approach to their conservation. *Forests*, 11, 1116. <https://doi.org/10.3390/f11011116>
- Vayreda, J., Gracia, M., Canadell, J. G., & Retana, J. (2012). Spatial patterns and predictors of forest carbon stocks in Western Mediterranean. *Ecosystems*, 15, 1258–1270. <https://doi.org/10.1007/s10021-012-9582-7>
- Velazquez, E., Martínez-Jaraíz, C., Wheeler, C., Mitchard, E. T. A., & Bravo, F. (2022). Forest expansion in abandoned agricultural lands has limited effect to offset carbon emissions from Central-North Spain. *Regional Environmental Change*, 22, 132.
- Vidal-Legaz, B., Martínez-Fernández, J., Sanchez Picon, A., & Pugnaire, F. I. (2013). Trade-offs between maintenance of ecosystem services and socio-economic development in rural mountainous communities in southern Spain: A dynamic simulation approach. *Journal of Environmental Management*, 131, 280–297.
- Villarreal Walker, R., & Beck, M. B. (2012). Understanding the metabolism of urban-rural ecosystems. A multi-sectoral systems analysis. *Urban Ecosystems*, 12, 809–848. <https://doi.org/10.1007/s11252-012-0241-8>
- Walker, B., Holling, C. S., Carpenter, S. R., & Kinzig, A. (2004). Resilience adaptability and transformability in social-ecological systems. *Ecology and Society*, 9, art.5. <http://www.ecologyandsociety.org/vol9/iss2/art5/>
- ## DATA SOURCES
- Arroyo, F. M., Lopez, H. S., Castillo-Valero, J. S., & Garcia-Cortijo, M. C. (2022). Rural development programmes: Lessons learnt, and knowledge advancement. A case study in Castilla-La Mancha (Spain). *Agricultural Economics*, 68, 393–402. https://agricecon.agriculturejournals.cz/artkey/age-202210-0004_rural-development-programmes-lessons-learnt-and-knowledge-advancement-a-case-study-in-castilla-la-mancha-sp.php
- Barroso, V. M., de Castro-Pardo, M., Martínez, P. F., & Azevedo, J. C. (2022). A regionalized IO-model to value seasonal recreational ecosystem services in a mountain National Park in Spain. *Journal of Mountain Science*, 19, 211–227. <https://link.springer.com/article/10.1007/s11629-021-6911-3>
- Bautista, S., Orr, B. J., Alloza, J. A., & Vallejo, R. V. (2010b). Evaluating the restoration of dryland ecosystems in the Northern Mediterranean. In G. Schneier-Madanes & M. F. Courel (Eds.), *Water and sustainability in arid regions* (pp. 295–310). Springer.
- Beguería, S. (2006). Changes in land cover and shallow landslide activity: A case study in the Spanish Pyrenees. *Geomorphology*, 74, 196–206. <https://www.sciencedirect.com/science/article/pii/S0169555X05002692>
- Beguería, S., Lopez-Moreno, J. I., Gomez-Villar, A., Rubio, V., Lana-Renault, N., & Garcia-Ruiz, J. M. (2006). Fluvial adjustments to soil erosion and plant cover changes in the Central Spanish Pyrenees. *Geografiska Annaler: Series A, Physical Geography*, 88(A), 177–186. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1468-0459.2006.00293.x>
- Beguería, S., López-Moreno, J. I., Lorente, A., Seeger, M., & García-Ruiz, J. M. (2003b). Assessing the effects of climate oscillations and land-use changes on streamflow in the Central Spanish Pyrenees. *Ambio*, 32, 283–286. <https://bio-one.org/journals/AMBIO-A-Journal-of-the-Human-Environment/volume-32/issue-4/0044-7447-32.4.283/Assessing-the-Effect-of-Climate-Oscillations-and-Land-use-Changes/10.1579/0044-7447-32.4.283.short>
- Bodi, M. B., Cerda, A., Mataix-Solera, J., & Doerr, S. H. (2012). A review of fire effects on vegetation and soil in the Mediterranean Basin. *Boletín Asociación Geógrafos Españoles*, 58, 33–55. (in Spanish).
- Bruno, D., Sorando, R., Farziro, B. A., Castellano, C., Céspedes, V., Gallardo, B., Jimenez, J. J., Lopez, M. V., Flores, R. L., Moret-Fernandez, D., Navarro, E., Picazo, F., Sevilla-Callejo, M., Tormo, J., Macus, J. J. V., Nicolou, J. M., & Comin, F. A. (2021b). Depopulation impacts on ecosystem services in Mediterranean rural areas. *Ecosystem Services*, 52, 101039. <https://www.sciencedirect.com/science/article/pii/S2212041621001273>
- Bryant, R. L., Paniagua, A., & Kizos, T. (2011). Conceptualizing 'shadow landscape' in political ecology and rural studies. *Land Use Policy*, 28, 460–471. <https://www.sciencedirect.com/science/article/abs/pii/S0264837710000967>
- Caceres-Feria, R., Hernandez-Ramirez, M., & Ruiz-Ballesteros, E. (2021b). Depopulation, community-based tourism, and community resilience in south-west Spain. *Journal of Rural Studies*, 88, 108–116. <https://www.sciencedirect.com/science/article/pii/S0743016721003144>
- Calafat-Marzal, C., Gallego-Salguero, A., Segura, M., & Calvet-Sanz, S. (2021). GIS-based and outranking approach to assess suitable pig farming areas in the Mediterranean region: Valencian community. *Animals*, 11, 1151. <https://www.mdpi.com/2076-2615/11/4/1151>
- Camarero, J. J., de Andres, E. G., Colangelo, M., & Loren, C. D. (2022). Growth history of pollarded black poplars in a continental Mediterranean region: A paradigm of vanishing landscapes. *Forest Ecology and Management*, 517, 120268. <https://www.sciencedirect.com/science/article/pii/S0378112722002626>
- Camarero, L., & Oliva, J. (2019). Thinking in rural gap: Mobility and social inequalities. *Palgrave Communications*, 5, 95. <https://www.nature.com/articles/s41599-019-0306-x>
- Carmona, C. P., Guerrero, I., Peco, B., Morales, M. B., Oñate, J. J., Pärt, T., Tschardtke, T., Lira, J., Avik, T., Emmerson, M., Berendse, F., Ceryngier, P., Bretagnolle, V., Weisser, W. W., & Bengtsson, J. (2020b). Agriculture intensification reduces plant taxonomic and functional diversity across European arable systems. *Functional Ecology*, 34, 1448–1460. <https://besjournals.onlinelibrary.wiley.com/action/showCitFormats?doi=10.1111%2F1365-2435.13608>
- Candel-Perez, D., Hernandez-Alonso, H., Castro, F., Sanguesa-Barreda, G., Mutke, S., Garcia-Hidalgo, M., Rozas, V., & Olano, J. M. (2022). 250-year reconstruction of pollarding events reveals sharp management changes in Iberian ash woodlands. *Trees-Structure and Function*, 36, 1909–1921. <https://link.springer.com/article/10.1007/s00468-022-02343-8>
- Castro, A. J., Martín-Lopez, B., Lopez, E., Plieninger, T., Alcaraz-Segura, D., Vaughn, C. C., & Cabello, J. (2015b). Do protected areas networks ensure the supply of ecosystem services? Spatial patterns of two nature reserve systems in semi-arid Spain. *Applied Geography*, 60, 1–9. <https://www.sciencedirect.com/science/article/abs/pii/S0143622815000430>
- Cerdá, A., & Lasanta, T. (2005b). Long-term erosional responses after fire in the Central Spanish Pyrenees: 1. Water and sediment yield. *Catena*, 60, 59–80. <https://www.sciencedirect.com/science/article/abs/pii/S0341816204001171>
- Cerdá, A., Giménez Morera, A., Burguet, M., Arcenegui, V., Gonzalez Peñaloza, F. A., García-Orenes, F., & Pereira, P. (2012b). El impacto del cultivo, el abandono y a intensificación de la agricultura en la pérdida de agua y suelo. El ejemplo de la vertiente norte de la Serra Grossa en el este peninsular. *Cuadernos de Investigación Geográfica*, 38, 75–94. (in Spanish).
- Colombo, S., Hanley, N., & Calatrava-Requena, J. (2005b). Designing policy for reducing the off-farm effects of soil erosion using choice experiments. *Journal of Agricultural Economics*, 56, 81–95. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1477-9552.2005.tb00123.x>
- de Molina, M. G. (2012). Environmental arguments for the renovation of the Agrarian History. *Vínculos de Historia*, 1, 95–114. (in Spanish).
- Delibes-Mateos, M., Glikman, J. A., Lafuente, R., Villafuerte, R., & Garrido, F. E. (2022b). Support to Iberian lynx reintroduction and perceived impacts: Assessments before and after reintroduction. *Conservation Science and Practice*, 4, 3605. <https://conbio.onlinelibrary.wiley.com/doi/full/10.1111/csp2.605>
- Debussche, M., Lepart, J., & Dervieux, A. (1999). Mediterranean landscape changes: Evidence from old postcards. *Global Ecology and Biogeography*, 8, 3–15. <https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2699.1999.00316.x>
- Moreno Díaz del Campo, F. J. (2021). The Guadiana River and the Real Dehesa de Zacatena at the end of the ancient regime. Drainage projects, environmental changes and institutional disputes. *Estudios Geográficos*, 82, e072 (in Spanish). <https://estudiosgeograficos.revistas.csic.es/index.php/estudiosgeograficos/article/view/1060>
- del Río, S., Alvarez-Esteban, R., Alonso-Redondo, R., Hidalgo, C., & Penas, A. (2021). A new integrated methodology for characterizing and assessing suitable areas for viticulture: A case study in Northwest Spain. *European Journal of Agronomy*, 131, 126391. <https://www.sciencedirect.com/science/article/pii/S1161030121001623>
- González Díaz, J. A., Celaya, R., García, F. F., Osoro, K., & García, R. R. (2019). Dynamics of rural landscapes in marginal areas of northern Spain: Past, present, and future. *Land Degradation & Development*, 30(1), 41–150. <https://onlinelibrary.wiley.com/doi/abs/10.1002/ldr.3201>
- González Díaz, J. A., Díaz, B. G., & Rosa-García, R. (2022). Role of socioeconomy and land management in the evolution of agrosilvopastoral landscapes

- in Northern Spain: The case study of Redes Biosphere Reserve. *Frontiers in Ecology and Evolution*, 10, 949093. <https://doi.org/10.3389/fevo.2022.949093>
- Doblas-Miranda, E., Rovira, P., Brotons, L., Martínez-Vilalta, J., Retana, J., & Vayreda, J. (2013b). Soil carbon stocks and their variability across the forests, shrublands and grasslands of peninsular Spain. *Biogeosciences*, 10, 8353–8361. <https://bg.copernicus.org/articles/10/8353/2013/>
- Donohue, I., Leira, M., Hobbs, W., Leon-Vintro, L., O'Reilly, J., & Irvine, K. (2010b). Rapid ecosystem recovery from diffuse pollution after the Great Irish Famine. *Ecological Applications*, 20, 1733–1743. <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/09-1966.1>
- Errea, M. P., Cortijos-Lopez, M., Llena, M., Nadal-Romero, E., Zabalza-Martinez, J., & Lasanta, T. (2023). From the local landscape organization to land abandonment: An analysis of landscape changes (1956–2017) in the Aisa Valley (Spanish Pyrenees). *Landscape Ecology*, 36, 3443–3462. <https://link.springer.com/article/10.1007/s10980-023-01675-1>
- Escribano-Avila, G., Calvino-Cancela, M., Pias, B., Virgos, E., Valladares, F., & Escudero, A. (2014). Diverse guilds provide complementary dispersal services in a woodland expansion process after land abandonment. *Journal of Applied Ecology*, 51, 1701–1711. <https://besjournals.onlinelibrary.wiley.com/doi/full/10.1111/1365-2664.12340>
- Franz, K. W., Romanowski, J., & Saavedra, D. (2011b). Effects of prospective landscape changes on species viability in Segre River valley, NE Spain. *Landscape and Urban Planning*, 100, 242–250. <https://www.sciencedirect.com/science/article/abs/pii/S016920461100020X>
- Frigole, J. (2012). Cosmologies, heritagization and eco-symbols in the Catalan Pyrenees within a global perspective. *Revista de Antropologia Social*, 21, 173–196. <https://www.redalyc.org/pdf/838/83824463007.pdf>
- García-Carmona, M., García-Robles, H., Turpín Torrano, C., Fernández Ondoño, E., Lorite Moreno, J., Sierra Aragón, M., & Martín Peinado, F. J. (2019b). Residual pollution and vegetation distribution in amended soils 20 years after a pyrite mine tailings spill (Aznalcóllar, Spain). *Science of the Total Environment*, 650, 933–940. <https://www.sciencedirect.com/science/article/abs/pii/S0048969718335344>
- García-Ruiz, J. M., Lana-Renault, N., Begueria, S., Lasanta, T., Regues, D., Nadal-Romero, E., Serrano-Muela, P., Lopez-Moreno, J. I., Alvera, B., Martí-Bono, C., & Alatorre, L. C. (2010). From plot to regional scales: Interactions of slope and catchment hydrological and geomorphic processes in the Spanish Pyrenees. *Geomorphology*, 120, 248–257. <https://www.sciencedirect.com/science/article/abs/pii/S0169555X10001571>
- García-Ruiz, J. M., Lasanta, T., Nadal-Romero, E., Lana-Renault, N., & Alvarez-Farizo, B. (2020). Rewilding and restoring cultural landscapes in Mediterranean mountains: Opportunities and challenges. *Land Use Policy*, 99, 104850. <https://www.sciencedirect.com/science/article/abs/pii/S0264837719313110>
- Garha, N. S. (2022). From decarbonization to depopulation: An emerging challenge for the carbon-intensive regions under the energy transition in Spain. *Sustainability*, 14, 14786. <https://www.mdpi.com/2071-1050/14/22/14786>
- Gonzalez-Rosado, M., Parras-Alcantara, L., Aguilera-Huertas, J., & Lozano-Garcia, B. (2021). Building an agroecological process towards agricultural sustainability: A case study from Southern Spain. *Agriculture*, 11, 1024. <https://www.mdpi.com/2077-0472/11/10/1024>
- Gracia, A., Sanchez, A. M., Jurado, F., & Mallor, C. (2020). Making use of sustainable local plant genetic resources: Would consumers support the recovery of a traditional purple carrot? *Sustainability*, 12, 6549. <https://www.mdpi.com/2071-1050/12/16/6549>
- Henle, K., Alard, D., Clitherow, J., Cobb, P., Firbank, L., Kull, T., McCracken, D., Moritz, R. F. A., Niemelä, J., Rebane, M., Wascher, D., Watt, A., & Young, J. (2008b). Identifying and managing the conflicts between agriculture and biodiversity conservation in Europe—A review. *Agriculture Ecosystems & Environment*, 124, 60–71. <https://www.sciencedirect.com/science/article/abs/pii/S0167880907002319>
- Ladrera, R., Belmar, O., Tomás, R., Prat, N., & Cañedo-Argüelles, M. (2019b). Agricultural impacts on streams near nitrate vulnerable zones: A case study in the Ebro basin, Northern Spain. *PLoS One*, 14, e0218582. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0218582>
- Larraz, B., & San-Martin, E. (2021b). A tale of two dams: The impact of reservoir management on rural depopulation in Central Spain. *Water Resources Management*, 35, 4769–4787. <https://link.springer.com/article/10.1007/s11269-021-02938-9>
- Lasanta, T., Arnaez, J., Pascual, N., Ruiz-Flano, P., Errea, M. P., & Lana-Renault, N. (2017). Space-time process and drivers of land abandonment in Europe. *Catena*, 149, 810–823. <https://www.sciencedirect.com/science/article/abs/pii/S0341816216300777>
- Llurdes, J. C., Sauri, D., & Cerdan, R. (2003b). Ten years wasted: The failure of siting waste facilities in central Catalonia, Spain. *Land Use Policy*, 20, 335–342. <https://www.sciencedirect.com/science/article/abs/pii/S0264837703000577>
- Lopez, I., & Pardo, M. (2018b). Tourism versus nature conservation: Reconciliation of common interests and objectives an analysis through Picos de Europa National Park. *Journal of Mountain Science*, 15, 2505–2516. <https://link.springer.com/article/10.1007/s11629-018-4943-0>
- Luque Revuelto, R. M. (2012). The recent demographic dynamics in the Sierra Morena of Cordoba. *Revista de Estudios Regionales*, 95, 121–147. (in Spanish). <https://www.redalyc.org/pdf/755/75525394004.pdf>
- MacDonald, D., Crabtree, J. R., Wiesinger, G., Dax, T., Stamou, N., Fleury, P., Lazpita, J. G., & Gibon, A. (2000b). Agricultural abandonment in mountain areas of Europe: Environmental consequences and policy response. *Journal of Environmental Management*, 59, 47–69. <https://www.sciencedirect.com/science/article/abs/pii/S030147979903353>
- Martinez-Abraín, A., Ferrer, X., Jimenez, J., & Fernandez-Calvo, I. C. (2021). The selection of anthropogenic habitat by wildlife as an ecological consequence of rural exodus: Empirical examples from Spain. *Animal Biodiversity and Conservation*, 44, 195–203. <https://ruc.udc.es/dspace/handle/2183/28512>
- Martinez, P. F., de Castro-Pardo, M., Barroso, V. M., & Azevedo, J. C. (2020). Assessing sustainable rural development based on ecosystem services vulnerability. *Land*, 9(7), 222. <https://www.mdpi.com/2073-445X/9/7/222>
- Morales-Molino, C., Tinner, W., Perea, R., Carrion, J. S., Colombaroli, D., Valbuena-Carabana, M., Zafra, E., & Gil, L. (2019). Unprecedented herbivory threatens rear-edge populations of Betula in southwestern Eurasia. *Ecology*, 100, e02833. <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecy.2833>
- Mugica, L., Canals, R. M., & San Emeterio, L. (2018). Changes in soil nitrogen dynamics caused by prescribed fires in dense gorse lands in SW Pyrenees. *Science of the Total Environment*, 639, 175–185. <https://www.sciencedirect.com/science/article/abs/pii/S0048969718317807>
- Navas, A., Machin, J., Begueria, S., Lopez-Vicente, M., & Gaspar, L. (2008b). Soil properties and physiographic factors controlling the natural vegetation regrowth in a disturbed catchment of the Central Spanish Pyrenees. *Agroforestry Systems*, 72, 173–185. <https://link.springer.com/article/10.1007/s10457-007-9085-2>
- Otero, I., Boada, M., Badia, A., Pla, E., Vayreda, J., Sabate, S., Gracia, C. A., & Peñuelas, J. (2011b). Loss of water availability and stream biodiversity under land abandonment and climate change in a Mediterranean catchment (Olzinelles, NE Spain). *Land Use Policy*, 28, 207–218. <https://www.sciencedirect.com/science/article/abs/pii/S0264837710000645>
- Palacio, S. (2021). From depopulation to rural repopulation in mountain areas. *Ecosistemas: Revista Científica y Técnica de Ecología y Medio Ambiente*, 30, 2164. <https://revistaecosistemas.net/index.php/ecosistemas/article/view/2164>
- Pareja, J., Baraza, E., Ibanez, M., Domenech, O., & Bartolome, J. (2020). The role of feral goats in maintaining firebreaks by using attractants. *Sustainability*, 12, 7144. <https://www.mdpi.com/2071-1050/12/17/7144>
- Pausas, J. G., & Fernandez-Munoz, S. (2012). Fire regime changes in the Western Mediterranean Basin: From fuel-limited to drought-driven fire regime. *Climatic Change*, 110, 215–226. <https://link.springer.com/article/10.1007/s10584-011-0060-6>
- Pereira, L. S., Louro, V., Rosário, L. D., & Almeida, A. (2006). Desertification, territory and people, a holistic approach in the Portuguese context. In W. G. Kepner, J. L. Rubio, D. A. Mouat, & F. Pedrazzini (Eds.), *Desertification in the Mediterranean Region. A Security Issue*. NATO Security Through Science Series (Vol. 3, pp. 269–289). Springer. https://link.springer.com/chapter/10.1007/1-4020-3760-0_11
- Perez-Barberia, F. J., Gomez, J. A., & Gordon, I. J. (2023b). Legislative hurdles to using traditional domestic livestock in rewilding programmes in Europe. *Ambio*, 52, 585–597. <https://link.springer.com/article/10.1007/s13280-022-01822-z>
- Perez-Granados, C., Serrano-Davies, E., & Nogueiras, V. (2018). Returning home after fire: How fire may help us manage the persistence of scrub-steppe specialist bird populations. *Biodiversity and Conservation*, 27, 3087–3102. <https://link.springer.com/article/10.1007/s10531-018-1586-y>
- Pisabarro, A., Pellitero, R., Serrano, E., & Lopez-Moreno, J. I. (2019b). Impacts of land abandonment and climate variability on runoff generation and sediment transport in the Pisuerga headwaters (Cantabrian Mountains, Spain). *Geografiska Annaler: Series A, Physical Geography*, 101, 211–224. <https://www.tandfonline.com/doi/abs/10.1080/04353676.2019.1591042>
- Plieninger, T. (2006). Habitat loss, fragmentation, and alteration—Quantifying the impact of land-use changes on a Spanish dehesa landscape by use of aerial

- photography and GIS. *Landscape Ecology*, 21, 91–105. <https://link.springer.com/article/10.1007/s10980-005-8294-1>
- Recio, M. R., Sand, H., & Virgos, E. (2020b). Promoting grazing or rewilding initiatives against rural exodus? The return of the wolf and other large carnivores must be considered. *Environmental Conservation*, 47, 269–276. <https://doi.org/10.1017/S0376892920000284>
- Regos, A., Domínguez, J., Gil-Tena, A., Brotons, L., Ninyerola, M., & Pons, X. (2016b). Rural abandoned landscapes and bird assemblages: Winners and losers in the rewilding of a marginal mountain area (NW Spain). *Regional Environmental Change*, 16, 199–211. <https://link.springer.com/article/10.1007/s10113-014-0740-7>
- Riera, S., Wansard, G., & Julià, R. (2004). 2000-year environmental history of a karstic lake in the Mediterranean pre-pyrenees: The Estanya lakes (Spain). *Catena*, 55, 293–324. <https://www.sciencedirect.com/science/article/abs/pii/S0341816203001073>
- Rodríguez-Rodríguez, D., Larrubia, R., & Sinoga, J. D. (2021). Are protected areas good for the human species? Effects of protected areas on rural depopulation in Spain. *Science of the Total Environment*, 763, 144399. <https://www.sciencedirect.com/science/article/abs/pii/S0048969720379304>
- Rodríguez-Rodríguez, D., & Vargas, R. L. (2022). Protected areas and rural depopulation in Spain: A multi-stakeholder perceptual study. *Land*, 11, 384. <https://www.mdpi.com/2073-445X/11/3/384>
- Romita, T., & Morales, S. N. (2014). New rural population: Rural users, nomadic and new inhabitants. *Gran Tour: Revista de Investigaciones Turísticas*, 10, 4–22. (in Spanish).
- Ropero, R., Runi, R., & Aguilera, P. A. (2019b). Modelling relationships between socioeconomy, landscape and water flows in Mediterranean agroecosystems: A case study in Adra catchment (Spain) using Bayesian networks. *Environmental and Ecological Statistics*, 26, 47–86. <https://link.springer.com/article/10.1007/s10651-019-00419-2>
- Rubio-Cuadrado, A., Camarero, J. J., Gordaliza, G. G., Cerioni, M., Montes, F., & Gil, L. (2020). Competition overrides climate as trigger of growth decline in a mixed Fagaceae Mediterranean rear-edge forest. *Annals of Forest Science*, 77, 94. <https://link.springer.com/article/10.1007/s13595-020-01004-5>
- San Emeterio, L., Mugica, L., Ugarte, M. D., Goicoa, T., & Canals, R. M. (2016). Sustainability of traditional pastoral fires in highlands under global change: Effects on soil function and nutrient cycling. *Agriculture Ecosystems & Environment*, 235, 155–163. <https://www.sciencedirect.com/science/article/abs/pii/S0167880916305011>
- Sanjuan, Y., Arnaez, J., Begueria, S., Lana-Renault, N., Lasanta, T., Gomez-Villar, A., Alvarez-Martinez, J., Coba-Perez, P., & Garcia-Ruiz, J. M. (2018). Woody plant encroachment following grazing abandonment in the subalpine belt: A case study in northern Spain. *Regional Environmental Change*, 18, 1103–1115. <https://link.springer.com/article/10.1007/s10113-017-1245-y>
- Sanjuan, Y., Gomez-Villar, A., Nadal-Romero, E., Alvarez-Martinez, J., Arnaez, J., Serrano-Muela, M. P., Rubiales, J. M., Gonzalez-Samperiz, P., & Garcia-Ruiz, J. M. (2016b). Linking land cover changes in the sub-alpine and montane belts to changes in a Torrential River. *Land Degradation & Development*, 27, 179–189. <https://onlinelibrary.wiley.com/doi/abs/10.1002/ldr.2294>
- Schaich, H., Kizos, T., Schneider, S., & Plieninger, T. (2015). Land change in eastern Mediterranean wood-pasture landscapes: The case of deciduous oak woodlands in Lesvos (Greece). *Environmental Management*, 56, 110–126. <https://link.springer.com/article/10.1007/s00267-015-0496-y>
- Seeger, M., & Ries, J. B. (2008). Soil degradation and soil surface process intensities on abandoned fields in Mediterranean mountain environments. *Land Degradation & Development*, 19, 488–501. <https://onlinelibrary.wiley.com/doi/abs/10.1002/ldr.854>
- Shakesby, R. A. (2011). Post-wildfire soil erosion in the Mediterranean: Review and future research directions. *Earth Science Reviews*, 105, 71–100. <https://www.sciencedirect.com/science/article/abs/pii/S001282521100002X>
- Silvestre, J., & Clar, E. (2010b). The demographic impact of irrigation projects: A comparison of two case studies of the Ebro basin, Spain, 1900–2001. *Journal of Historical Geography*, 36, 315–326. <https://www.sciencedirect.com/science/article/abs/pii/S0305748809001145>
- Solomun, M. K., Barger, N., Cerda, A., Keesstra, S., & Markovic, M. (2018). Assessing land condition as a first step to achieving land degradation neutrality: A case study of the Republic of Srpska. *Environmental Science & Policy*, 90, 19–27. <https://www.sciencedirect.com/science/article/abs/pii/S1462901118306580>
- Tenza-Peral, A., Perez-Ibarra, I., Breceda, A., Martínez-Fernández, J., & Gimenez, A. (2022). Can local policy options reverse the decline process of small and marginalized rural areas influenced by global change? *Environmental Science & Policy*, 127, 57–65. <https://www.sciencedirect.com/science/article/abs/pii/S1462901121002872>
- Underwood, E. C., Viers, J. H., Klausmeyer, K. R., Cox, R. L., & Shaw, M. R. (2009b). Threats and biodiversity in the Mediterranean biome. *Diversity and Distribution*, 15, 188–197. <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1472-4642.2008.00518.x>
- Urbano, B., Barquero, M., & Gonzalez-Andres, F. (2022). The environmental impact of fresh tomatoes consumed in cities: A comparative LCA of long-distance transportation and local production. *Scientia Horticulturae*, 301, 111126. <https://www.sciencedirect.com/science/article/pii/S0304423822002473>
- Valenzuela, V. G., & Holl, A. (2023). Growth and decline in rural Spain: An exploratory analysis. *European Planning Studies*, 32(2), 430–453. <https://doi.org/10.1080/09654313.2023.2179390>
- Varga, D. (2020b). Are agrarian areas in Mediterranean Mountain regions becoming extinct? A methodological approach to their conservation. *Forests*, 11, 1116. <https://www.mdpi.com/1999-4907/11/10/1116>
- Vayreda, J., Gracia, M., Canadell, J. G., & Retana, J. (2012b). Spatial patterns and predictors of forest carbon stocks in Western Mediterranean. *Ecosystems*, 15, 1258–1270. <https://link.springer.com/article/10.1007/s10021-012-9582-7>
- Vazquez-Hisado, J. C. (2018). The process towards sustainable mobility in Seville: Historic town center, pedestrians and cyclists. *Ciudad y Territorio. Estudios Territoriales*, 50, 263–276.
- Velazquez, E., Martínez-Jaraíz, C., Wheeler, C., Mitchard, E. T. A., & Bravo, F. (2022b). Forest expansion in abandoned agricultural lands has limited effect to offset carbon emissions from Central-North Spain. *Regional Environmental Change*, 22, 132. <https://link.springer.com/article/10.1007/s10113-022-01978-0>
- Vicente-Serrano, S. M., Lasanta, T., & Romo, A. (2004). Analysis of spatial and temporal evolution of vegetation cover in the Spanish central pyrenees: Role of human management. *Environmental Management*, 34, 802–818. <https://link.springer.com/article/10.1007/s00267-003-0022-5>
- Vidal-Legaz, B., Martínez-Fernández, J., Sánchez Picon, A., & Pugnaire, F. I. (2013b). Trade-offs between maintenance of ecosystem services and socio-economic development in rural mountainous communities in southern Spain: A dynamic simulation approach. *Journal of Environmental Management*, 131, 280–297. <https://www.sciencedirect.com/science/article/abs/pii/S0301479713006361>
- Zorrilla-Miras, P., Suarez, M., Gonzalez-Garcia, A., Gomez, C. S., & Bea, M. (2020). Water supply from Sierra del Rincón Biosphere Reserve to the Community of Madrid (Spain): An ecosystem services approach. *Ecosistemas: Revista Científica y Técnica de Ecología y Medio Ambiente*, 29, 1938. <https://www.revistaecosistemas.net/index.php/ecosistemas/article/view/1938>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Data S1. Supplementary material.

How to cite this article: Lloret, F., Escudero, A., Lloret, J., & Valladares, F. (2024). An ecological perspective for analysing rural depopulation and abandonment. *People and Nature*, 00, 1–17. <https://doi.org/10.1002/pan3.10606>